PROVIDER BULLETIN



Plantation, FL 33324 www.AetnaBetterHealth.com/Florida

Date:	April 30, 2021
Purpose:	Provider Bulletin: Educate providers on
	proper coding for tobaccouse
	prevention and cessation
Subject:	Tobacco Cessation Billing Update
Products:	MMA, LTC, FHK
From:	<u>Provider Relations</u>

Dear Provider,

Aetna Better Health of Florida (ABHFL) would like to remind you of the billing and coding guidelines for CPT codes 99406 and 99407. These two codes are used to report counseling, by a physician or non-physician practitioner, for smoking and tobacco use cessation counseling.

Code	Description
99406	Smoking and tobacco use cessation counseling visit; intermediate, greater than 3 minutes up to 10 minutes
99407	Smoking and tobacco use cessation counseling visit; intermediate, greater than 10 minutes

OBs and Midwives billing 99406/99407

- These codes are for actual counseling, not just screening for tobacco use
- The claims must include a diagnosis from the charts below
- Modifier 25 should be appended to the E/M code to indicate separate services

PCPs billing 99406/99407

- Routine office visit with counseling Modifier 25 should be appended to the E/M code to indicate separate services
- Preventative visit with counseling 99406 and 99407 are <u>not</u> payable. Counseling is considered part of the preventative visit.
 - Note: Providers may bill a preventative visit and an E/M visit together using Modifier 25 on the E/M code if there is a specific problem/issue that they addressed during the preventative visit, however, the counseling is not payable.



Diagnosis Codes allowed for 99406/99407:

When billing for these services providers must use an ICD-10 F17 code or a Z code. The F codes are used if the patient is dependent on tobacco. The Z codes are used if there is <u>not</u> dependence on tobacco. The Z codes cannot be combined with an F17 code.

Aetna Better Health of Florida also reimburses for tobacco cessation counseling provided to pregnant women. The codes specific to maternal and newborn health (O99) and the toxic effects of tobacco (T65) are used by medical providers to reflect the biological impact of tobacco use.

Please see below a list of Dx codes appropriate when billing CPT code 99406 or 99407.

FCODES

ICD-10	Description: All with Nicotine Dependence
Diagnosis Code	
F17.200*	Product unspecified, uncomplicated
F17.201*	Product unspecified, in remission
F17.203	Product unspecified, with withdrawal
F17.208	Product unspecified, with other nicotine-induced disorders
F17.209	Product unspecified, with unspecified nicotine-induced disorders
F17.210*	Cigarettes, uncomplicated
F17.211*	Cigarettes, in remission
F17.213	Cigarettes, with withdrawal
F17.218	Cigarettes, with other nicotine-induced disorders
F17.219	Cigarettes, with unspecified nicotine-induced disorders
F17.220*	Chewing tobacco, uncomplicated
F17.221*	Chewing tobacco, in remission
F17.223	Chewing tobacco, with withdrawal
F17.228	Chewing tobacco, with other nicotine-induced disorders
F17.229	Chewing tobacco, with unspecified nicotine-induced disorders



F17.290*	Other tobacco product, uncomplicated
F17.291*	Other tobacco product, in remission
F17.293	Other tobacco product, with withdrawal
F17.298	Other tobacco product, with other nicotine-induced disorders
F17.299	Other tobacco product, with unspecified nicotine-induced disorders

ZCODES

ICD-10	Description: All with Nicotine Dependence
Diagnosis Code	
Z57.31	Occupational exposure to environmental tobacco smoke
	May not be used with Z77.22 exposure to environmental smoke
Z77.22	Contact with and suspected exposure to environmental smoke
	May not be used with a F17.2 tobacco dependence or Z72 tobacco use code.
Z71.6	Counseling and Medicaid Advice – tobacco abuse counseling
Z72.0	Problems Related to Lifestyle and tobacco use not otherwise specified
Z87.891	Personal history of nicotine dependence
	May not be used with F17.2 current nicotine dependence code.
Z13.89	Encounter for screening for other disorder. Use for tobacco use screening.

Maternal Tobacco Use and Newborn Exposure

• For maternal use add an additional F17 code to indicate type of tobacco

ICD-10	Description: All with Nicotine Dependence
Diagnosis Code	
O99.330	Smoking (tobacco) complicating pregnancy, unspecified trimester
O99.331	Smoking (tobacco) complicating pregnancy, first trimester
O99.332	Smoking (tobacco) complicating pregnancy, second trimester
O99.333	Smoking (tobacco) complicating pregnancy, third trimester
O99.334	Smoking (tobacco) complicating childbirth



O99.335	Smoking (tobacco) complicating puerperium
P04.2	Newborn suspected to be affected by exposure in utero to tobacco smoke May not be used with P96.81 newborn exposure to environmental tobacco smoke.
P96.81	Exposure to (parental) (environmental) tobacco smoke in the perinatal period

Please note: the plan may request medical records at any time to support the services rendered.

We appreciate the excellent care you provide to our members. If you have any questions, please feel free to contact us via e-mail: **FLMedicaidProviderRelations@Aetna.com**. You can also fax us at 1-844-235-1340 or call us through our Provider Relations telephone line: 1-844-528-5815.

Thank you

Provider Relations Department

Aetna Better Health of Florida

CONFIDENTIALITY NOTICE This message is intended only for the user of the individual or entity to which it is addressed and may contain confidential and proprietary information. If you are not the intended recipient of the employee or agent responsible for delivering the message to the intended recipient, you are hereby notified that any dissemination, distribution, or copying of this communication is prohibited. If you received this communication in error, please notify the sender at the phone number above. **NOTICE TO RECIPIENT(S) OF INFORMATION:** Information disclosed to you pertaining to alcohol or drug abuse treatment is protected by federal confidentiality rules (42 CFR Part 2), which prohibit any further disclosure of this information by you without express written consent of the person to whom it pertains of as otherwise permitted by 42 CFR Part 2. A general authorization for the release of medical or other information is NOT sufficient for this purpose. The federal rules restrict any use of the information to criminally investigate or prosecute any alcohol or drug abuse patient.