

## 2020 Healthcare Effectiveness Data & Information Set results (HEDIS®)

At Aetna Better Health of Kentucky (ABHKY), our goal is to assist you in getting members the care they need to improve their health. How well are we doing? One way to find out is a tool called HEDIS®. This chart shows our HEDIS® results for 2020. It also shows the 50th percentile of HEDIS® results for Medicaid in the U.S. overall for 2019. In most cases, the higher the number, the better the results.

2020 HEDIS Results		
Measure/Data Element	2020 Final HEDIS Rate	NCQA 2020 50th %tile
<b>Effectiveness of Care: Prevention and Screening</b>		
<b>Adult BMI Assessment (aba)</b>	97.32%	91.33
<b>Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (wcc)</b>		
<i>BMI Percentile</i>	81.02%	80.5
<i>Counseling for Nutrition</i>	61.56%	71.55
<i>Counseling for Physical Activity</i>	58.15%	66.79
<b>Childhood Immunization Status (cis)</b>		
<i>DTaP</i>	75.67%	77.62
<i>IPV</i>	90.75%	89.78
<i>MMR</i>	88.81%	89.08
<i>HiB</i>	88.08%	88.08
<i>Hepatitis B</i>	91.48%	89.78
<i>VZV</i>	88.08%	88.81
<i>Pneumococcal Conjugate</i>	74.21%	78.10
<i>Hepatitis A</i>	88.56%	85.64
<i>Rotavirus</i>	64.96%	72.02
<i>Influenza</i>	35.77%	49.88
<i>Combination #2</i>	73.48%	74.00
<i>Combination #3</i>	68.37%	71.05
<i>Combination #4</i>	67.88%	68.86
<i>Combination #5</i>	53.77%	61.31
<i>Combination #6</i>	31.39%	42.94
<i>Combination #7</i>	53.28%	59.49
<i>Combination #8</i>	31.14%	41.91

<i>Combination #9</i>	28.71%	37.35
<i>Combination #10</i>	28.47%	37.47
<b>Immunizations for Adolescents (ima) Revised</b>		
<i>Meningococcal</i>	83.70%	84.91
<i>Tdap/Td</i>	90.75%	88.65
<i>HPV</i>	34.31%	38.93
<i>Combination #1</i>	82.97%	82.33
<i>Combination #2</i>	30.66%	36.86
<b>Lead Screening in Children (lsc)</b>	64.23%	73.11
<b>Breast Cancer Screening (bcs)</b>	47.98%	58.82
<b>Cervical Cancer Screening (ccs)</b>	58.15%	61.31
<b>Non-Recommended Cervical Cancer Screening in Adolescent Females (ncs) <i>Inverted Rate (lower is better)</i></b>	1.4%	0.71
<b>Chlamydia Screening in Women (chl)</b>		
<i>16-20 Years</i>	48.65%	53.71
<i>21-24 Years</i>	60.81%	65.40
<i>Total</i>	53.07%	58.44
<b>Effectiveness of Care: Respiratory Conditions</b>		
<b>Appropriate Testing for Children with Pharyngitis (cwp)</b>	79.22%	82.26
<b>Appropriate Treatment for Children With URI (uri)</b>	68.51%	91.45
<b>Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis (aab)</b>	29.32%	50.77
<b>Use of Spirometry Testing in the Assessment and Diagnosis of COPD (spr)</b>	25.94%	28.77
<b>Pharmacotherapy Management of COPD Exacerbation (pce)</b>		
<i>Systemic Corticosteroid</i>	86.62%	71.05
<i>Bronchodilator</i>	93.01%	84.71
<b>Medication Management for People With Asthma (mma)</b>		
<i>5-11 Years - Medication Compliance 50%</i>	59.66%	NA

5-11 Years - Medication Compliance 75%	36.70%	31.8
12-18 Years - Medication Compliance 50%	59.86%	NA
12-18 Years - Medication Compliance 75%	35.96%	31.44
19-50 Years - Medication Compliance 50%	67.45%	NA
19-50 Years - Medication Compliance 75%	43.84%	42.79
51-64 Years - Medication Compliance 50%	82.35%	NA
51-64 Years - Medication Compliance 75%	62.94%	55.1
Total - Medication Compliance 50%	63.22%	NA
Total - Medication Compliance 75%	40.02%	38.59
<b>Asthma Medication Ratio (amr)</b>		
5-11 Years	85.35%	73.61
12-18 Years	76.32%	65.39
19-50 Years	51.93%	53.33
51-64 Years	51.53%	56.23
Total	69.69%	62.43
<b>Effectiveness of Care: Cardiovascular Conditions</b>		
<b>Controlling High Blood Pressure (cbp)</b>	66.67%	61.8
<b>Persistence of Beta-Blocker Treatment After a Heart Attack (pbh)</b>	86.76%	80.37
<b>Effectiveness of Care: Diabetes</b>		
<b>Comprehensive Diabetes Care (cdc)</b>		
Hemoglobin A1c (HbA1c) Testing	88.49%	88.79
<b>Inverted Rate (Lower is better)</b> HbA1c Poor Control (>9.0%)	33.75%	37.47
HbA1c Control (<8.0%)	57.08%	51.73
HbA1c Control (<7.0%)	41.12%	36.38
Eye Exam (Retinal) Performed	50.86%	58.64
Medical Attention for Nephropathy	87.56%	90.14
Blood Pressure Control (<140/90 mm Hg)	71.54%	63.75
<b>Effectiveness of Care: Musculoskeletal</b>		

Conditions		
<b>Use of Imaging Studies for Low Back Pain (lbp)</b>	64.96%	74.19
<b>Effectiveness of Care: Behavioral Health</b>		
<b>Antidepressant Medication Management (amm)</b>		
<i>Effective Acute Phase Treatment</i>	50.68%	53.57
<i>Effective Continuation Phase Treatment</i>	33.62%	38.18
<b>Follow-Up Care for Children Prescribed ADHD Medication (add)</b>		
<i>Initiation Phase</i>	53.06%	42.95
<i>Continuation and Maintenance (C&amp;M) Phase</i>	61.93%	54.73
<b>Follow-Up After Hospitalization for Mental Illness (fuh)</b>		
<i>30-Day Follow-Up</i>	62.02%	59.15
<i>7-Day Follow-Up</i>	40.30%	36.68
<b>Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medication (ssd)</b>	83.84%	82.09
<b>Diabetes Monitoring for People With Diabetes and Schizophrenia (smd)</b>	70.39%	70.82
<b>Cardiovascular Monitoring for People With Cardiovascular Disease and Schizophrenia (smc)</b>	65.38%	78.38
<b>Adherence to Antipsychotic Medications for Individuals With Schizophrenia (saa)</b>	57.90%	62.06

<b>Metabolic Monitoring for Children and Adolescents on Antipsychotics (apm)</b>		
<i>Blood Glucose Testing</i>	48.90%	N/A
<i>Cholesterol Testing</i>	31.62%	N/A
<i>Blood Glucose Testing and Cholesterol Testing</i>	30.23%	N/A
<b>Access/Availability of Care</b>		
<b>Adults' Access to Preventive/Ambulatory Health Services (aap)</b>		
<i>20-44 Years</i>	79.87%	78.61
<i>45-64 Years</i>	85.28%	86.48
<i>65+ Years</i>	77.85%	89.13
<b>Children and Adolescents' Access to Primary Care Practitioners (cap)</b>		
<i>12-24 Months</i>	97.99%	96.1
<i>25 Months - 6 Years</i>	92.51%	88.4
<i>7-11 Years</i>	96.10%	91.42
<i>12-19 Years</i>	94.73%	90.21
<b>Annual Dental Visit (adv)</b>		
<i>2-3 Years</i>	44.04%	43.94
<i>4-6 Years</i>	71.65%	66.24
<i>7-10 Years</i>	73.03%	70.
<i>11-14 Years</i>	66.45%	65.93
<i>15-18 Years</i>	55.42%	55.87
<i>19-20 Years</i>	35.48%	39.52
<i>Total</i>	62.77%	60.15

<b>Initiation and Engagement of AOD Dependence Treatment (iet)</b>		
<i>Initiation of AOD Treatment: 13-17 Years</i>	51.31%	42.03
<i>Engagement of AOD Treatment: 13-17 Years</i>	17.28%	13.65
<i>Initiation of AOD Treatment: 18+ Years</i>	51.47%	43.03
<i>Engagement of AOD Treatment: 18+ Years</i>	24.07%	13.91
<i>Initiation of AOD Treatment: Total</i>	51.46%	41.76
<i>Engagement of AOD Treatment: Total</i>	23.64%	14.18
<b>Prenatal and Postpartum Care (ppc)</b>		
<i>Timeliness of Prenatal Care</i>	88.81%	89.05
<i>Postpartum Care</i>	72.26%	76.4
<b>Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (app)</b>		
<i>1-11 Years</i>	68.00%	63.64
<i>12-17 Years</i>	66.67%	64.97
<i>Total</i>	67.31%	64.89
<b>Well-Child Visits in the First 15 Months of Life (w15) <i>Lower is Better for 0 to 5 Visits</i></b>		
<i>0 Visits</i>	0.24%	1.46
<i>1 Visit</i>	2.19%	1.7
<i>2 Visits</i>	2.92%	2.68
<i>3 Visits</i>	3.16%	4.14
<i>4 Visits</i>	8.27%	7.54
<i>5 Visits</i>	13.63%	13.89
<i>6+ Visits</i>	69.59%	67.88
<b>Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life (w34)</b>	67.88%	74.7
<b>Adolescent Well-Care Visits (awc)</b>	48.91%	57.18
<b>Statin Therapy for Patients With Cardiovascular Disease (spc) <i>Second Year Measure</i></b>		
<i>Received Statin Therapy - 21-75years (Male)</i>	83.54%	81.88

<i>Statin Adherence 80% - 21-75 years (Male)</i>	66.72%	68.99
<i>Received Statin Therapy - 40-75years (Female)</i>	78.95%	77.21
<i>Statin Adherence 80% - 40-75 years (Female)</i>	63.05%	67.39
<i>Received Statin Therapy - Total</i>	81.48%	79.94
<i>Statin Adherence 80% - Total</i>	65.12%	67.94
<b>Statin Therapy for Patients With Diabetes (spd)</b>		
<i>Received Statin Therapy</i>	65.30%	65.92
<i>Statin Adherence 80%</i>	61.77%	64.25