

2021 Healthcare Effectiveness Data & Information Set results (HEDIS®)

At Aetna Better Health of Kentucky (ABHK), our goal is to help you get the care you need to improve your health. That’s why we remind you about care you should get. How well are we doing? One way to find out is a tool called HEDIS®. This chart shows our HEDIS® results for 2021. It also shows the 50th percentile of HEDIS® results for Medicaid in the U.S. overall for 2020. In most cases, the higher the number, the better the results.

2021 HEDIS Results		
Measure/Data Element	2021 Final HEDIS Rate	NCQA 2021 50th %tile
Effectiveness of Care: Prevention and Screening		
Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (wcc)		
<i>BMI Percentile</i>	84.43%	76.64
<i>Counseling for Nutrition</i>	56.45%	70.11
<i>Counseling for Physical Activity</i>	52.55%	66.18
Childhood Immunization Status (cis)		
<i>DTaP</i>	77.37%	74.45
<i>IPV</i>	91.97%	88.32
<i>MMR</i>	87.59%	88.08
<i>HiB</i>	88.32%	87.06
<i>Hepatitis B</i>	93.43%	88.56
<i>VZV</i>	87.35%	87.35
<i>Pneumococcal Conjugate</i>	80.78%	76.16
<i>Hepatitis A</i>	87.83%	84.67
<i>Rotavirus</i>	72.51%	72.08
<i>Influenza</i>	39.90%	50.61
<i>Combination #2</i>	74.70%	71.18
<i>Combination #3</i>	72.75%	67.88
<i>Combination #4</i>	71.29%	66.91
<i>Combination #5</i>	60.58%	59.37
<i>Combination #6</i>	36.25%	42.09
<i>Combination #7</i>	59.85%	58.64
<i>Combination #8</i>	36.01%	41.43

<i>Combination #9</i>	31.87%	37.96
<i>Combination #10</i>	31.87%	38.20
Immunizations for Adolescents (ima) Revised		
<i>Meningococcal</i>	82.97%	84.18
<i>Tdap/Td</i>	89.29%	87.46
<i>HPV</i>	37.71%	38.44
<i>Combination #1</i>	82.73%	76.97
<i>Combination #2</i>	34.55%	36.74
Lead Screening in Children (lsc)	64.48%	71.53
Breast Cancer Screening (bcs)	44.28%	53.93
Cervical Cancer Screening (ccs)	56.20%	59.12
Non-Recommended Cervical Cancer Screening in Adolescent Females (ncs) <i>Inverted Rate (lower is better)</i>	0.93%	0.57
Chlamydia Screening in Women (chl)		
<i>16-20 Years</i>	46.07%	50.46
<i>21-24 Years</i>	59.07%	60.65
<i>Total</i>	51.10%	54.91
Effectiveness of Care: Respiratory Conditions		
Appropriate Testing for Children with Pharyngitis (cwp)	80.36%	76.44
Appropriate Treatment for Children With URI (uri)	71.25%	88.86
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis (aab)	31.54%	54.06
Use of Spirometry Testing in the Assessment and Diagnosis of COPD (spr)	24.18%	26.12
Pharmacotherapy Management of COPD Exacerbation (pce)		
<i>Systemic Corticosteroid</i>	85.38%	70.33
<i>Bronchodilator</i>	90.58%	85.08
Asthma Medication Ratio (amr)		
<i>5-11 Years</i>	86.23%	79.49
<i>12-18 Years</i>	76.29%	69.18
<i>19-50 Years</i>	55.87%	56.30
<i>51-64 Years</i>	51.54%	57.14
<i>Total</i>	69.03%	64.78

Effectiveness of Care: Cardiovascular Conditions		
Controlling High Blood Pressure (cbp)	60.10%	55.35
Persistence of Beta-Blocker Treatment After a Heart Attack (pbh)	80.00%	81.21
Effectiveness of Care: Diabetes		
Comprehensive Diabetes Care (cdc)		
<i>Hemoglobin A1c (HbA1c) Testing</i>	80.29%	82.97
<i>Inverted Rate (Lower is better)</i> <i>HbA1c Poor Control (>9.0%)</i>	38.44%	43.19
<i>HbA1c Control (<8.0%)</i>	49.15%	46.83
<i>Eye Exam (Retinal) Performed</i>	46.72%	51.36
<i>Blood Pressure Control (<140/90 mm Hg)</i>	63.26%	58.52
Effectiveness of Care: Musculoskeletal Conditions		
Use of Imaging Studies for Low Back Pain (lbp)	67.42%	75.61
Effectiveness of Care: Behavioral Health		
Antidepressant Medication Management (amm)		
<i>Effective Acute Phase Treatment</i>	51.56%	56.66
<i>Effective Continuation Phase Treatment</i>	35.30%	40.28
Follow-Up Care for Children Prescribed ADHD Medication (add)		
<i>Initiation Phase</i>	55.45%	44.51
<i>Continuation and Maintenance (C&M) Phase</i>	65.00%	55.96
Follow-Up After Hospitalization for Mental Illness (fuh)		
<i>30-Day Follow-Up</i>	62.72%	60.08
<i>7-Day Follow-Up</i>	40.81%	38.95
Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medication (ssd)	77.83%	76.64
Diabetes Monitoring for People With Diabetes and Schizophrenia (smd)	66.08%	64.40

Cardiovascular Monitoring for People With Cardiovascular Disease and Schizophrenia (smc)	57.89%	73.43
Adherence to Antipsychotic Medications for Individuals With Schizophrenia (saa)	57.74%	63.46
Metabolic Monitoring for Children and Adolescents on Antipsychotics (apm)		
<i>Blood Glucose Testing</i>	41.85%	48.49
<i>Cholesterol Testing</i>	26.82%	31.99
<i>Blood Glucose Testing and Cholesterol Testing</i>	25.60%	30.58
Access/Availability of Care		
Adults' Access to Preventive/Ambulatory Health Services (aap)		
<i>20-44 Years</i>	78.51%	75.32
<i>45-64 Years</i>	83.86%	83.18
<i>65+ Years</i>	69.21%	82.72
<i>Total</i>	80.16%	78.3
Annual Dental Visit (adv)		
<i>2-3 Years</i>	34.03%	30.74
<i>4-6 Years</i>	55.58%	50.59
<i>7-10 Years</i>	56.81%	54.26
<i>11-14 Years</i>	51.45%	51.11
<i>15-18 Years</i>	43.58%	44.72
<i>19-20 Years</i>	27.90%	30.08
<i>Total</i>	48.51%	45.77
Initiation and Engagement of AOD Dependence Treatment (iet)		
<i>Initiation of AOD Treatment: 13-17 Years</i>	35.69%	44.03
<i>Engagement of AOD Treatment: 13-17 Years</i>	9.54%	12.41
<i>Initiation of AOD Treatment: 18+ Years</i>	50.55%	42.7
<i>Engagement of AOD Treatment: 18+ Years</i>	23.75%	13.86
<i>Initiation of AOD Treatment: Total</i>	49.60%	43.07
<i>Engagement of AOD Treatment: Total</i>	22.85%	13.98
Prenatal and Postpartum Care (ppc)		
<i>Timeliness of Prenatal Care</i>	86.37%	85.89

<i>Postpartum Care</i>	76.40%	76.4
Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (app)		
<i>1-11 Years</i>	64.54%	62.38
<i>12-17 Years</i>	61.04%	63.48
<i>Total</i>	62.65%	63.03
Well-Child Visits in the First 30 Months of Life (w30)		
<i>First 15 Months</i>	55.92%	54.92
<i>15 Months - 30 Months</i>	66.25%	70.67
Child and Adolescent Well Care Visit (wcv)	36.16%	45.31
Statin Therapy for Patients With Cardiovascular Disease (spc) <i>Second Year Measure</i>		
<i>Received Statin Therapy - 21-75 years (Male)</i>	80.63%	82.02
<i>Statin Adherence 80% - 21-75 years (Male)</i>	71.16%	71.77
<i>Received Statin Therapy - 40-75 years (Female)</i>	79.35%	77.73
<i>Statin Adherence 80% - 40-75 years (Female)</i>	69.11%	72.63
<i>Received Statin Therapy - Total</i>	80.07%	80.34
<i>Statin Adherence 80% - Total</i>	70.27%	72.21
Statin Therapy for Patients With Diabetes (spd)		
<i>Received Statin Therapy</i>	64.97%	66.47
<i>Statin Adherence 80%</i>	67.97%	68.75