

ABIRATERONE

Products Affected

- Abiraterone Acetate
- Zytiga TABS 500MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Node-positive (N1), non-metastatic (M0) prostate cancer
Exclusion Criteria	N/A
Required Medical Information	The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ACITRETIN

Products Affected

- Acitretin

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Prevention of non-melanoma skin cancers in high risk individuals, Lichen planus, Keratosis follicularis (Darier Disease)
Exclusion Criteria	N/A
Required Medical Information	Psoriasis: The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to methotrexate or cyclosporine.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ACTIMMUNE

Products Affected

- Actimmune

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Mycosis fungoides, Sezary syndrome.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ADEMPAS

Products Affected

- Adempas

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): PAH was confirmed by right heart catheterization. For PAH new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, AND 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. For chronic thromboembolic pulmonary hypertension (CTEPH) (WHO Group 4): 1) Patient has persistent or recurrent CTEPH after pulmonary endarterectomy (PEA), OR 2) Patient has inoperable CTEPH with the diagnosis confirmed by right heart catheterization AND by computed tomography (CT), magnetic resonance imaging (MRI), or pulmonary angiography.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AIMOVIG

Products Affected

- Aimovig

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the preventive treatment of migraine, initial: 1) The patient experienced an inadequate treatment response with a 4-week trial of any one of the following: Antiepileptic drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants OR 2) The patient experienced an intolerance or has a contraindication that would prohibit a 4-week trial of any one of the following: Antiepileptic drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants. For preventive treatment of migraine, continuation: The patient received at least 3 months of treatment with the requested drug, and the patient had a reduction in migraine days per month from baseline.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial: 3 months, Continuation: Plan Year
Other Criteria	N/A

AKEEGA

Products Affected

- Akeega

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ALECENSA

Products Affected

- Alecensa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent ALK-positive non-small cell lung cancer (NSCLC), brain metastases from ALK-positive NSCLC.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ALOSETRON

Products Affected

- Alosetron Hydrochloride

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For severe diarrhea-predominant irritable bowel syndrome (IBS): 1) The requested drug is being prescribed for a biological female or a person that self-identifies as a female, 2) chronic IBS symptoms lasting at least 6 months, 3) gastrointestinal tract abnormalities have been ruled out, AND 4) inadequate response to one conventional therapy (e.g., antispasmodics, antidepressants, antidiarrheals).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ALPHA1-PROTEINASE INHIBITOR

Products Affected

- Prolastin-c

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For alpha1-proteinase inhibitor deficiency: Patient must have 1) clinically evident emphysema and 2) pretreatment serum alpha1-proteinase inhibitor level less than 11 micromol/L (80 mg/dL by radial immunodiffusion or 50 mg/dL by nephelometry).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ALUNBRIG

Products Affected

- Alunbrig

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent ALK-positive non-small cell lung cancer (NSCLC), brain metastases from ALK-positive NSCLC, Inflammatory myofibroblastic tumor (IMT) with ALK translocation.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AMBRISENTAN

Products Affected

- Ambrisentan

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): Diagnosis was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) Pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ARCALYST

Products Affected

- Arcalyst

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Prevention of gout flares in patients initiating or continuing urate-lowering therapy.
Exclusion Criteria	N/A
Required Medical Information	For prevention of gout flares in patients initiating or continuing urate-lowering therapy (e.g., allopurinol) (new starts): 1) two or more gout flares within the previous 12 months, AND 2) inadequate response, intolerance or contraindication to maximum tolerated doses of a non-steroidal anti-inflammatory drug (NSAID) and colchicine, AND 3) concurrent use with urate-lowering therapy. For prevention of gout flares in patients initiating or continuing urate-lowering therapy (e.g., allopurinol) (continuation): 1) patient must have achieved or maintained a clinical benefit (i.e., a fewer number of gout attacks or fewer flare days) compared to baseline, AND 2) continued use of urate-lowering therapy concurrently with the requested drug. For recurrent pericarditis: patient must have had an inadequate response, intolerance or contraindication to maximum tolerated doses of an NSAID and colchicine.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	For prevention of gout flares: 4 months. Other: Plan Year
Other Criteria	N/A

ARMODAFINIL

Products Affected

- Armodafinil

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1) The patient has a diagnosis of narcolepsy and the diagnosis is confirmed by sleep lab evaluation OR 2) The patient has a diagnosis of Shift Work Disorder (SWD) OR 3) The patient has a diagnosis of obstructive sleep apnea (OSA) and the diagnosis is confirmed by polysomnography.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ASPARLAS

Products Affected

- Asparlas

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	21 years of age or younger
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AUBAGIO

Products Affected

- Aubagio

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AUSTEDO

Products Affected

- Austedo
- Austedo Xr
- Austedo Xr Patient Titration Kit

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Tourette's syndrome
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AUVELITY

Products Affected

- Auvelity

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For Major Depressive Disorder (MDD): The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to two of the following: serotonin and norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), mirtazapine, bupropion.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AVONEX

Products Affected

- Avonex INJ 30MCG/0.5ML
- Avonex Pen

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AYVAKIT

Products Affected

- Ayvakit

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Myeloid and lymphoid neoplasms with eosinophilia, gastrointestinal stromal tumor (GIST) for unresectable, recurrent, or metastatic disease without platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation.
Exclusion Criteria	N/A
Required Medical Information	For myeloid and lymphoid neoplasms with eosinophilia, the patient meets all of the following criteria: 1) The disease is FIP1L1- PDGFRA rearrangement-positive, AND 2) The disease harbors a PDGFRA D842A mutation, AND 3) The disease is resistant to imatinib. For GIST, the patient meets either of the following criteria: 1) The disease harbors PDGFRA exon 18 mutation, including PDGFRA D842V mutations, OR 2) The requested drug will be used after failure on at least two Food and Drug Administration (FDA)-approved therapies in unresectable, recurrent, or metastatic disease without PDGFRA exon 18 mutation. For systemic mastocytosis: 1) The patient has a diagnosis of indolent systemic mastocytosis or advanced systemic mastocytosis (including aggressive systemic mastocytosis [ASM], systemic mastocytosis with associated hematological neoplasm [SM-AHN], and mast cell leukemia [MCL]) AND 2) The patient has a platelet count of greater than or equal to 50,000/microliter (mcL).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BALVERSA

Products Affected

- Balversa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent primary carcinoma of the urethra, recurrent or persistent urothelial carcinoma of the bladder.
Exclusion Criteria	N/A
Required Medical Information	For urothelial carcinoma: Disease has susceptible fibroblast growth factor receptor 3 (FGFR3) or fibroblast growth factor receptor 2 (FGFR2) genetic alterations AND the requested drug will be used as subsequent therapy for any of the following: a) locally advanced or metastatic urothelial carcinoma, b) recurrent primary carcinoma of the urethra, c) stage II urothelial carcinoma of the bladder if tumor is present following reassessment of tumor status 2-3 months after primary treatment with bladder preserving concurrent chemoradiotherapy, d) urothelial carcinoma of the bladder with metastatic or local recurrence post cystectomy, or e) urothelial carcinoma of the bladder with muscle invasive local recurrence or persistent disease in a preserved bladder.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BANZEL

Products Affected

- Rufinamide

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	1 year of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BENLYSTA

Products Affected

- Benlysta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	For patients new to therapy: severe active central nervous system lupus.
Required Medical Information	For systemic lupus erythematosus (SLE): 1) Patient is currently receiving a stable standard therapy regimen (e.g., corticosteroid or antimalarial) for SLE OR 2) patient is not currently receiving a stable standard therapy regimen for SLE because the patient experienced an intolerance or has a contraindication to standard therapy regimens. For lupus nephritis: 1) patient is currently receiving a stable standard therapy regimen (e.g., corticosteroid) for lupus nephritis OR 2) patient is not currently receiving a stable standard therapy regimen for lupus nephritis because the patient experienced an intolerance or has a contraindication to standard therapy regimens.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BERINERT

Products Affected

- Berinert

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Short-term preprocedural prophylaxis for hereditary angioedema (HAE) attacks
Exclusion Criteria	N/A
Required Medical Information	For hereditary angioedema (HAE): The requested drug is being used for the treatment of acute angioedema attacks. Patient has HAE with C1 inhibitor deficiency or dysfunction confirmed by laboratory testing OR patient has HAE with normal C1 inhibitor confirmed by laboratory testing. For patients with HAE with normal C1 inhibitor, EITHER 1) Patient tested positive for an F12, angiotensin-1, plasminogen, kininogen-1 (KNG1), heparan sulfate-glucosamine 3-O-sulfotransferase 6 (HS3ST6), or myoferlin (MYOF) gene mutation OR 2) Patient has a family history of angioedema and the angioedema was refractory to a trial of high-dose antihistamine therapy for at least one month.
Age Restrictions	5 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
Coverage Duration	Plan Year
Other Criteria	N/A

BESREMI

Products Affected

- Besremi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BETASERON

Products Affected

- Betaseron

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BEXAROTENE

Products Affected

- Bexarotene CAPS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Mycosis fungoides, Sezary syndrome, CD30-positive primary cutaneous anaplastic large cell lymphoma, CD30-positive lymphomatoid papulosis.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BOSENTAN

Products Affected

- Bosentan

- Tracleer TBSO

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): Diagnosis was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) Pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BOSULIF

Products Affected

- Bosulif

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL), myeloid and/or lymphoid neoplasms with eosinophilia and ABL1 rearrangement in the chronic phase or blast phase
Exclusion Criteria	N/A
Required Medical Information	For chronic myeloid leukemia (CML) or acute lymphoblastic leukemia (ALL), including patients who have received a hematopoietic stem cell transplant: 1) Diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene, and 2) If patient experienced resistance to an alternative tyrosine kinase inhibitor, patient is negative for all of the following mutations: T315I, G250E, V299L, and F317L. For CML, including patients newly diagnosed with CML and patients who have received a hematopoietic stem cell transplant: patient has experienced resistance or intolerance to imatinib or dasatinib.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BRAFTOVI

Products Affected

- Braftovi CAPS 75MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Adjuvant systemic therapy for cutaneous melanoma
Exclusion Criteria	N/A
Required Medical Information	For colorectal cancer: The patient must meet both of the following criteria: 1) Tumor is positive for BRAF V600E mutation, 2) The requested drug will be used for either of the following: a) as subsequent therapy for advanced or metastatic disease, or b) as primary treatment for unresectable metachronous metastases. For cutaneous melanoma: The patient must meet all of the following criteria: 1) Tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K), 2) The requested drug will be used in combination with binimetinib, and 3) The requested drug will be used for either of the following: a) unresectable or metastatic disease, or b) adjuvant systemic therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BRIVIACT

Products Affected

- Brivact ORAL SOLN
- Brivact TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1) The patient has experienced an inadequate treatment response, intolerance, or contraindication to a generic anticonvulsant AND 2) The patient has experienced an inadequate treatment response, intolerance, or contraindication to any of the following: Aptiom, Vimpat, Xcopri, Spritam.
Age Restrictions	1 month of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BRIVIACT INJ

Products Affected

- Briviact INJ

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1) The patient has experienced an inadequate treatment response, intolerance, or contraindication to a generic anticonvulsant AND 2) The patient has experienced an inadequate treatment response, intolerance, or contraindication to any of the following: Aptiom, Vimpat, Xcopri, Spritam.
Age Restrictions	1 month of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BRUKINSA

Products Affected

- Brukinsa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For marginal zone lymphoma: 1) the requested drug is being used for the treatment of relapsed or refractory disease AND the patient has received at least one anti-CD20-based regimen, OR 2) the requested drug is being used for the treatment of refractory or progressive disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BUPRENORPHINE

Products Affected

- Buprenorphine Hcl SUBL

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug is being prescribed for the treatment of opioid use disorder AND patient meets one of the following: 1) The patient is pregnant or breastfeeding, and the requested drug is being prescribed for induction therapy and/or subsequent maintenance therapy for treatment of opioid use disorder OR 2) The requested drug is being prescribed for induction therapy for transition from opioid use to treatment of opioid use disorder OR 3) The requested drug is being prescribed for maintenance therapy for treatment of opioid use disorder in a patient who is intolerant to naloxone.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BUPRENORPHINE PATCH

Products Affected

- Buprenorphine PTWK

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug is being prescribed for pain associated with cancer, sickle cell disease, a terminal condition, or pain being managed through palliative care OR the patient meets all of the following: 1) The requested drug is being prescribed for pain severe enough to require daily, around-the-clock, long-term treatment in a patient who has been taking an opioid AND 2) The patient can safely take the requested dose based on their history of opioid use [Note: This drug should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.] AND 3) The patient has been evaluated and the patient will be monitored for the development of opioid use disorder AND 4) This request is for continuation of therapy for a patient who has been receiving an extended-release opioid agent for at least 30 days OR the patient has taken an immediate-release opioid for at least one week.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BYDUREON AET

Products Affected

- Bydureon Bcise

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	10 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	The Prior Authorization only applies to patients whose claim is not submitted with an ICD-10 code indicating a diagnosis of type 2 diabetes mellitus OR to patients who do not have a history of an antidiabetic drug (EXCLUDING glucagon-like peptide receptor agonists [GLP-1 RAs] and combination glucose-dependent insulinotropic polypeptide [GIP] and GLP-1 RAs).

BYETTA AET

Products Affected

- Byetta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	10 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	The Prior Authorization only applies to patients whose claim is not submitted with an ICD-10 code indicating a diagnosis of type 2 diabetes mellitus OR to patients who do not have a history of an antidiabetic drug (EXCLUDING glucagon-like peptide receptor agonists [GLP-1 RAs] and combination glucose-dependent insulinotropic polypeptide [GIP] and GLP-1 RAs).

CABOMETYX

Products Affected

- Cabometyx

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Non-small cell lung cancer, Ewing sarcoma, osteosarcoma, gastrointestinal stromal tumor
Exclusion Criteria	N/A
Required Medical Information	For renal cell carcinoma: The disease is advanced, relapsed, or stage IV. For non-small cell lung cancer: 1) the disease is rearranged during transfection (RET) positive AND 2) the disease is recurrent, advanced, or metastatic. For hepatocellular carcinoma: the requested drug will be used as subsequent treatment. For gastrointestinal stromal tumor (GIST): the disease is unresectable, recurrent, or metastatic AND the patient has failed on an FDA-approved therapy (e.g., imatinib, sunitinib, regorafenib, ripretinib). For Ewing sarcoma and osteosarcoma: the requested drug will be used as subsequent therapy. For differentiated thyroid cancer (DTC) (follicular, papillary, Hurthle cell): 1) The disease is locally advanced or metastatic disease, 2) the disease has progressed after a vascular endothelial growth factor receptor (VEGFR)- targeted therapy, AND 3) the patient is refractory to radioactive iodine therapy (RAI) or ineligible for RAI.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CALCIPOTRIENE

Products Affected

- Calcipotriene CREA
- Calcipotriene OINT
- Calcipotriene SOLN
- Calcitrene
- Calcitriol OINT
- Enstilar

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For Treatment of Psoriasis: The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to a topical steroid.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CALQUENCE

Products Affected

- Calquence

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Waldenstrom macroglobulinemia, lymphoplasmacytic lymphoma, gastric mucosa-associated lymphoid tissue (MALT) lymphoma, non-gastric MALT lymphoma (noncutaneous), nodal marginal zone lymphoma, splenic marginal zone lymphoma
Exclusion Criteria	N/A
Required Medical Information	For gastric MALT lymphoma, non-gastric MALT lymphoma, nodal marginal zone lymphoma, and splenic marginal zone lymphoma: the requested drug is being used for the treatment of refractory or progressive disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CAPRELSA

Products Affected

- Caprelsa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Differentiated thyroid carcinoma: papillary, follicular, and Hurthle cell.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CARAC

Products Affected

- Fluorouracil CREA 0.5%

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Patients who are pregnant or may become pregnant. Patients with dihydropyrimidine dehydrogenase (DPD) enzyme deficiency.
Required Medical Information	If being used as part of a compounded product, all active ingredients in the compounded product are Food and Drug Administration (FDA) approved for topical use.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A

CARBAGLU

Products Affected

- Carglumic Acid

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For N-acetylglutamate synthase (NAGS) deficiency: Diagnosis of NAGS deficiency was confirmed by enzymatic, biochemical, or genetic testing.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CAYSTON

Products Affected

- Cayston

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For treatment of respiratory symptoms in cystic fibrosis patients: 1) Pseudomonas aeruginosa is present in the patient's airway cultures OR 2) The patient has a history of pseudomonas aeruginosa infection or colonization in the airways.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CERDELGA

Products Affected

- Cerdelga

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For type 1 Gaucher disease (GD1): 1) The diagnosis was confirmed by an enzyme assay demonstrating a deficiency of beta-glucocerebrosidase enzyme activity or by genetic testing, and 2) The patient's CYP2D6 metabolizer status has been established using an FDA-cleared test, and 3) The patient is a CYP2D6 extensive metabolizer, an intermediate metabolizer, or a poor metabolizer.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CHANTIX

Products Affected

- Varenicline Starting Month Box
- Varenicline Tartrate

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	This Prior Authorization requirement applies after the patient has received a cumulative 24 weeks of therapy in a Plan Year.

CHLORDIAZEPOXIDE - 65

Products Affected

- Chlordiazepoxide Hcl CAPS 10MG, 5MG
- Chlordiazepoxide Hydrochloride CAPS 25MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For all indications: the prescriber must acknowledge the benefit of therapy with the prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For the management of anxiety disorders: 1) the requested drug is being used concurrently with a selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the SSRI/SNRI becomes effective for the symptoms of anxiety, OR 2) the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to AT LEAST TWO agents from the following classes: a) selective serotonin reuptake inhibitors (SSRIs), or b) serotonin-norepinephrine reuptake inhibitors (SNRIs).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Short-term relief anxiety-preop apprehens and anx-1 mo, Anxiety Disorder-4 mo, Alc Withdrawal-PlanYR
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older.

CLOBAZAM

Products Affected

- Clobazam

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CLOMIPRAMINE

Products Affected

- Clomipramine Hydrochloride

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Depression, Panic Disorder
Exclusion Criteria	N/A
Required Medical Information	1) The requested drug is being prescribed for one of the following: a) Obsessive-Compulsive Disorder (OCD), b) Panic Disorder AND 2) The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to any of the following: a) a serotonin and norepinephrine reuptake inhibitor (SNRI), b) a selective serotonin reuptake inhibitor (SSRI) OR 3) The requested drug is being prescribed for Depression AND 4) The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to two of the following: a) serotonin and norepinephrine reuptake inhibitors (SNRIs), b) selective serotonin reuptake inhibitors (SSRIs), c) mirtazapine, d) bupropion
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CLORAZEPATE

Products Affected

- Clorazepate Dipotassium TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For all indications: The prescriber must acknowledge the benefit of therapy with this prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For the management of anxiety disorders: 1) The requested drug is being used concurrently with a selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the SSRI/SNRI becomes effective for the symptoms of anxiety, OR 2) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to AT LEAST TWO agents from the following classes: a) selective serotonin reuptake inhibitors (SSRIs), b) serotonin-norepinephrine reuptake inhibitors (SNRIs).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Short-term relief anxiety-1 month, Anxiety Disorders-4 months, All other Diagnoses-Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older.

CLOZAPINE ODT

Products Affected

- Clozapine Odt

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

COLY-MYCIN

Products Affected

- Colistimethate Sodium INJ

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug is being used intravenously (IV) or intramuscularly (IM). Administration will not be via nebulizer (Note: CMS endorsed compendia do not support inhalation/nebulization of colistimethate.). The infection is proven or strongly suspected to be caused by susceptible bacteria based on: 1) culture and susceptibility information OR 2) local epidemiology and susceptibility patterns.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist.
Coverage Duration	Initial approval: 3 months. Renewal: Plan Year
Other Criteria	N/A

COMETRIQ

Products Affected

- Cometriq

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Non-small cell lung cancer (NSCLC), differentiated thyroid carcinoma: papillary, follicular, and Hurthle cell.
Exclusion Criteria	N/A
Required Medical Information	For NSCLC: The requested medication is used for NSCLC when the patient's disease expresses rearranged during transfection (RET) gene rearrangements.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

COPIKTRA

Products Affected

- Copiktra

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL): the patient has relapsed or refractory disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

COTELLIC

Products Affected

- Cotellic

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma), Erdheim-Chester disease, Langerhans cell histiocytosis, Rosai-Dorfman disease
Exclusion Criteria	N/A
Required Medical Information	For adjuvant treatment of melanoma, and central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma): The patient must meet both of the following criteria: 1) The tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K), and 2) The requested drug will be used in combination with vemurafenib. For unresectable or metastatic melanoma: The patient must meet both of the following criteria: 1) The tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K), and 2) The requested drug will be used in combination with vemurafenib (with or without atezolizumab).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CYSTAGON

Products Affected

- Cystagon

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Diagnosis of nephropathic cystinosis was confirmed by ANY of the following: 1) the presence of increased cystine concentration in leukocytes, OR 2) genetic testing, OR 3) demonstration of corneal cystine crystals by slit lamp examination.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CYSTARAN

Products Affected

- Cystaran

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient meets both of the following: 1) Diagnosis of cystinosis was confirmed by ANY of the following: a) the presence of increased cystine concentration in leukocytes, OR b) genetic testing, OR c) demonstration of corneal cystine crystals by slit lamp examination, AND 2) the patient has corneal cystine crystal accumulation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DALFAMPRIDINE

Products Affected

- Dalfampridine Er

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For multiple sclerosis, patient must meet the following: For new starts, prior to initiating therapy, patient demonstrates sustained walking impairment. For continuation of therapy: patient must have experienced an improvement in walking speed OR other objective measure of walking ability since starting the requested drug.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DAURISMO

Products Affected

- Daurismo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Post induction therapy following response to previous therapy with the same regimen for acute myeloid leukemia (AML). Relapsed/refractory AML as a component of repeating the initial successful induction regimen.
Exclusion Criteria	N/A
Required Medical Information	For acute myeloid leukemia: 1) the requested drug must be used in combination with cytarabine, 2) the patient is 75 years of age or older OR has comorbidities that preclude intensive chemotherapy, and 3) the requested drug will be used as treatment for induction therapy, post-induction therapy, or relapsed or refractory disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DEFERASIROX

Products Affected

- Deferasirox

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For chronic iron overload due to blood transfusions: pretreatment serum ferritin level is greater than 1000 mcg/L.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DEMSEER

Products Affected

- Metyrosine

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to an alpha-adrenergic antagonist.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DESVENLAFAXINE

Products Affected

- Desvenlafaxine Er

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to TWO of the following: a) serotonin and norepinephrine reuptake inhibitors (SNRIs), b) selective serotonin reuptake inhibitors (SSRIs), c) mirtazapine, d) bupropion
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DHE NASAL

Products Affected

- Dihydroergotamine Mesylate NASAL SOLN

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Coverage will be denied when used in conjunction with potent CYP3A4 inhibitors (e.g., ritonavir, nelfinavir, indinavir, erythromycin, clarithromycin).
Required Medical Information	The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one triptan 5-HT1 receptor agonist.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DIACOMIT

Products Affected

- Diacomit

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	6 months of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DIAZEPAM

Products Affected

- Diazepam CONC
- Diazepam INJ 5MG/ML
- Diazepam SOLN 5MG/5ML
- Diazepam TABS
- Diazepam Intensol

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For all indications: The prescriber must acknowledge the benefit of therapy with this prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For the management of anxiety disorders: 1) The requested drug is being used concurrently with a selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the SSRI/SNRI becomes effective for the symptoms of anxiety, OR 2) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to AT LEAST TWO agents from the following classes: a) selective serotonin reuptake inhibitors (SSRIs), b) serotonin-norepinephrine reuptake inhibitors (SNRIs).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Short-term relief anx-1 mo, skeletal muscle spasm-3 mo, Anx Disorders-4 mo, Other Diagnoses-PlanYR
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older.

DOPTELET

Products Affected

- Doptelet

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For thrombocytopenia in patients with chronic liver disease: Untransfused platelet count prior to a scheduled procedure is less than 50,000/mcL. For chronic immune thrombocytopenia (ITP): 1) For new starts: a) Patient has had an inadequate response or is intolerant to prior therapy such as corticosteroids or immunoglobulins, AND b) Untransfused platelet count at any point prior to the initiation of the requested medication is less than 30,000/mcL OR 30,000 to 50,000/mcL with symptomatic bleeding or risk factor(s) for bleeding (e.g., undergoing a medical or dental procedure where blood loss is anticipated, comorbidities such as peptic ulcer disease and hypertension, anticoagulation therapy, profession or lifestyle that predisposes patient to trauma). 2) For continuation of therapy, platelet count response to the requested drug: a) Current platelet count is less than or equal to 200,000/mcL OR b) Current platelet count is greater than 200,000/mcL and less than or equal to 400,000/mcL and dosing will be adjusted to a platelet count sufficient to avoid clinically important bleeding.
Age Restrictions	18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Chronic liver disease: 1 month, ITP initial: 6 months, ITP reauthorization: Plan Year
Other Criteria	N/A

DRIZALMA

Products Affected

- Drizalma Sprinkle

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Cancer pain, chemotherapy-induced neuropathic pain
Exclusion Criteria	N/A
Required Medical Information	1) The patient has tried duloxetine capsules OR 2) The patient is unable to take duloxetine capsules for any reason (e.g., difficulty swallowing capsules, requires nasogastric administration).
Age Restrictions	Generalized Anxiety Disorder - 7 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DRONABINOL

Products Affected

- Dronabinol

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For nausea and vomiting associated with cancer chemotherapy: The patient has experienced an inadequate treatment response, intolerance, or contraindication to one oral 5-HT ₃ receptor antagonist.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

DUPIXENT

Products Affected

- Dupixent

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For atopic dermatitis (AD), initial therapy: 1) Patient has moderate-to-severe disease, 2) Patient has had an inadequate treatment response to either a topical corticosteroid or a topical calcineurin inhibitor, OR topical corticosteroids and topical calcineurin inhibitors are not advisable for the patient. For AD, continuation of therapy: the patient achieved or maintained positive clinical response. For moderate-to-severe asthma, initial therapy: Patient meets either of the following: 1) patient is oral corticosteroid dependent and asthma remains inadequately controlled despite current treatment with both of the following medications: a) high-dose inhaled corticosteroid and b) an additional controller (long acting beta2-agonist, long-acting muscarinic antagonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies, OR 2) patient has a baseline blood eosinophil count of at least 150 cells per microliter and their asthma remains inadequately controlled despite current treatment with both of the following medications: a) medium-to-high-dose inhaled corticosteroid and b) additional controller (long acting beta2-agonist, long-acting muscarinic antagonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies. For moderate-to-severe asthma, continuation of therapy: asthma control has improved on treatment with the requested drug. For chronic rhinosinusitis with nasal polyposis (CRSwNP): 1) the requested drug is used as add-on maintenance treatment, AND 2) the patient has experienced an inadequate treatment response to Xhance (fluticasone).
Age Restrictions	Atopic Dermatitis: 6 months of age or older, Asthma: 6 years of age or older, Chronic Rhinosinusitis with Nasal Polyposis and Prurigo Nodularis: 18 years of age or older, Eosinophilic Esophagitis: 12 years of age or older
Prescriber Restrictions	N/A

Coverage Duration	AD, initial: 4 months, PN, initial: 6 months, All other: Plan Year
Other Criteria	For eosinophilic esophagitis (EoE), initial therapy: 1) diagnosis has been confirmed by esophageal biopsy, 2) patient weighs at least 40 kilograms, 3) patient experienced an inadequate treatment response, intolerance, or patient has a contraindication to a topical corticosteroid (e.g., fluticasone propionate or budesonide). For EoE, continuation of therapy: the patient achieved or maintained a positive clinical response. For prurigo nodularis (PN), initial therapy: Patient has had an inadequate treatment response to a topical corticosteroid OR topical corticosteroids are not advisable for the patient. For PN, continuation of therapy: The patient achieved or maintained a positive clinical response.

ELIGARD

Products Affected

- Eligard

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent androgen receptor positive salivary gland tumors
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

EMSAM

Products Affected

- Emsam

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1) Patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to TWO of the following: a) serotonin and norepinephrine reuptake inhibitors (SNRIs), b) selective serotonin reuptake inhibitors (SSRIs), c) mirtazapine, d) bupropion OR 2) Patient is unable to swallow oral formulations.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ENBREL

Products Affected

- Enbrel
- Enbrel Mini
- Enbrel Sureclick

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Hidradenitis suppurativa
Exclusion Criteria	N/A
Required Medical Information	For moderately to severely active rheumatoid arthritis (new starts only): 1) Inadequate response, intolerance or contraindication to methotrexate (MTX) OR 2) Inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. For active ankylosing spondylitis (new starts only): 1) Inadequate response to a non-steroidal anti-inflammatory drug (NSAID) trial OR 2) Intolerance or contraindication to NSAIDs. For moderate to severe plaque psoriasis (new starts only): 1) At least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis AND 2) Patient meets any of the following: a) Patient has experienced an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, OR b) Pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, OR c) Patient has severe psoriasis that warrants a biologic as first-line therapy (i.e. at least 10% of the BSA or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected). For hidradenitis suppurativa (new starts only): patient has severe, refractory disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

EPCLUSA

Products Affected

- Epclusa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hepatitis C virus (HCV): Infection confirmed by presence of HCV RNA in the serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of human immunodeficiency virus (HIV) coinfection, presence or absence of resistance-associated substitutions where applicable, transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current American Association for the Study of Liver Diseases (AASLD) treatment guidelines.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Criteria will be applied consistent with current AASLD-IDSA guidance.
Other Criteria	N/A

EPIDIOLEX

Products Affected

- Epidiolex

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	1 year of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

EPO

Products Affected

- Procrit

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Anemia due to myelodysplastic syndromes (MDS), anemia in congestive heart failure (CHF), anemia in rheumatoid arthritis (RA), anemia due to hepatitis C treatment (ribavirin in combination with either interferon alfa or peginterferon alfa)
Exclusion Criteria	Patients receiving chemotherapy with curative intent. Patients with myeloid cancer.
Required Medical Information	Requirements regarding hemoglobin (Hgb) values exclude values due to a recent transfusion. For initial approval: 1) for all uses except anemia due to chemotherapy or myelodysplastic syndrome (MDS): patient has adequate iron stores (for example, a transferrin saturation [TSAT] greater than or equal to 20%), AND 2) for all uses except surgery: pretreatment (no erythropoietin treatment in previous month) Hgb is less than 10 g/dL (less than 9 g/dL for anemia in congestive heart failure), AND 3) for MDS: pretreatment serum erythropoietin level is 500 international units/L or less. For reauthorizations (patient received erythropoietin treatment in previous month) in all uses except surgery: 1) patient has received at least 12 weeks of erythropoietin therapy, AND 2) patient responded to erythropoietin therapy, AND 3) current Hgb is less than 12 g/dL, AND 4) for all uses except anemia due to chemotherapy or MDS: patient has adequate iron stores (for example, a transferrin saturation [TSAT] greater than or equal to 20%).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	16 weeks

Other Criteria	Coverage includes use in anemia in patients whose religious beliefs forbid blood transfusions. Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual (e.g., used for treatment of anemia for a patient with chronic renal failure who is undergoing dialysis, or furnished from physician's supply incident to a physician service).
-----------------------	--

EPRONTIA

Products Affected

- Eprontia

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For treatment of partial-onset seizures: 1)The patient has experienced an inadequate treatment response, intolerance, or contraindication to a generic anticonvulsant AND 2) the patient has experienced an inadequate treatment response, intolerance, or contraindication to any of the following: Aptiom, Vimpat, Xcopri, Spritam. For monotherapy treatment of primary generalized tonic-clonic seizures: 1) The patient has experienced an inadequate treatment response or intolerance to topiramate tablets or capsules, OR 2) The patient has difficulty swallowing solid oral dosage forms (e.g., tablets, capsules). For adjunctive treatment of primary generalized tonic-clonic seizures: 1) The patient has experienced an inadequate treatment response, intolerance, or contraindication to a generic anticonvulsant AND 2) If the patient is 4 years of age or older, the patient has experienced an inadequate treatment response, intolerance, or contraindication to Spritam or Vimpat. For the preventative treatment of migraines: 1) The patient has experienced an inadequate treatment response or intolerance to topiramate tablets or capsules, OR 2) The patient has difficulty swallowing solid oral dosage forms (e.g., tablets, capsules).
Age Restrictions	Epilepsy: 2 years of age or older, Migraine: 12 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ERGOTAMINE

Products Affected

- Ergotamine Tartrate/caffeine

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Coverage will be denied when used in conjunction with potent CYP3A4 inhibitors (e.g., ritonavir, nelfinavir, indinavir, erythromycin, clarithromycin).
Required Medical Information	The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least ONE triptan 5-HT1 agonist.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ERIVEDGE

Products Affected

- Erivedge

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Adult medulloblastoma
Exclusion Criteria	N/A
Required Medical Information	Adult medulloblastoma: patient has received chemotherapy previously AND has tumor(s) with mutations in the sonic hedgehog pathway
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ERLEADA

Products Affected

- Erleada

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ERLOTINIB

Products Affected

- Erlotinib Hydrochloride TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent non-small cell lung cancer (NSCLC), recurrent chordoma, relapsed or stage IV renal cell carcinoma (RCC), brain metastases from non-small cell lung cancer (NSCLC), recurrent pancreatic cancer.
Exclusion Criteria	N/A
Required Medical Information	For NSCLC (including brain metastases from NSCLC): 1) the disease is recurrent, advanced, or metastatic and 2) the patient has sensitizing EGFR mutation-positive disease. For pancreatic cancer: the disease is locally advanced, unresectable, recurrent, or metastatic.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ESBRIET

Products Affected

- Esbriet CAPS

- Pirfenidone

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For idiopathic pulmonary fibrosis (Initial Review Only): 1) a high-resolution computed tomography (HRCT) study of the chest or a lung biopsy reveals the usual interstitial pneumonia (UIP) pattern, OR 2) HRCT study of the chest reveals a result other than the UIP pattern (e.g., probable UIP, indeterminate for UIP) and the diagnosis is supported either by a lung biopsy or by a multidisciplinary discussion between at least a radiologist and pulmonologist who are experienced in idiopathic pulmonary fibrosis if a lung biopsy has not been conducted.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

EVEROLIMUS

Products Affected

- Everolimus TABS 10MG, 2.5MG, 5MG, 7.5MG

- Everolimus TBSO

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Classic Hodgkin lymphoma, thymomas and thymic carcinomas, previously treated Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma, soft tissue sarcoma (perivascular epithelioid cell tumors (PEComa) and lymphangioliomyomatosis subtypes), gastrointestinal stromal tumors, neuroendocrine tumors of the thymus, well differentiated grade 3 neuroendocrine tumors, thyroid carcinoma (papillary, Hurthle cell, and follicular), endometrial carcinoma, histiocytic neoplasms (Rosai-Dorfman Disease, Erdheim-Chester Disease, Langerhans Cell Histiocytosis)
Exclusion Criteria	N/A
Required Medical Information	For breast cancer: 1) The disease is recurrent, advanced, or metastatic hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, AND 2) The requested drug is prescribed in combination with exemestane, fulvestrant, or tamoxifen, AND 3) The requested drug is used for subsequent treatment. For renal cell carcinoma: The disease is relapsed, advanced, or stage IV. For subependymal giant cell astrocytoma (SEGA): The requested drug is given as adjuvant treatment. For gastrointestinal stromal tumor: The disease is recurrent, unresectable, or metastatic AND the patient failed an FDA-approved therapy (e.g., imatinib, sunitinib, regorafenib, ripretinib). For symptomatic or relapsed/refractory Erdheim-Chester Disease (ECD), symptomatic or relapsed/refractory Rosai-Dorfman Disease, and Langerhans Cell Histiocytosis (LCH): the patient must have a phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha (PIK3CA) mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A

Coverage Duration	Plan Year
Other Criteria	N/A

EXKIVITY

Products Affected

- Exkivity

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FANAPT

Products Affected

- Fanapt

- Fanapt Titration Pack

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For treatment of schizophrenia: 1) The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following generic products: aripiprazole, asenapine, olanzapine, quetiapine, risperidone, ziprasidone AND 2) The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following brand products: Latuda, Rexulti, Secuado, Vraylar.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FARYDAK

Products Affected

- Farydak

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For multiple myeloma: the disease needs to have been previously treated.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FASENRA

Products Affected

- Fasenra

- Fasenra Pen

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For severe asthma: For initial therapy: 1) Either a) Patient has baseline blood eosinophil count of at least 150 cells per microliter OR b) Patient is dependent on systemic corticosteroids, and 2) Patient has a history of severe asthma despite current treatment with both of the following medications: a) inhaled corticosteroid and b) additional controller (long-acting beta2-agonist, long-acting muscarinic antagonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies. For continuation of therapy: Asthma control has improved on treatment with the requested drug, as demonstrated by a reduction in the frequency and/or severity of symptoms and exacerbations or a reduction in the daily maintenance oral corticosteroid dose.
Age Restrictions	12 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FENTANYL PATCH

Products Affected

- Fentanyl PT72

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug is being prescribed for pain associated with cancer, sickle cell disease, a terminal condition, or pain being managed through palliative care OR the patient meets all of the following: 1) The requested drug is being prescribed for pain severe enough to require daily, around-the-clock, long-term treatment in a patient who has been taking an opioid AND 2) The patient can safely take the requested dose based on their history of opioid use [Note: This drug should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.] AND 3) The patient has been evaluated and the patient will be monitored for the development of opioid use disorder AND 4) This request is for continuation of therapy for a patient who has been receiving an extended-release opioid agent for at least 30 days OR the patient has taken an immediate-release opioid for at least one week.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FETZIMA

Products Affected

- Fetzima

- Fetzima Titration Pack

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to TWO of the following: serotonin and norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), mirtazapine, bupropion.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FINTEPLA

Products Affected

- Fintepla

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FLUOROURACIL

Products Affected

- Fluoroplex CREA

- Fluorouracil CREA 5%

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Patients who are pregnant or may become pregnant. Fluorouracil cream 5 percent only; Patients with dihydropyrimidine dehydrogenase (DPD) enzyme deficiency.
Required Medical Information	Applies to new starts only. If being used as part of a compounded product, all active ingredients in the compounded product are Food and Drug Administration (FDA) approved for topical use.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A

FORTAMET/GLUMETZA

Products Affected

- Metformin Hydrochloride Er TB24 500MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient has experienced an intolerance to generic Glucophage XR.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FORTEO

Products Affected

- Forteo INJ 600MCG/2.4ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>For postmenopausal osteoporosis: patient has ONE of the following (1 or 2): 1) a history of fragility fracture, OR 2) A pre-treatment T-score of less than or equal to -2.5 or pre-treatment T-score greater than -2.5 and less than -1 with a high pre-treatment Fracture Risk Assessment Tool (FRAX) fracture probability AND patient has ANY of the following: a) indicators for higher fracture risk (e.g., advanced age, frailty, glucocorticoid therapy, very low T-scores, or increased fall risk), OR b) patient has failed prior treatment with or is intolerant to a previous injectable osteoporosis therapy OR c) patient has had an oral bisphosphonate trial of at least 1-year duration or there is a clinical reason to avoid treatment with an oral bisphosphonate. For primary or hypogonadal osteoporosis in men: patient has one of the following: 1) a history of osteoporotic vertebral or hip fracture, OR 2) pre-treatment T-score of less than or equal to -2.5, OR 3) pre-treatment T-score greater than -2.5 and less than -1 with a high pre-treatment FRAX fracture probability. For glucocorticoid-induced osteoporosis: Patient has had an oral bisphosphonate trial of at least 1-year duration unless patient has a contraindication or intolerance to an oral bisphosphonate, AND patient meets ANY of the following: 1) patient has a history of fragility fracture, OR 2) a pre-treatment T-score of less than or equal to -2.5, OR 3) pre-treatment T-score greater than -2.5 and less than -1 with a high pre-treatment FRAX fracture probability.</p> <p>Continuation of therapy: If the patient has received greater than or equal to 24 months of therapy with any parathyroid hormone analog: 1) The patient remains at or has returned to having a high risk for fracture, AND 2) The benefit of therapy with this prescribed medication outweighs the potential risks for this patient.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A

Coverage Duration	Initial: 24 months, Continuation: Plan Year
Other Criteria	Patient has high FRAX fracture probability if the 10 year probability is either greater than or equal to 20 percent for any major osteoporotic fracture or greater than or equal to 3 percent for hip fracture. If glucocorticoid treatment is greater than 7.5 mg (prednisone equivalent) per day, the estimated risk score generated with FRAX should be multiplied by 1.15 for major osteoporotic fracture and 1.2 for hip fracture.

FOTIVDA

Products Affected

- Fotivda

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For advanced renal cell carcinoma: the following criteria must be met: 1) The disease is relapsed or refractory, 2) The requested drug must be used after at least two prior systemic therapies, and 3) The patient has experienced disease progression or an intolerable adverse event with a trial of cabozantinib (Cabometyx).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FYCOMPA

Products Affected

- Fycompa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For treatment of partial-onset seizures: 1) The patient experienced an inadequate treatment response, intolerance, or contraindication to a generic anticonvulsant AND 2) The patient has experienced an inadequate treatment response, intolerance, or contraindication to any of the following: Aptiom, Vimpat, Xcopri, Spritam. For adjunctive treatment of primary generalized tonic-clonic seizures: 1) The patient experienced an inadequate treatment response, intolerance, or contraindication to a generic anticonvulsant AND 2) The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following: Vimpat, Spritam.
Age Restrictions	Partial-onset seizures: 4 years of age or older. Primary generalized tonic-clonic seizures: 12 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

GATTEX

Products Affected

- Gattex

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For short bowel syndrome (SBS) initial therapy: Adult patients were dependent on parenteral support for at least 12 months. For SBS continuation: Requirement for parenteral support has decreased from baseline while on therapy with the requested drug.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, gastrointestinal surgeon, or nutritional support specialist.
Coverage Duration	Plan Year
Other Criteria	N/A

GAVRETO

Products Affected

- Gavreto

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent rearranged during transfection (RET) rearrangement-positive non-small cell lung cancer
Exclusion Criteria	N/A
Required Medical Information	For non-small cell lung cancer, patient must meet all of the following: 1) The disease is recurrent, advanced, or metastatic, and 2) The tumor is rearranged during transfection (RET) fusion-positive or RET rearrangement-positive.
Age Restrictions	Non-small cell lung cancer: 18 years of age or older. Medullary thyroid cancer and thyroid cancer: 12 years of age or older.
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

GILENYA

Products Affected

- Fingolimod
- Gilenya CAPS 0.5MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

GILOTRIF

Products Affected

- Gilotrif

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For non-small cell lung cancer (NSCLC): Patient meets either of the following: 1) Patient has metastatic squamous NSCLC that progressed after platinum-based chemotherapy, OR 2) Patient has sensitizing epidermal growth factor receptor (EGFR) mutation-positive disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

GLATIRAMER

Products Affected

- Copaxone INJ 20MG/ML, 40MG/ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

GROWTH HORMONE

Products Affected

- Genotropin

- Genotropin Miniquick

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Pediatric patients with closed epiphyses
Required Medical Information	Pediatric growth hormone deficiency (GHD): Patient (pt) is a neonate or was diagnosed with GHD as a neonate OR meets any of the following: 1) younger than 2.5 years old (yo) with pre-treatment (pre-tx) height (ht) more than 2 standard deviations (SD) below mean and slow growth velocity OR 2) 2.5 yo or older AND one of the following: a) pre-tx 1-year ht velocity more than 2 SD below mean OR b) pre-tx ht more than 2 SD below mean and 1-year ht velocity more than 1 SD below mean, AND patient meets any of the following: 1) failed 2 pre-tx growth hormone (GH) stimulation tests (peak below 10 ng/mL), OR 2) pituitary/central nervous system (CNS) disorder (e.g., genetic defects, acquired structural abnormalities, congenital structural abnormalities) and pre-tx insulin-like growth factor-1 (IGF-1) more than 2 SD below mean. Turner syndrome: 1) Confirmed by karyotyping AND 2) pre-tx ht is less than the 5th percentile for age. Small for gestational age (SGA): 1) Birth weight (wt) less than 2500g at gestational age (GA) greater than 37 weeks, OR birth wt or length below 3rd percentile for GA or at least 2 SD below mean for GA, AND 2) did not manifest catch-up growth by age 2.
Age Restrictions	SGA: 2 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist, pediatric endocrinologist, nephrologist, infectious disease specialist, gastroenterologist/nutritional support specialist, or geneticist.
Coverage Duration	Plan Year

Other Criteria	<p>Adult GHD: Pt meets any of the following: 1) failed 2 pre-tx GH stimulation tests, OR 2) pre-tx IGF-1 more than 2 SD below mean AND failed 1 pre-tx GH stimulation test. (Note: Stimulation tests include: a) insulin tolerance test [ITT] [peak GH less than or equal to 5 ng/ml], or b) Macrilen-stimulation test [peak GH level less than 2.8ng/ml], or c) glucagon-stimulation test [GST] [peak GH level less than or equal to 3 ng/ml] for pt with a body mass index [BMI] 25-30 kg/m² and high pretest probability of GHD [e.g., acquired structural abnormalities] or BMI less than 25 kg/m², or d) GST [peak GH level less than or equal to 1 ng/ml] in pt with BMI 25-30 kg/m² and low pretest probability of GHD or BMI greater than 30 kg/m²), OR 3) organic hypothalamic-pituitary disease (e.g., suprasellar mass with previous surgery and cranial irradiation) with 3 or more pituitary hormone deficiencies AND pre-tx IGF-1 more than 2 SD below mean, OR 4) genetic or structural hypothalamic-pituitary defects, OR 5) childhood-onset GHD with congenital (genetic or structural) abnormality of the hypothalamus/pituitary/CNS. Renewal for pediatric GHD, TS, SGA, and adult GHD: Patient is experiencing improvement.</p>
-----------------------	---

HAEGARDA

Products Affected

- Haegarda

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hereditary angioedema: The requested drug is being used for the prevention of acute angioedema attacks. Patient has hereditary angioedema (HAE) with C1 inhibitor deficiency or dysfunction confirmed by laboratory testing OR patient has hereditary angioedema with normal C1 inhibitor confirmed by laboratory testing. For patients with HAE with normal C1 inhibitor, EITHER 1) Patient tested positive for an F12, angiopoietin-1, plasminogen, kininogen-1 (KNG1), heparan sulfate-glucosamine 3-O-sulfotransferase 6 (HS3ST6), or myoferlin (MYOF) gene mutation OR 2) Patient has a family history of angioedema and the angioedema was refractory to a trial of high-dose antihistamine therapy for at least one month.
Age Restrictions	6 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
Coverage Duration	Plan Year
Other Criteria	N/A

HARVONI

Products Affected

- Harvoni

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hepatitis C virus (HCV): Infection confirmed by presence of HCV RNA in the serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of human immunodeficiency virus (HIV) coinfection, presence or absence of resistance-associated substitutions where applicable, transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current American Association for the Study of Liver Diseases (AASLD) treatment guidelines.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Criteria applied consistent w/ current AASLD-IDSA guidance. Reminder for 8wk option if appropriate.
Other Criteria	N/A

HETLIOZ

Products Affected

- HetlioZ

- Tasimelteon

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For Non-24-Hour Sleep-Wake Disorder: 1) for initial therapy and continuation of therapy: a) diagnosis of total blindness in both eyes (e.g., nonfunctioning retinas) and b) unable to perceive light in either eye, AND 2) if currently on therapy with the requested drug, patient must meet at least one of the following: a) increased total nighttime sleep or b) decreased daytime nap duration. For nighttime sleep disturbances in Smith-Magenis Syndrome (SMS): 1) for initial therapy and continuation therapy, the patient has a confirmed diagnosis of SMS AND 2) if currently on therapy with the requested drug, the patient experiences improvement in the quality of sleep since starting therapy.
Age Restrictions	Non-24: 18 years of age or older. SMS: 16 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with sleep disorder specialist or neurologist
Coverage Duration	Initiation: 6 Months, Renewal: Plan Year
Other Criteria	N/A

HETLIOZ LQ

Products Affected

- Hetlioz Lq

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For nighttime sleep disturbances in Smith-Magenis Syndrome (SMS): 1) for initial therapy and continuation therapy, the patient has a confirmed diagnosis of SMS AND 2) if currently on therapy with the requested drug, the patient experiences improvement in the quality of sleep since starting therapy.
Age Restrictions	3 to 15 years of age
Prescriber Restrictions	Prescribed by or in consultation with a sleep disorder specialist or neurologist
Coverage Duration	Initiation: 6 Months, Renewal: Plan Year
Other Criteria	N/A

HIGH RISK MEDICATION - 65

Products Affected

- Dicyclomine Hydrochloride INJ
- Dihydroergotamine Mesylate INJ
- Dipyridamole TABS
- Disopyramide Phosphate CAPS
- Guanfacine Er TB24 2MG
- Guanfacine Hydrochloride TABS
- Guanfacine Hydrochloride TB24 1MG, 3MG, 4MG
- Ketorolac Tromethamine TABS
- Methscopolamine Bromide TABS
- Perphenazine/amitriptyline
- Thioridazine Hcl TABS 100MG, 10MG, 25MG, 50MG
- Trimethobenzamide Hydrochloride

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM - ANTICONVULSANTS - 65

Products Affected

- Phenobarbital ELIX 20MG/5ML
- Phenobarbital TABS 100MG, 15MG, 16.2MG, 30MG, 32.4MG, 60MG, 64.8MG, 97.2MG
- Phenobarbital Sodium INJ 130MG/ML, 65MG/ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Epilepsy
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM - HYPNOTICS - 65

Products Affected

- Zaleplon

- Zolpidem Tartrate TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For insomnia: 1) The patient meets one of the following: a) the patient has a contraindication to the non-HRM (non-High Risk Medication) alternative drug doxepin (3 mg or 6 mg) OR b) The non-HRM (non-High Risk Medication) alternative drug doxepin (3 mg or 6 mg) has been tried AND the patient experienced an inadequate treatment response OR intolerance to the non-HRM (non-High Risk Medication) alternative drug doxepin (3 mg or 6 mg) AND 2) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient AND 3) If the patient is using two or more additional central nervous system (CNS) active medications (e.g., lorazepam, quetiapine, sertraline, clonazepam, escitalopram, alprazolam) with the requested drug, the prescriber has determined that taking multiple central nervous system (CNS) active medications is medically necessary for the patient [Note: Use of multiple central nervous system (CNS) active medications in older adults is associated with an increased risk of falls.].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) APPLIES TO GREATER THAN CUMULATIVE 90 DAYS OF THERAPY PER YEAR.

HRM - SCOPOLAMINE - 65

Products Affected

- Scopolamine PT72 1MG/3DAYS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Excessive salivation
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM-ANTIPARKINSON - 65

Products Affected

- Benztropine Mesylate TABS
- Trihexyphenidyl Hcl SOLN
- Trihexyphenidyl Hydrochloride

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. EPS (extrapyramidal symptoms): 1) The patient has not tried the non-HRM alternative drug amantadine AND 2) The patient has a contraindication to the non-HRM alternative drug amantadine OR 3) The patient has tried the non-HRM alternative drug amantadine AND 4) The patient experienced an inadequate treatment response OR intolerance to the non-HRM alternative drug amantadine. Parkinson's: 1) The patient has tried two of the following non-HRM alternative drugs: amantadine, carbidopa/levodopa, pramipexole, or ropinirole. AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: amantadine, carbidopa/levodopa, pramipexole, or ropinirole.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM-CYPROHEPTADINE - 65

Products Affected

- Cyproheptadine Hcl SYRP
- Cyproheptadine Hydrochloride TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Pruritus, spasticity due to spinal cord injury
Exclusion Criteria	N/A
Required Medical Information	The prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. For rhinitis: 1) The patient has tried two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM-HYDROXYZINE - 65

Products Affected

- Hydroxyzine Hcl TABS 50MG
- Hydroxyzine Hydrochloride SYRP
- Hydroxyzine Hydrochloride TABS 10MG, 25MG
- Hydroxyzine Pamoate CAPS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For anxiety: 1) The patient has tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline, or venlafaxine extended-release AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline, or venlafaxine extended-release OR 3) The patient has not tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release AND 4) The patient has acute anxiety. For all indications: 1) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. AND 2) If the patient is taking one or more additional anticholinergic medications (e.g., oxybutynin, meclizine, paroxetine, amitriptyline, dicyclomine, cyclobenzaprine) with the requested drug, the prescriber has determined that taking multiple anticholinergic medications is medically necessary for the patient [Note: Use of multiple anticholinergic medications in older adults is associated with an increased risk of cognitive decline.].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	This Prior Authorization only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)
-----------------------	--

HRM-HYDROXYZINE INJ - 65

Products Affected

- Hydroxyzine Hcl INJ 25MG/ML

- Hydroxyzine Hydrochloride INJ

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. For alcohol withdrawal syndrome: 1) The patient has not tried one of the following alternative drugs: clorazepate or lorazepam AND 2) The patient has a contraindication to one of the following alternative drugs: clorazepate or lorazepam OR 3) The patient has tried one of the following alternative drugs: clorazepate or lorazepam AND 4) The patient experienced an inadequate treatment response OR intolerance to one of the following alternative drugs: clorazepate or lorazepam. For anxiety: 1) The patient has tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release OR 3) The patient has not tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release AND 4) The patient has acute anxiety.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	<p>This Prior Authorization only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)</p>

HRM-PROMETHAZINE - 65

Products Affected

- Promethazine Hcl INJ
- Promethazine Hcl SUPP 12.5MG, 25MG
- Promethazine Hcl TABS 12.5MG
- Promethazine Hcl Plain
- Promethazine Hydrochloride TABS 25MG, 50MG
- Promethegan

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. For rhinitis: 1) The patient has tried two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRMS - CARBINOXAMINE

Products Affected

- Carbinoxamine Maleate SOLN
- Carbinoxamine Maleate TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. For rhinitis: 1) The patient has tried two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRMS - CLEMASTINE- 65

Products Affected

- Clemastine Fumarate TABS 2.68MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. For rhinitis: 1) The patient has tried two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRMS - TCAS

Products Affected

- Protriptyline Hcl

- Trimipramine Maleate CAPS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For depression: 1) The patient tried two of the following alternative drugs: SSRIs (selective serotonin reuptake inhibitors), SNRIs (serotonin-norepinephrine reuptake inhibitors), bupropion, mirtazapine, or trazodone AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following alternative drugs: SSRIs (selective serotonin reuptake inhibitors), SNRIs (serotonin-norepinephrine reuptake inhibitors), bupropion, mirtazapine, or trazodone AND 3) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM-SKELETAL MUSCLE RELAXANTS - 65

Products Affected

- Chlorzoxazone TABS 500MG
- Cyclobenzaprine Hydrochloride TABS 10MG, 5MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient AND 2) If the patient is using one or more additional anticholinergic medications (e.g., oxybutynin, meclizine, paroxetine, amitriptyline, dicyclomine, hydroxyzine) with the requested drug, the prescriber has determined that taking multiple anticholinergic medications is medically necessary for the patient [Note: Use of multiple anticholinergic medications in older adults is associated with an increased risk of cognitive decline.].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	This Prior Authorization only applies to patients 70 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.). APPLIES TO GREATER THAN CUMULATIVE 90 DAYS OF THERAPY PER YEAR.

HUMIRA

Products Affected

- Humira INJ 10MG/0.1ML, 20MG/0.2ML, 40MG/0.4ML, 40MG/0.8ML
- Humira Pediatric Crohns Disease Starter Pack INJ 0, 80MG/0.8ML
- Humira Pen
- Humira Pen-cd/uc/hs Starter
- Humira Pen-pediatric Uc Starter Pack
- Humira Pen-ps/uv Starter

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Axial spondyloarthritis, Behcet's syndrome
Exclusion Criteria	N/A
Required Medical Information	<p>For moderately to severely active rheumatoid arthritis (new starts only): 1) Inadequate response, intolerance or contraindication to methotrexate (MTX) OR 2) Inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. For active ankylosing spondylitis and axial spondyloarthritis (new starts only): 1) Inadequate response to a non-steroidal anti-inflammatory drug (NSAID) trial OR 2) Intolerance or contraindication to NSAIDs. For moderate to severe plaque psoriasis (new starts only): 1) At least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) Patient meets any of the following: a) Patient has experienced an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, OR b) Pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, OR c) Patient has severe psoriasis that warrants a biologic as first-line therapy (i.e., at least 10% of the BSA or crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected). For moderately to severely active Crohn's disease (new starts only): 1) Inadequate response to at least one conventional therapy (e.g., corticosteroids), OR 2) Intolerance or contraindication to conventional therapy. For moderately to severely active ulcerative colitis (new starts only): 1) Inadequate response to at least one conventional therapy (e.g., corticosteroids), OR 2) Intolerance or contraindication to conventional therapy.</p>
Age Restrictions	N/A

Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

HYPNOTIC-BENZODIAZEPINES - 65

Products Affected

- Triazolam TABS 0.125MG
- Temazepam CAPS 15MG, 22.5MG, 7.5MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge the benefit of therapy with the requested drug outweighs the potential risks to the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to doxepin (3 mg or 6 mg).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization requirement applies only to patients 70 years of age or older. APPLIES TO GREATER THAN CUMULATIVE 90 DAYS OF THERAPY PER YEAR.

IBRANCE

Products Affected

- Ibrance

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Unresectable well-differentiated/dedifferentiated liposarcoma of the retroperitoneum, recurrent hormone receptor-positive human epidermal growth factor receptor 2 (HER2)-negative breast cancer
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ICATIBANT

Products Affected

- Icatibant Acetate

- Sajazir

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hereditary angioedema (HAE): The requested drug is being used for the treatment of acute angioedema attacks. Patient has HAE with C1 inhibitor deficiency or dysfunction confirmed by laboratory testing OR patient has HAE with normal C1 inhibitor confirmed by laboratory testing. For patients with HAE with normal C1 inhibitor, EITHER 1) Patient tested positive for an F12, angiotensin-1, plasminogen, kininogen-1 (KNG1), heparan sulfate-glucosaminase 3-O-sulfotransferase 6 (HS3ST6), or myoferlin (MYOF) gene mutation OR 2) Patient has a family history of angioedema and the angioedema was refractory to a trial of high-dose antihistamine therapy for at least one month.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
Coverage Duration	Plan Year
Other Criteria	N/A

ICLUSIG

Products Affected

- Iclusig

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Myeloid and/or lymphoid neoplasms with eosinophilia and FGFR1 or ABL1 rearrangement in the chronic phase or blast phase
Exclusion Criteria	N/A
Required Medical Information	For chronic myeloid leukemia (CML) or acute lymphoblastic leukemia (ALL), including patients who have received a hematopoietic stem cell transplant: diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene. For CML, including patients who have received a hematopoietic stem cell transplant: 1) patient has accelerated or blast phase CML and no other kinase inhibitor is indicated, OR 2) patient has chronic phase CML and has experienced resistance or intolerance to at least 2 prior kinase inhibitors AND at least one of those was imatinib or dasatinib, OR 3) patient is positive for the T315I mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

IDHIFA

Products Affected

- Idhifa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Newly-diagnosed acute myeloid leukemia
Exclusion Criteria	N/A
Required Medical Information	For acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation: 1) patient is 60 years of age or older with newly-diagnosed AML and meets one of the following: a) patient is not a candidate for intensive induction therapy, or b) patient declines intensive induction chemotherapy, OR 2) patient is 60 years of age or older and the requested drug will be used as post-induction therapy following response to induction therapy with the requested drug OR 3) patient has relapsed or refractory AML.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

IMATINIB

Products Affected

- Imatinib Mesylate

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Desmoid tumors, pigmented villonodular synovitis/tenosynovial giant cell tumor (PVNS/TGCT), recurrent chordoma, melanoma, Kaposi sarcoma, chronic myelomonocytic leukemia, chronic graft versus host disease (cGVHD), T-cell acute lymphoblastic leukemia with ABL-class translocation, aggressive systemic mastocytosis for well-differentiated systemic mastocytosis (WDSM) or when eosinophilia is present with FIP1L1-PDGFR A fusion gene, myeloid and/or lymphoid neoplasms with eosinophilia and ABL1, FIP1L1-PDGFR A, or PDGFR B rearrangement in the chronic phase or blast phase
Exclusion Criteria	N/A
Required Medical Information	For chronic myeloid leukemia (CML) or Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL), including patients who have received a hematopoietic stem cell transplant: diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene. For CML: patient did not fail (excluding failure due to intolerance) prior therapy with a tyrosine kinase inhibitor. For melanoma: c-Kit mutation is positive.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

IMBRUVICA

Products Affected

- Imbruvica

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Hairy cell leukemia, lymphoplasmacytic lymphoma, primary central nervous system (CNS) lymphoma, AIDS-related B-cell lymphoma, diffuse large B-cell lymphoma, post-transplant lymphoproliferative disorders, high-grade B-cell lymphoma, mantle cell lymphoma, marginal zone lymphoma (including extranodal marginal zone lymphoma of the stomach, extranodal marginal zone lymphoma of nongastric sites, nodal marginal zone lymphoma, splenic marginal zone lymphoma)
Exclusion Criteria	N/A
Required Medical Information	For mantle cell lymphoma: 1) the requested drug will be used as second-line or subsequent therapy, OR 2) the requested drug will be used in combination with rituximab as pretreatment to induction therapy with RHyperCVAD (rituximab, cyclophosphamide, vincristine, doxorubicin, and dexamethasone) regimen, OR 3) the requested drug will be used as aggressive induction therapy. For marginal zone lymphoma (including extranodal marginal zone lymphoma of the stomach, extranodal marginal zone lymphoma of nongastric sites, nodal marginal zone lymphoma, and splenic marginal zone lymphoma): the requested drug will be used as second-line or subsequent therapy. For hairy cell leukemia: the requested drug will be used as a single agent for disease progression. For primary CNS lymphoma: 1) the disease is relapsed or refractory OR 2) the requested drug is used for induction therapy as a single agent. For diffuse large B-cell lymphoma and high-grade B-cell lymphoma: the requested drug will be used as second-line or subsequent therapy. For AIDS-related B-cell lymphoma: the requested drug will be used as a single agent and as second-line or subsequent therapy for relapsed disease. For post-transplant lymphoproliferative disorders: the requested drug will be used in patients who have received prior chemoimmunotherapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A

Coverage Duration	Plan Year
Other Criteria	N/A

INBRIJA

Products Affected

- Inbrija

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For initial treatment of off episodes in Parkinson's disease: 1) The patient is currently being treated with oral carbidopa/levodopa. 2) Patient does not have any of the following: asthma, chronic obstructive pulmonary disease (COPD), or other chronic underlying lung disease. For continuation treatment of off episodes in Parkinson's disease: The patient is experiencing improvement on the requested drug.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

INCRELEX

Products Affected

- Increlex

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Pediatric patients with closed epiphyses
Required Medical Information	For growth failure due to severe primary insulin-like growth factor-1 (IGF-1) deficiency or growth hormone (GH) gene deletion in patients who have developed neutralizing antibodies to GH, patient meets all of the following prior to beginning therapy with the requested drug (new starts only): 1) height 3 or more standard deviations (SD) below the mean for children of the same age and gender AND 2) basal IGF-1 level 3 or more SD below the mean for children of the same age and gender AND 3) provocative growth hormone test showing a normal or elevated growth hormone level. For growth failure due to severe primary IGF-1 deficiency or GH gene deletion in patients who have developed neutralizing antibodies to GH, continuation of therapy: patient is experiencing improvement.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

INLYTA

Products Affected

- Inlyta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Thyroid carcinoma (papillary, Hurthle cell, or follicular)
Exclusion Criteria	N/A
Required Medical Information	For renal cell carcinoma: The disease is advanced, relapsed, or stage IV.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

INQOVI

Products Affected

- Inqovi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

INREBIC

Products Affected

- Inrebic

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and janus kinase 2 (JAK2) rearrangement, accelerated phase myelofibrosis, blast phase myelofibrosis/acute myeloid leukemia
Exclusion Criteria	N/A
Required Medical Information	For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and JAK2 rearrangement: the disease is in chronic or blast phase.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

IR BEFORE ER

Products Affected

- Hydrocodone Bitartrate Er T24A
- Hysingla Er
- Methadone Hcl INJ
- Methadone Hcl ORAL SOLN
- Methadone Hcl TABS
- Methadone Hydrochloride CONC
- Morphine Sulfate Er CP24 100MG, 10MG, 120MG, 20MG, 30MG, 45MG, 50MG, 60MG, 75MG, 80MG, 90MG
- Morphine Sulfate Er TBCR
- Tramadol Hcl Er TB24
- Tramadol Hydrochloride Er

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug is being prescribed for pain associated with cancer, sickle cell disease, a terminal condition, or pain being managed through palliative care OR the patient meets all of the following: 1) The requested drug is being prescribed for pain severe enough to require daily, around-the-clock, long-term treatment in a patient who has been taking an opioid AND 2) The patient can safely take the requested dose based on their history of opioid use [Note: This drug should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.] AND 3) The patient has been evaluated and the patient will be monitored for the development of opioid use disorder AND 4) This request is for continuation of therapy for a patient who has been receiving an extended-release opioid agent for at least 30 days OR the patient has taken an immediate-release opioid for at least one week.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

IRESSA

Products Affected

- Gefitinib

- Iressa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Sensitizing epidermal growth factor receptor (EGFR) mutation-positive recurrent non-small cell lung cancer (NSCLC).
Exclusion Criteria	N/A
Required Medical Information	For non-small cell lung cancer (NSCLC): 1) disease must be metastatic, advanced, or recurrent AND 2) patient must have a sensitizing EGFR mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ISOTRETINOIN

Products Affected

- Accutane
- Amnesteem
- Claravis
- Isotretinoin CAPS
- Zenatane

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Refractory acne vulgaris, severe refractory rosacea, neuroblastoma, cutaneous T-cell lymphoma (CTCL) (e.g., mycosis fungoides, Sezary syndrome), high risk for developing skin cancer (squamous cell cancers), transient acantholytic dermatosis (Grover's Disease), keratosis follicularis (Darier Disease), lamellar ichthyosis, pityriasis rubra pilaris.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ITRACONAZOLE

Products Affected

- Itraconazole CAPS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Coccidioidomycosis, Coccidioidomycosis prophylaxis in HIV infection, Cryptococcosis, Microsporidiosis, Talaromycosis (formerly Penicilliosis), Histoplasmosis prophylaxis in HIV infection, Invasive fungal infection prophylaxis in liver transplant, chronic granulomatous disease (CGD), and hematologic malignancy, Sporotrichosis, Pityriasis versicolor, Tinea versicolor, Tinea corporis, Tinea cruris, Tinea capitis, Tinea manuum, Tinea pedis.
Exclusion Criteria	N/A
Required Medical Information	The requested drug will be used orally. For the treatment of onychomycosis due to dermatophytes (Tinea unguium), the diagnosis has been confirmed by a fungal diagnostic test (e.g., potassium hydroxide [KOH] preparation, fungal culture, or nail biopsy).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Disseminated/CNS histoplasmosis, Histoplasmosis/Coccidioidomycosis/CGD ppx: 12 mths. Others: 6 mths
Other Criteria	N/A

IVERMECTIN TAB

Products Affected

- Ivermectin TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Ascariasis, Cutaneous larva migrans, Mansonelliasis, Scabies, Gnathostomiasis, Pediculosis
Exclusion Criteria	N/A
Required Medical Information	The requested drug is not being prescribed for the prevention or treatment of coronavirus disease 2019 (COVID-19).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 month
Other Criteria	N/A

IVIG

Products Affected

- Bivigam INJ 10%, 5GM/50ML
- Flebogamma Dif
- Gammagard Liquid
- Gammagard S/d Iga Less Than 1mcg/ml
- Gammaked INJ 10GM/100ML, 1GM/10ML, 20GM/200ML, 5GM/50ML
- Gammaplex INJ 10GM/100ML, 10GM/200ML, 20GM/200ML, 20GM/400ML, 5GM/100ML, 5GM/50ML
- Gamunex-c
- Octagam INJ 10GM/100ML, 10GM/200ML, 2.5GM/50ML, 20GM/200ML, 25GM/500ML, 2GM/20ML, 30GM/300ML, 5GM/100ML, 5GM/50ML
- Panzyga
- Privigen

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For B-cell chronic lymphocytic leukemia (CLL): 1) serum IgG less than 500 mg/dL OR 2) a history of recurrent bacterial infections. For bone marrow transplant/hematopoietic stem cell transplant (BMT/HSCT): 1) IVIG is requested within the first 100 days post-transplant OR 2) serum IgG less than 400 mg/dL. For pediatric human immunodeficiency virus (HIV) infection: 1) serum IgG less than 400 mg/dL OR 2) history of recurrent bacterial infections. For dermatomyositis and polymyositis: 1) at least one standard first-line treatment (corticosteroid or immunosuppressant) has been tried but was unsuccessful or not tolerated OR 2) patient is unable to receive standard therapy because of a contraindication or other clinical reason. For pure red cell aplasia (PRCA): PRCA is secondary to parvovirus B19 infection.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.
-----------------------	--

JAKAFI

Products Affected

- Jakafi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Lower-risk myelofibrosis, accelerated phase myelofibrosis, blast phase myelofibrosis/acute myeloid leukemia, acute lymphoblastic leukemia (ALL), chronic myelomonocytic leukemia (CMML)-2, BCR-ABL negative atypical chronic myeloid leukemia (aCML), essential thrombocythemia, and myeloid, lymphoid or mixed lineage neoplasms with eosinophilia and JAK2 rearrangement
Exclusion Criteria	N/A
Required Medical Information	For polycythemia vera: patient had an inadequate response or intolerance to interferon therapy or hydroxyurea. For acute lymphoblastic leukemia: patient has a cytokine receptor-like factor 2 (CRLF2) mutation or a mutation associated with activation of the Janus kinase/signal transducers and activators of transcription (JAK/STAT) pathway. For CMML-2: the requested drug is used in combination with a hypomethylating agent. For BCR-ABL negative aCML: the requested drug is used as a single agent or in combination with a hypomethylating agent. For essential thrombocythemia: patient had an inadequate response or loss of response to hydroxyurea, interferon therapy, or anagrelide. For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and JAK2 rearrangement: the disease is in chronic or blast phase.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

JAYPIRCA

Products Affected

- Jaypirca

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KALYDECO

Products Affected

- Kalydeco PACK 13.4MG, 25MG, 50MG, 75MG
- Kalydeco TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For cystic fibrosis (CF): The requested medication will not be used in combination with other medications containing ivacaftor.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KESIMPTA

Products Affected

- Kesimpta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KETOCONAZOLE

Products Affected

- Ketoconazole TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Cushing's syndrome
Exclusion Criteria	Acute or chronic liver disease. Concurrent use with drugs that are contraindicated with ketoconazole tablets: dofetilide, quinidine, pimozide, cisapride, methadone, disopyramide, dronedarone, ranolazine, ergot alkaloids, irinotecan, lurasidone, oral midazolam, alprazolam, triazolam, felodipine, nisoldipine, tolvaptan, eplerenone, lovastatin, simvastatin, or colchicine.
Required Medical Information	The potential benefits outweigh the risks of treatment with oral ketoconazole. For systemic fungal infections, the patient has any of the following diagnoses: blastomycosis, coccidioidomycosis, histoplasmosis, chromomycosis, or paracoccidioidomycosis. For Cushing's syndrome: the requested drug is being prescribed for a patient who cannot tolerate surgery or where surgery has not been curative.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

KEVZARA

Products Affected

- Kevzara

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderately to severely active rheumatoid arthritis (new starts only): 1) Inadequate response, intolerance or contraindication to methotrexate (MTX) OR 2) Inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. For polymyalgia rheumatica (PMR) (new starts only): 1) The patient has experienced an inadequate treatment response to corticosteroids OR 2) The patient has experienced a disease flare while attempting to taper corticosteroids.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KISQALI

Products Affected

- Kisqali
- Kisqali Femara 200 Dose
- Kisqali Femara 400 Dose
- Kisqali Femara 600 Dose

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent hormone receptor-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer, in combination with an aromatase inhibitor, or fulvestrant.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KORLYM

Products Affected

- Korlym

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	Plan Year
Other Criteria	N/A

KRAZATI

Products Affected

- Krazati

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KRISTALOSE

Products Affected

- Kristalose

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1) The patient has experienced an inadequate treatment response to a one month trial of generic lactulose solution, 2) The patient has experienced an intolerance that would prohibit a one month trial generic lactulose solution, or 3) the patient has a contraindication to an inactive ingredient which is not contained in the requested drug.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KUVAN

Products Affected

- Javygtor

- Sapropterin Dihydrochloride

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For phenylketonuria (PKU): For patients who have not yet received a therapeutic trial of the requested drug, the patient's pretreatment, including before dietary management, phenylalanine level is greater than 6 mg/dL (360 micromol/L). For patients who completed a therapeutic trial of the requested drug, the patient must have experienced improvement (e.g., reduction in blood phenylalanine levels, improvement in neuropsychiatric symptoms).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial: 2 months. All others: Plan Year.
Other Criteria	N/A

LAPATINIB

Products Affected

- Lapatinib Ditosylate

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Brain metastases from human epidermal growth factor receptor 2 (HER2)-positive breast cancer, recurrent HER2-positive breast cancer, recurrent epidermal growth factor receptor (EGFR)-positive chordoma, HER2-amplified and RAS and BRAF wild-type colorectal cancer in combination with trastuzumab.
Exclusion Criteria	N/A
Required Medical Information	For breast cancer, the patient meets all the following: a) the disease is recurrent, advanced, or metastatic (including brain metastases), b) the disease is human epidermal growth factor receptor 2 (HER2)-positive, c) the requested drug will be used in combination with any of the following: 1) aromatase inhibitor, 2) capecitabine, OR 3) trastuzumab.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LENVIMA

Products Affected

- Lenvima 10 Mg Daily Dose
- Lenvima 12mg Daily Dose
- Lenvima 14 Mg Daily Dose
- Lenvima 18 Mg Daily Dose
- Lenvima 20 Mg Daily Dose
- Lenvima 24 Mg Daily Dose
- Lenvima 4 Mg Daily Dose
- Lenvima 8 Mg Daily Dose

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Medullary thyroid carcinoma, recurrent endometrial carcinoma, thymic carcinoma
Exclusion Criteria	N/A
Required Medical Information	For differentiated thyroid cancer (follicular, papillary, or Hurthle cell): disease is not amenable to radioactive iodine therapy and unresectable, locally recurrent, persistent, or metastatic. For hepatocellular carcinoma: disease is unresectable or inoperable, local, metastatic or with extensive liver tumor burden. For renal cell carcinoma, the disease is advanced, relapsed, or stage IV. For endometrial carcinoma, the patient meets ALL of the following: 1) The disease is advanced, recurrent, or metastatic, 2) The patient experienced disease progression following prior systemic therapy, AND 3) The patient is not a candidate for curative surgery or radiation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LEUPROLIDE

Products Affected

- Leuprolide Acetate INJ 1MG/0.2ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Use in combination with growth hormone for children with growth failure and advancing puberty, recurrent androgen receptor positive salivary gland tumors
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LIDOCAINE PATCHES

Products Affected

- Lidocaine PTCH 5%

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Pain associated with diabetic neuropathy, pain associated with cancer-related neuropathy (including treatment-related neuropathy [e.g., neuropathy associated with radiation treatment or chemotherapy]).
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LINEZOLID

Products Affected

- Linezolid

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The infection is proven or strongly suspected to be caused by susceptible bacteria based on: 1) culture and susceptibility information OR 2) local epidemiology and susceptibility patterns. The patient will be using the requested drug orally or intravenously.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed or directed by an Infectious Disease specialist when being converted from intravenous (IV) linezolid (Zyvox)
Coverage Duration	28 days
Other Criteria	N/A

LONSURF

Products Affected

- Lonsurf

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For colorectal cancer: The disease is advanced or metastatic. For gastric or gastroesophageal junction adenocarcinoma, all of the following criteria must be met: 1) The disease is unresectable locally advanced, recurrent, or metastatic, and 2) The patient has been previously treated with at least two prior lines of chemotherapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LORBRENA

Products Affected

- Lorbrena

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Anaplastic lymphoma kinase (ALK)-positive recurrent non-small cell lung cancer (NSCLC), repressor of silencing (ROS)-1 rearrangement-positive recurrent, advanced, or metastatic NSCLC.
Exclusion Criteria	N/A
Required Medical Information	For recurrent, advanced, or metastatic non-small cell lung cancer: 1) Disease is anaplastic lymphoma kinase (ALK)-positive OR 2) Disease is positive for repressor of silencing (ROS)-1 rearrangement and the requested drug is being used following disease progression on crizotinib, entrectinib, or ceritinib.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LUMAKRAS

Products Affected

- Lumakras

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LUPRON PED

Products Affected

- Lupron Depot-ped INJ 45MG
- Lupron Depot-ped (1-month)
- Lupron Depot-ped (3-month)

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For central precocious puberty (CPP), patients not currently receiving therapy must meet all of the following criteria: 1) Diagnosis of CPP was confirmed by a pubertal response to a gonadotropin releasing hormone (GnRH) agonist test OR a pubertal level of a third generation luteinizing hormone (LH) assay, 2) Assessment of bone age versus chronological age supports the diagnosis of CPP, and 3) The onset of secondary sexual characteristics occurred prior to 8 years of age for female patients OR prior to 9 years of age for male patients.
Age Restrictions	CPP: Patient must be less than 12 years old if female and less than 13 years old if male.
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LUPRON-ENDOMETRIOSIS

Products Affected

- Lupron Depot (1-month) INJ 3.75MG
- Lupron Depot (3-month) INJ 11.25MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Breast cancer, epithelial ovarian cancer/fallopian tube cancer/primary peritoneal cancer
Exclusion Criteria	N/A
Required Medical Information	For uterine fibroids, patient must meet one of the following: 1) Diagnosis of anemia (e.g., hematocrit less than or equal to 30 percent and/or hemoglobin less than or equal to 10g/dL), OR 2) the requested medication will be used prior to surgery for uterine fibroids. For breast cancer, the requested drug is used for hormone receptor (HR)-positive disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Fibroids: 3 months (mo), max 6 mo total. Endometriosis: 6 mo, max 12 mo total. Others: Plan Year
Other Criteria	N/A

LYNPARZA

Products Affected

- Lynparza TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent HER2-negative, BRCA 1/2-germline mutated breast cancer, recurrent or metastatic HER2-positive, BRCA 1/2-germline mutated breast cancer, uterine leiomyosarcoma.
Exclusion Criteria	N/A
Required Medical Information	For recurrent or metastatic breast cancer: the disease is BRCA 1/2-germline mutated. For prostate cancer: 1) The patient has a BRCA mutation and the requested drug will be used in combination with abiraterone and either prednisone or prednisolone OR 2) The patient has progressed on prior treatment with an androgen receptor-directed therapy. For epithelial ovarian, fallopian tube, or primary peritoneal cancer: The requested drug is used for maintenance therapy for stage II-IV or recurrent disease who are in complete or partial response to chemotherapy. For uterine leiomyosarcoma: 1) the requested drug is used as second-line therapy AND 2) the patient has BRCA-altered disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LYRICA CR

Products Affected

- Pregabalin Er

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to gabapentin.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LYTGOBI

Products Affected

- Lytgobi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Extrahepatic cholangiocarcinoma
Exclusion Criteria	N/A
Required Medical Information	For cholangiocarcinoma: 1) patient has a diagnosis of unresectable, locally advanced or metastatic cholangiocarcinoma, 2) patient has received a previous treatment, AND 3) patient disease has a fibroblast growth factor receptor 2 (FGFR2) gene fusion or other rearrangement.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

MAVYRET

Products Affected

- Mavyret

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Decompensated cirrhosis/moderate or severe hepatic impairment (Child Turcotte Pugh [CTP] class B or C).
Required Medical Information	For hepatitis C virus (HCV): Infection confirmed by presence of HCV RNA in the serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [CTP class B or C]), presence or absence of human immunodeficiency virus (HIV) coinfection, presence or absence of resistance-associated substitutions where applicable, transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current American Association for the Study of Liver Diseases (AASLD) treatment guidelines.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Criteria will be applied consistent with current AASLD-IDSA guidance
Other Criteria	N/A

MEKINIST

Products Affected

- Mekinist

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Brain metastases from melanoma, uveal melanoma, central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma), low grade serous ovarian cancer, ovarian borderline epithelial tumors (low malignant potential) with invasive implants, Langerhans cell histiocytosis, Erdheim-Chester disease, Rosai-Dorfman disease, gallbladder cancer, intrahepatic cholangiocarcinoma, extrahepatic cholangiocarcinoma
Exclusion Criteria	N/A
Required Medical Information	For adjuvant treatment of melanoma,: 1) The tumor is positive for a BRAF V600 activating mutation (e.g., V600E or V600K), and 2) The requested drug will be used in combination with dabrafenib. For unresectable or metastatic melanoma: 1) The tumor is positive for a BRAF V600 activating mutation (e.g., V600E or V600K), and 2) The requested drug will be used as a single agent or in combination with dabrafenib. For brain metastases from melanoma, central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma), non-small cell lung cancer, solid tumors, and anaplastic thyroid cancer: 1) The tumor is positive for a BRAF V600E mutation, and 2) The requested drug will be used in combination with dabrafenib. For uveal melanoma, the requested drug will be used as a single agent. For low grade serous ovarian cancer and ovarian borderline epithelial tumors (low malignant potential) with invasive implants: The requested drug will be used to treat persistent or recurrent disease. For gallbladder cancer, intrahepatic cholangiocarcinoma, and extrahepatic cholangiocarcinoma: 1) The tumor is positive for a BRAF V600E mutation, 2) the disease is unresectable or metastatic, and 3) The requested drug will be used in combination with dabrafenib.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	N/A
-----------------------	-----

MEKTOVI

Products Affected

- Mektovi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Adjuvant systemic therapy for cutaneous melanoma
Exclusion Criteria	N/A
Required Medical Information	For cutaneous melanoma: The patient must meet all of the following criteria: 1) Tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K), 2) The requested drug will be used in combination with encorafenib, and 3) The requested drug will be used for either of the following: a) unresectable or metastatic disease, or b) adjuvant systemic therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

MEMANTINE

Products Affected

- Memantine Hcl Titration Pak
- Memantine Hydrochloride SOLN 2MG/ML
- Memantine Hydrochloride TABS
- Memantine Hydrochloride Er

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This edit only applies to patients less than 30 years of age.

MEPRON

Products Affected

- Atovaquone SUSP

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Babesiosis, Toxoplasmosis, Pneumocystis jirovecii pneumonia prophylaxis in pediatric patients, mild-to-moderate Pneumocystis jirovecii pneumonia treatment in pediatric patients.
Exclusion Criteria	N/A
Required Medical Information	For the treatment of mild-to-moderate Pneumocystis jirovecii pneumonia (PCP): the patient had an intolerance or has a contraindication to sulfamethoxazole/trimethoprim (SMX-TMP). For the prevention of PCP and primary toxoplasmosis prophylaxis indications: A) the patient had an intolerance or has a contraindication to SMX-TMP, AND B) the patient has had a CD4 cell count of less than 200 cells per cubic millimeter within the past 3 months. For secondary toxoplasmosis prophylaxis: the patient has had a CD4 cell count of less than 200 cells per cubic millimeter within the past 6 months. For babesiosis treatment: the requested drug is used concurrently with azithromycin.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Secondary toxoplasmosis prophylaxis: 6 months, All other indications: 3 months
Other Criteria	N/A

MODAFINIL

Products Affected

- Modafinil

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1) The patient has a diagnosis of narcolepsy and the diagnosis is confirmed by sleep lab evaluation OR 2) The patient has a diagnosis of Shift Work Disorder (SWD) OR 3) The patient has a diagnosis of obstructive sleep apnea (OSA) and the diagnosis is confirmed by polysomnography
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NATPARA

Products Affected

- Natpara

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Acute postsurgical hypoparathyroidism (within 6 months of surgery) and expected recovery from hypoparathyroidism.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NAYZILAM

Products Affected

- Nayzilam

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	12 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NERLYNX

Products Affected

- Nerlynx

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent human epidermal growth factor receptor 2 (HER2)-positive breast cancer, brain metastases from HER2-positive breast cancer.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NEXAVAR

Products Affected

- Nexavar

- Sorafenib Tosylate TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Acute myeloid leukemia, soft tissue sarcoma (angiosarcoma, desmoid tumors/aggressive fibromatosis, and solitary fibrous tumor subtypes), gastrointestinal stromal tumor, medullary thyroid carcinoma, osteosarcoma, recurrent chordoma, epithelial ovarian cancer, fallopian tube cancer, primary peritoneal cancer, lymphoid, myeloid, or mixed lineage neoplasms with eosinophilia
Exclusion Criteria	N/A
Required Medical Information	For acute myeloid leukemia: the disease is FMS-like tyrosine kinase 3-internal tandem duplication (FLT3-ITD) mutation-positive AND either of the following is met (1 OR 2): 1) the requested drug will be used as maintenance therapy after hematopoietic stem cell transplant, OR 2) the requested drug is used in combination with azacitidine or decitabine for low-intensity treatment induction or post-induction therapy AND either a) the patient has a physiologic age of 60 years of age or older or b) the disease is relapsed/refractory. For thyroid carcinoma: histology is follicular, papillary, Hurthle cell or medullary. For gastrointestinal stromal tumor (GIST): the disease is unresectable, recurrent, or metastatic AND the patient has failed on an FDA-approved therapy (e.g., imatinib, sunitinib, regorafenib, ripretinib). For renal cell carcinoma: the patient meets ALL of the following: 1) The disease is advanced, AND 2) The patient has experienced disease progression or an intolerable adverse event with a trial of cabozantinib or axitinib. For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia: 1) The disease has a FLT3 rearrangement AND 2) The disease is in chronic or blast phase.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	N/A
-----------------------	-----

NINLARO

Products Affected

- Ninlaro

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Relapsed/refractory systemic light chain amyloidosis, Waldenstrom macroglobulinemia, lymphoplasmacytic lymphoma
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NITISINONE

Products Affected

- Nitisinone

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hereditary tyrosinemia type 1 (HT-1): Diagnosis of HT-1 is confirmed by one of the following: 1) biochemical testing (e.g., detection of succinylacetone in urine) or 2) DNA testing (mutation analysis).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NORTHERA

Products Affected

- Droxidopa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For neurogenic orthostatic hypotension (nOH): Prior to initial therapy, patient has a persistent, consistent decrease in systolic blood pressure of at least 20 mmHg OR decrease in diastolic blood pressure of at least 10 mmHg within 3 minutes of standing or head-up tilt test. For continuation of therapy for nOH, patient must experience a sustained reduction in symptoms of nOH (i.e., decrease in dizziness, lightheadedness, or feeling faint). For both initial and continuation of therapy for nOH, the requested drug will be used for patients with neurogenic orthostatic hypotension associated with one of the following diagnoses: 1) primary autonomic failure due to Parkinson's disease, multiple system atrophy, or pure autonomic failure, OR 2) dopamine beta-hydroxylase deficiency, OR 3) non-diabetic autonomic neuropathy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A

NUBEQA

Products Affected

- Nubeqa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NUEDEXTA

Products Affected

- Nuedexta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NUPLAZID

Products Affected

- Nuplazid CAPS
- Nuplazid TABS 10MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hallucinations and delusions associated with Parkinson's disease psychosis, the diagnosis of Parkinson's disease must be made prior to the onset of psychotic symptoms.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NURTEC

Products Affected

- Nurtec

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute migraine treatment: The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to one triptan 5-HT1 receptor agonist . Preventive treatment of migraine, initial: The patient meets either of the following: 1) The patient experienced an inadequate treatment response with a 4-week trial of any one of the following: antiepileptic drugs (AEDs), beta-adrenergic blocking agents, antidepressants OR 2) The patient experienced an intolerance or has a contraindication that would prohibit a 4-week trial of any one of the following: antiepileptic drugs (AEDs), beta-adrenergic blocking agents, antidepressants. Preventive treatment of migraine, continuation: The patient received at least 3 months of treatment with the requested drug, and the patient had a reduction in migraine days per month from baseline.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Preventive treatment of migraine - initial: 3 months, All other indications: Plan Year
Other Criteria	N/A

OCTREOTIDE

Products Affected

- Octreotide Acetate INJ 1000MCG/ML, 100MCG/ML, 200MCG/ML, 500MCG/ML, 50MCG/ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Tumor control of thymomas and thymic carcinomas.
Exclusion Criteria	N/A
Required Medical Information	For acromegaly (initial): 1) patient has a high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range, and 2) patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For acromegaly continuation of therapy: patient's IGF-1 level has decreased or normalized since initiation of therapy. For tumor control of thymomas and thymic carcinomas, the requested drug will be used for any of the following: 1) locally advanced or metastatic disease, 2) postoperatively following tumor resection.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ODOMZO

Products Affected

- Odomzo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

OFEV

Products Affected

- Ofev

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

OJJAARA

Products Affected

- Ojjaara

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ONCASPAR

Products Affected

- Oncaspar

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Extranodal natural killer/T-cell lymphoma, hepatosplenic gamma-delta T-cell lymphoma
Exclusion Criteria	N/A
Required Medical Information	For extranodal natural killer/T-cell lymphoma, acute lymphoblastic leukemia, and hepatosplenic gamma-delta T-cell lymphoma: the requested drug must be used in conjunction with multi-agent chemotherapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ONUREG

Products Affected

- Onureg

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

OPSUMIT

Products Affected

- Opsumit

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): Diagnosis was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) Pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ORACEA

Products Affected

- Doxycycline CPDR
- Oracea

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient has experienced an inadequate treatment response or intolerance to formulary generic topical metronidazole or the patient has a contraindication that would prohibit a trial of formulary generic topical metronidazole.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ORAL-INTRANASAL FENTANYL

Products Affected

- Fentanyl Citrate Oral Transmucosal

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1) The requested drug is indicated for the treatment of breakthrough CANCER-related pain only. The requested drug is being prescribed for the management of breakthrough pain in a CANCER patient with underlying CANCER pain AND 2) The International Classification of Diseases (ICD) diagnosis code provided supports the CANCER-RELATED diagnosis. [Note: For drug coverage approval, ICD diagnosis code provided MUST support the CANCER-RELATED diagnosis.] AND 3) The patient is currently receiving, and will continue to receive, around-the-clock opioid therapy for underlying CANCER pain AND 4) The requested drug is intended only for use in opioid tolerant patients. The patient can safely take the requested dose based on their current opioid use history. [Note: Patients considered opioid tolerant are those who are taking around-the-clock medicine consisting of at least 60 mg of oral morphine per day, at least 25 mcg per hour of transdermal fentanyl, at least 30 mg of oral oxycodone per day, at least 60 mg of oral hydrocodone per day, at least 8 mg of oral hydromorphone per day, at least 25 mg of oral oxymorphone per day, or an equianalgesic dose of another opioid medication daily for one week or longer.].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ORGOVYX

Products Affected

- Orgovyx

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ORKAMBI

Products Affected

- Orkambi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For cystic fibrosis (CF): The requested medication will not be used in combination with other medications containing ivacaftor.
Age Restrictions	1 year of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ORSERDU

Products Affected

- Orserdu

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent hormone receptor positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: 1) the disease is estrogen receptor (ER) positive, human epidermal growth factor receptor 2 (HER2)-negative, and ESR1 mutated AND 2) the patient meets either of the following: a) the disease is advanced, recurrent, or metastatic AND the patient has disease progression following at least one line of endocrine therapy OR b) the disease had no response to preoperative systemic therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

OTEZLA

Products Affected

- Otezla

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For plaque psoriasis (new starts only): Patient meets either of the following: 1) Inadequate treatment response or intolerance to ANY of the following: a) a topical therapy (e.g., topical corticosteroids, calcineurin inhibitors, vitamin D analogs), b) phototherapy (e.g., UVB, PUVA), or c) pharmacologic treatment with methotrexate, cyclosporine, or acitretin, OR 2) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

OXANDROLONE

Products Affected

- Oxandrolone TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Cachexia associated with AIDS (HIV wasting), To enhance growth in patients with Turners Syndrome
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Turners Syndrome: Plan Year, All other diagnoses: 6 months
Other Criteria	Coverage will be denied if request is for an indication excluded from Medicare Part D.

OXAZEPAM - 65

Products Affected

- Oxazepam

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For all indications: The prescriber must acknowledge the benefit of therapy with this prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For the management of anxiety disorders, anxiety associated with depression, and the management of anxiety, tension, agitation and irritability in older patients: 1) The requested drug is being used concurrently with a selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the SSRI/SNRI becomes effective for the symptoms of anxiety, OR 2) The patient has experienced an inadequate treatment response, intolerance, or a has a contraindication to AT LEAST TWO agents from the following classes: a) selective serotonin reuptake inhibitors (SSRIs), b) serotonin-norepinephrine reuptake inhibitors (SNRIs).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Short-term relief anxiety-1 month, Anxiety Disorders-4 months, Alcohol Withdrawal-Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older.

OZEMPIC AET

Products Affected

- Ozempic

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	The Prior Authorization only applies to patients whose claim is not submitted with an ICD-10 code indicating a diagnosis of type 2 diabetes mellitus OR to patients who do not have a history of an antidiabetic drug (EXCLUDING glucagon-like peptide receptor agonists [GLP-1 RAs] and combination glucose-dependent insulintropic polypeptide [GIP] and GLP-1 RAs).

PEGASYS

Products Affected

- Pegasys

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Myeloproliferative neoplasm (essential thrombocythemia, polycythemia vera, symptomatic lower-risk myelofibrosis), systemic mastocytosis, adult T-cell leukemia/lymphoma, mycosis fungoides/sezary syndrome, primary cutaneous CD30+ T-cell lymphoproliferative disorders, hairy cell leukemia, Erdheim-Chester disease.
Exclusion Criteria	N/A
Required Medical Information	For chronic hepatitis C: Hepatitis C virus (HCV) confirmed by presence of hepatitis C virus HCV RNA in serum prior to starting treatment and the planned treatment regimen.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	HCV: 12-48wks. Criteria applied consistent w/current AASLD/IDSA guidance. HBV: 48wks. Other: Plan Yr
Other Criteria	N/A

PEMAZYRE

Products Affected

- Pemazyre

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and FGFR1 rearrangement: the disease is in chronic or blast phase.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PHENYL BUTYRATE

Products Affected

- Sodium Phenylbutyrate POWD 3GM/TSP
- Sodium Phenylbutyrate TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For urea cycle disorders (UCD): Diagnosis of UCD was confirmed by enzymatic, biochemical or genetic testing.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PIQRAY

Products Affected

- Piqray 200mg Daily Dose
- Piqray 250mg Daily Dose
- Piqray 300mg Daily Dose

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, PIK3CA-mutated breast cancer in combination with fulvestrant.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

POMALYST

Products Affected

- Pomalyst

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Relapsed/refractory systemic light chain amyloidosis, primary central nervous system (CNS) lymphoma, POEMS syndrome.
Exclusion Criteria	N/A
Required Medical Information	For multiple myeloma: The patient has previously received at least two prior therapies for multiple myeloma, including an immunomodulatory agent AND a proteasome inhibitor. For Kaposi sarcoma, patient meets one of the following: 1) patient has acquired immunodeficiency syndrome (AIDS), or 2) patient is negative for human immunodeficiency virus (HIV).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PRALUENT

Products Affected

- Praluent

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PREGABALIN

Products Affected

- Pregabalin CAPS

- Pregabalin SOLN

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Cancer-related neuropathic pain, cancer treatment-related neuropathic pain
Exclusion Criteria	N/A
Required Medical Information	For the management of postherpetic neuralgia, the management of neuropathic pain associated with diabetic peripheral neuropathy, cancer-related neuropathic pain, and cancer treatment-related neuropathic pain: The patient has experienced an inadequate treatment response, intolerance, or contraindication to gabapentin.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PRETOMANID

Products Affected

- Pretomanid

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PREVYMIS

Products Affected

- Prevymis TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For prophylaxis of cytomegalovirus (CMV) infection or disease in hematopoietic stem cell transplant (HSCT): 1) the patient is CMV-seropositive, AND 2) the patient is a recipient of an allogeneic HSCT. For prophylaxis of CMV disease in kidney transplant: 1) the patient is CMV-seronegative, AND 2) the patient is a high risk recipient of kidney transplant.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	7 months
Other Criteria	N/A

PROMACTA

Products Affected

- Promacta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>For chronic or persistent immune thrombocytopenia (ITP): 1) For new starts: a) Patient (pt) has had an inadequate response or is intolerant to a prior therapy such as corticosteroids or immunoglobulins, b) Untransfused platelet (plt) count at any point prior to the initiation of the requested medication is less than 30,000/mcL OR 30,000-50,000/mcL with symptomatic bleeding or risk factor(s) for bleeding (e.g., undergoing a medical or dental procedure where blood loss is anticipated, comorbidities such as peptic ulcer disease and hypertension, anticoagulation therapy, profession or lifestyle that predisposes pt to trauma) AND c) For chronic ITP only: pt has had an inadequate response or intolerance to avatrombopag. 2) For continuation of therapy, plt count response to the requested drug: a) Current plt count is less than or equal to 200,000/mcL OR b) Current plt count is greater than 200,000/mcL to less than or equal to 400,000/mcL and dosing will be adjusted to a plt count sufficient to avoid clinically important bleeding. For thrombocytopenia associated with chronic hepatitis C: 1) For new starts: the requested drug is used for initiation and maintenance of interferon-based therapy. 2) For continuation of therapy: pt is receiving interferon-based therapy. For severe aplastic anemia (AA): 1) For new starts: a) Pt will use the requested drug with standard immunosuppressive therapy for first line treatment OR b) the pt had an insufficient response to immunosuppressive therapy. 2) For continuation of therapy : 1) Current plt count is 50,000-200,000/mcL OR 2) Current plt count is less than 50,000/mcL and pt has not received appropriately titrated therapy for at least 16 weeks, OR 3) Current plt count is less than 50,000/mcL and pt is transfusion-independent, OR 4) Current plt count is greater than 200,000/mcL to less than or equal to 400,000/mcL and dosing will be adjusted to achieve and maintain an appropriate target plt count.</p>
Age Restrictions	N/A

Prescriber Restrictions	N/A
Coverage Duration	HCV: 6mo, ITP/AA initial: 6mo, ITP reauth: Plan Year, AA reauth: APR-Plan Year, IPR-16 wks
Other Criteria	APR: adequate platelet response (greater than 50,000/mcL), IPR: inadequate platelet response (less than 50,000/mcL).

PULMOZYME

Products Affected

- Pulmozyme SOLN 2.5MG/2.5ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For cystic fibrosis: Diagnosis of cystic fibrosis was confirmed by appropriate diagnostic or genetic testing.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

QINLOCK

Products Affected

- Qinlock

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

QUETIAPINE XR

Products Affected

- Quetiapine Fumarate Er

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Maintenance monotherapy treatment in bipolar I disorder, monotherapy treatment of generalized anxiety disorder, monotherapy treatment of major depressive disorder
Exclusion Criteria	N/A
Required Medical Information	For schizophrenia, acute treatment of manic or mixed episodes associated with bipolar I disorder, both as monotherapy and as an adjunct to lithium or divalproex, the acute treatment of depressive episodes associated with bipolar disorder, maintenance treatment of bipolar I disorder, as an adjunct to lithium or divalproex, adjunctive treatment of major depressive disorder, or maintenance monotherapy treatment in bipolar I disorder: The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following generic products: A) aripiprazole, B) asenapine, C) olanzapine, D) quetiapine immediate-release, E) risperidone, F) ziprasidone. For all indications: If the patient is 65 years of age or older AND is using two or more additional central nervous system (CNS) active medications (e.g., lorazepam, sertraline, clonazepam, escitalopram, alprazolam, zolpidem) with the requested drug, the prescriber determined that taking multiple central nervous system (CNS) active medications is medically necessary. [Note: Use of multiple central nervous system (CNS) active medications in older adults is associated with an increased risk of falls.].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

QUININE SULFATE

Products Affected

- Quinine Sulfate CAPS 324MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Babesiosis, uncomplicated Plasmodium vivax malaria.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 month
Other Criteria	N/A

REGRANEX

Products Affected

- Regranex

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	20 weeks
Other Criteria	N/A

RETEVMO

Products Affected

- Retevmo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent rearranged during transfection (RET)-rearrangement positive non-small cell lung cancer, Langerhans Cell Histiocytosis with a RET gene fusion, symptomatic or relapsed/refractory Erdheim-Chester Disease with a RET gene fusion, symptomatic or relapsed/refractory Rosai-Dorfman Disease with a RET gene fusion.
Exclusion Criteria	N/A
Required Medical Information	For non-small cell lung cancer, patient must meet all of the following: 1) The disease is recurrent, advanced or metastatic, and 2) Tumor is RET fusion-positive or RET rearrangement-positive.
Age Restrictions	Non-small cell lung cancer: 18 years of age or older. Medullary thyroid cancer and thyroid cancer: 12 years of age or older.
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

REVLIMID

Products Affected

- Lenalidomide

- Revlimid

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Systemic light chain amyloidosis, classical Hodgkin lymphoma, myelodysplastic syndrome without the 5q deletion cytogenetic abnormality, myelofibrosis-associated anemia, POEMS syndrome, myeloproliferative neoplasms, Kaposi Sarcoma, Langerhans cell histiocytosis, peripheral T-Cell lymphomas not otherwise specified, angioimmunoblastic T-cell lymphoma (AITL), enteropathy-associated T-cell lymphoma, monomorphic epitheliotropic intestinal T-cell lymphoma, nodal peripheral T-cell lymphoma, adult T-cell leukemia/lymphoma, hepatosplenic T-cell lymphoma, primary central nervous system (CNS) lymphoma, chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL), acquired immunodeficiency syndrome (AIDS)-related non-germinal center diffuse large B-cell lymphoma, monomorphic post-transplant lymphoproliferative disorder, diffuse large B-cell lymphoma, multicentric Castlemans disease, high-grade B-cell lymphomas, histologic transformation of nodal marginal zone lymphoma to diffuse large B-cell lymphoma, histologic transformation of follicular lymphoma to diffuse large B-cell lymphoma.
Exclusion Criteria	N/A
Required Medical Information	For myelodysplastic syndrome (MDS): Lower risk MDS with symptomatic anemia per the Revised International Prognostic Scoring System (IPSS-R), International Prognostic Scoring System (IPSS), or World Health organization (WHO) classification-based Prognostic Scoring System (WPSS).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

REZLIDHIA

Products Affected

- Rezlidhia

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

REZUROCK

Products Affected

- Rezurock

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	12 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

RINVOQ

Products Affected

- Rinvoq

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderately to severely active rheumatoid arthritis (new starts only): patient has experienced an inadequate treatment response, intolerance or has a contraindication to methotrexate (MTX) AND at least one tumor necrosis factor (TNF) inhibitor (e.g., Enbrel [etanercept], Humira [adalimumab]). For active psoriatic arthritis (new starts only): patient has experienced an inadequate treatment response, intolerance or has a contraindication to at least one TNF inhibitor (e.g., Enbrel [etanercept], Humira [adalimumab]). For moderately to severely active ulcerative colitis (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one TNF inhibitor (e.g., Humira [adalimumab]). For moderately to severely active Crohn's disease (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one TNF inhibitor (e.g., Humira [adalimumab]). For atopic dermatitis, continuation of therapy: the patient achieved or maintained positive clinical response. For active ankylosing spondylitis (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one TNF inhibitor (e.g., Enbrel [etanercept], Humira [adalimumab]). For non-radiographic axial spondyloarthritis (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one TNF inhibitor.
Age Restrictions	Atopic dermatitis: 12 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Atopic dermatitis (initial): 4 months, All others: Plan Year
Other Criteria	N/A

ROZLYTREK

Products Affected

- Rozlytrek

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent ROS1-positive non-small cell lung cancer (NSCLC), Non-metastatic neurotrophic tyrosine receptor kinase (NTRK) gene fusion-positive solid tumors, first-line treatment of NTRK gene fusion-positive solid tumors.
Exclusion Criteria	N/A
Required Medical Information	For all neurotrophic tyrosine receptor kinase (NTRK) gene fusion-positive solid tumors, the disease is without a known acquired resistance mutation. For ROS1-positive non-small cell lung cancer, the patient has recurrent, advanced, or metastatic disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

RUBRACA

Products Affected

- Rubraca

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Uterine leiomyosarcoma, advanced (stage II-IV) epithelial ovarian, fallopian tube, or primary peritoneal cancer
Exclusion Criteria	N/A
Required Medical Information	For metastatic castration-resistant prostate cancer with a deleterious breast cancer susceptibility gene (BRCA) mutation (germline and/or somatic): 1) patient has been treated with androgen receptor-directed therapy, 2) patient has been treated with a taxane-based chemotherapy or the patient is not fit for chemotherapy, 3) the requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy. For maintenance treatment of BRCA mutated epithelial ovarian, fallopian tube, primary peritoneal cancer: 1) the patient has advanced (stage II-IV) disease and is in complete or partial response to primary therapy or 2) the patient has recurrent disease and is in complete or partial response to platinum-based chemotherapy. For uterine leiomyosarcoma: 1) the requested drug is used as second-line therapy AND 2) the patient has BRCA-altered disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

RYBELSUS AET

Products Affected

- Rybelsus

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	The Prior Authorization only applies to patients whose claim is not submitted with an ICD-10 code indicating a diagnosis of type 2 diabetes mellitus OR to patients who do not have a history of an antidiabetic drug (EXCLUDING glucagon-like peptide receptor agonists [GLP-1 RAs] and combination glucose-dependent insulinotropic polypeptide [GIP] and GLP-1 RAs).

RYDAPT

Products Affected

- Rydapt

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Relapsed or refractory acute myeloid leukemia (AML), myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and FGFR1 or FLT3 rearrangements, post-induction therapy for AML, re-induction in residual disease for AML
Exclusion Criteria	N/A
Required Medical Information	For acute myeloid leukemia (AML): AML is FLT3 mutation-positive. For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and FGFR1 or FLT3 rearrangements: the disease is in chronic or blast phase.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SANDOSTATIN LAR

Products Affected

- Sandostatin Lar Depot

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Tumor control of thymomas and thymic carcinomas, tumor control of neuroendocrine tumors (NETs) of the pancreas, gastrointestinal tract, lung, thymus, unresected primary gastrinoma, well-differentiated grade 3 NETs, pheochromocytoma/paraganglioma.
Exclusion Criteria	N/A
Required Medical Information	For acromegaly (initial): 1) patient has a high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range, and 2) patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For acromegaly (continuation of therapy): patient's IGF-1 level has decreased or normalized since initiation of therapy. The requested drug will be used for tumor control for any of the following: 1) neuroendocrine tumor (NET) of the gastrointestinal tract or pancreas in patients with locoregional advanced disease and/or distant metastatic disease, OR 2) NET of the thymus or lung in patients with locoregional unresectable disease and/or distant metastatic disease, OR 3) unresected primary gastrinoma, OR 4) well-differentiated grade 3 unresectable locally advanced or metastatic NET with favorable biology (e.g., relatively low Ki-67 [less than 55%] and somatostatin receptor [SSR] positive imaging), OR 5) thymomas or thymic carcinomas when the following criteria are met: a) locally advanced or metastatic disease or b) postoperatively following tumor resection, OR 6) pheochromocytomas or paragangliomas when the following criteria are met: a) symptomatic locally unresectable disease with SSR positive imaging or b) secreting tumor in distant metastatic disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	N/A
-----------------------	-----

SCSEMBLIX

Products Affected

- Scemblix

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For chronic myeloid leukemia (CML) in the chronic phase: 1) the diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene AND the patient meets either of the following: A) the patient has previously been treated with 2 or more tyrosine kinase inhibitors (TKIs) AND at least one of those was imatinib or dasatinib, OR B) the patient is positive for the T315I mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SIGNIFOR

Products Affected

- Signifor

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	Plan Year
Other Criteria	N/A

SILDENAFIL

Products Affected

- Sildenafil INJ

- Sildenafil Citrate TABS 20MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): Diagnosis was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) If the request is for an adult, pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SIRTURO

Products Affected

- Sirturo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist.
Coverage Duration	Plan Year
Other Criteria	N/A

SKYRIZI

Products Affected

- Skyrizi INJ 150MG/ML, 75MG/0.83ML

- Skyrizi Pen

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) patient meets any of the following: a) patient has experienced an inadequate treatment response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, or b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, or c) patient has severe psoriasis that warrants a biologic as first-line therapy (i.e., at least 10% of the body surface area or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected). For moderately to severely active Crohn's disease (new starts only): 1) Inadequate response to at least one conventional therapy (e.g., corticosteroids), OR 2) Intolerance or contraindication to conventional therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SKYRIZI-CD

Products Affected

- Skyrizi INJ 180MG/1.2ML, 360MG/2.4ML, 600MG/10ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) patient meets any of the following: a) patient has experienced an inadequate treatment response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, or b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, or c) patient has severe psoriasis that warrants a biologic as first-line therapy (i.e., at least 10% of the body surface area or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected). For moderately to severely active Crohn's disease (new starts only): 1) Inadequate response to at least one conventional therapy (e.g., corticosteroids), OR 2) Intolerance or contraindication to conventional therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SOMATULINE DEPOT

Products Affected

- Somatuline Depot

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Tumor control of neuroendocrine tumors (NETs) of the lung, thymus or unresected primary gastrinoma, well-differentiated grade 3 neuroendocrine tumors not of gastroenteropancreatic origin, pheochromocytoma/paraganglioma.
Exclusion Criteria	N/A
Required Medical Information	For acromegaly (initial): 1) patient has a high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range, and 2) patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For acromegaly continuation of therapy: patient's IGF-1 level has decreased or normalized since initiation of therapy. For tumor control of neuroendocrine tumors (NETs) of the thymus or lung: patient has locoregional unresectable disease and/or distant metastatic disease. For tumor control of well-differentiated grade 3 unresectable locally advanced or metastatic NETs (not of gastroenteropancreatic origin): patient has favorable biology (e.g., relatively low Ki-67 [less than 55%] and somatostatin receptor [SSR] positive imaging). For tumor control of pheochromocytomas or paragangliomas: 1) patient has symptomatic locally unresectable disease with SSR positive imaging or 2) patient has distant metastases that are secreting tumors.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SOMAVERT

Products Affected

- Somavert

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For acromegaly (initial): 1) patient has a high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range, and 2) patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For acromegaly (continuation of therapy): patient's IGF-1 level has decreased or normalized since initiation of therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SPRITAM

Products Affected

- Spritam

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Partial-Onset Seizures: 4 years of age or older. Myoclonic Seizures: 12 years of age or older. Primary Generalized Tonic-Clonic Seizures: 6 years of age or older.
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SPRYCEL

Products Affected

- Sprycel

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Gastrointestinal stromal tumor (GIST), metastatic chondrosarcoma, recurrent chordoma, T-cell acute lymphoblastic leukemia (ALL), and Philadelphia (Ph)-like B-ALL, myeloid and/or lymphoid neoplasms with eosinophilia and ABL1 rearrangement in the chronic phase or blast phase
Exclusion Criteria	N/A
Required Medical Information	For chronic myeloid leukemia (CML), including patients who have received a hematopoietic stem cell transplant: 1) Diagnosis was confirmed by detection of the Philadelphia (Ph) chromosome or BCR-ABL gene, and 2) If patient experienced resistance to an alternative tyrosine kinase inhibitor, patient is negative for all of the following mutations: T315I/A, F317L/V/I/C, and V299L. For acute lymphoblastic leukemia (ALL), the patient has a diagnosis of one of the following: 1) Philadelphia chromosome positive ALL, including patients who have received a hematopoietic stem cell transplant: diagnosis that has been confirmed by detection of the Ph chromosome or BCR-ABL gene, and if patient experienced resistance to an alternative tyrosine kinase inhibitor, patient is negative for all of the following mutations: T315I/A, F317L/V/I/C, and V299L, OR 2) Ph-like B-ALL with ABL-class kinase fusion, OR 3) relapsed or refractory T-cell ALL with ABL-class translocation. For GIST, 1) the disease has progressed on imatinib in patients with PDGFRA D842V mutation, OR 2) the patient has failed at least 2 FDA-approved therapies (e.g., imatinib, sunitinib, regorafenib, ripretinib)
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

STELARA

Products Affected

- Stelara

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderate to severe plaque psoriasis (new starts): At least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

STIVARGA

Products Affected

- Stivarga

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Progressive gastrointestinal stromal tumors (GIST), osteosarcoma, glioblastoma, angiosarcoma, retroperitoneal/intra-abdominal soft tissue sarcoma, rhabdomyosarcoma, and soft tissue sarcomas of the extremities, body wall, head and neck, advanced colorectal cancer.
Exclusion Criteria	N/A
Required Medical Information	For gastrointestinal stromal tumors: The disease is progressive, locally advanced, unresectable, or metastatic.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SUTENT

Products Affected

- Sunitinib Malate

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Thyroid carcinoma (follicular, medullary, papillary, and Hurthle cell), soft tissue sarcoma (angiosarcoma, solitary fibrous tumor, and alveolar soft part sarcoma subtypes), recurrent chordoma, thymic carcinoma, lymphoid, myeloid, or mixed lineage neoplasms with eosinophilia
Exclusion Criteria	N/A
Required Medical Information	For renal cell carcinoma (RCC): 1) The disease is relapsed, advanced, or stage IV OR 2) the requested drug is being used as adjuvant treatment for patients that are at high risk of recurrent RCC following nephrectomy. For gastrointestinal stromal tumor (GIST): the disease is unresectable, recurrent, or metastatic AND the patient has failed on an FDA-approved therapy (e.g., imatinib, sunitinib, regorafenib, ripretinib). For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia: 1) The disease has a FLT3 rearrangement AND 2) The disease is in chronic or blast phase.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SYMLIN

Products Affected

- Symlinpen 120
- Symlinpen 60

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SYMPAZAN

Products Affected

- Sympazan

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SYNRIBO

Products Affected

- Synribo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TABRECTA

Products Affected

- Tabrecta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent non-small cell lung cancer (NSCLC).
Exclusion Criteria	N/A
Required Medical Information	For recurrent, advanced, or metastatic NSCLC: Tumor is positive for mesenchymal-epithelial transition (MET) exon 14 skipping mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TADALAFIL (PAH)

Products Affected

- Alyq
- Tadalafil TABS 20MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): Diagnosis was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, and 3) Pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TAFINLAR

Products Affected

- Tafinlar

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Brain metastases from melanoma, thyroid carcinoma (papillary carcinoma, follicular carcinoma, and Hurthle cell carcinoma), central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma), gallbladder cancer, extrahepatic cholangiocarcinoma, intrahepatic cholangiocarcinoma, Langerhans cell histiocytosis, Erdheim-Chester disease
Exclusion Criteria	N/A
Required Medical Information	For brain metastases from melanoma, adjuvant treatment of melanoma, and central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma): 1) The tumor is positive for a BRAF V600 activating mutation (e.g., V600E or V600K), and 2) The requested drug will be used in combination with trametinib. For unresectable or metastatic melanoma: 1) The tumor is positive for a BRAF V600 activating mutation (e.g., V600E or V600K), and 2) The requested drug will be used as a single agent or in combination with trametinib. For non-small cell lung cancer: 1) The tumor is positive for a BRAF V600E mutation, and 2) The requested drug will be used as a single agent or in combination with trametinib. For thyroid carcinoma with papillary, follicular, or Hurthle histology: The tumor is positive for BRAF activating mutation (e.g., V600E or V600K). For Langerhans Cell Histiocytosis and Erdheim-Chester Disease: The disease is positive for a BRAF V600E mutation. For gallbladder cancer, extrahepatic cholangiocarcinoma, and intrahepatic cholangiocarcinoma: 1) The disease is positive for a BRAF V600E mutation and 2) The disease is unresectable or metastatic and 3) The requested drug will be used in combination with trametinib. For solid tumors:) The tumor is positive for a BRAF V600E mutation, and 2) The requested drug will be used in combination with trametinib.
Age Restrictions	N/A
Prescriber Restrictions	N/A

Coverage Duration	Plan Year
Other Criteria	N/A

TAGRISSEO

Products Affected

- Tagrisso

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Sensitizing epidermal growth factor receptor (EGFR) mutation-positive recurrent non-small cell lung cancer (NSCLC), brain metastases from sensitizing EGFR mutation-positive NSCLC, leptomeningeal metastases from EGFR mutation-positive NSCLC.
Exclusion Criteria	N/A
Required Medical Information	For NSCLC, the requested drug is used in any of the following settings: 1) The patient meets both of the following: a) patient has metastatic, advanced, or recurrent NSCLC (including brain and/or leptomeningeal metastases from NSCLC) and b) patient has a sensitizing EGFR mutation OR 2) The patient meets both of the following: a) request is for adjuvant treatment of NSCLC following tumor resection and b) patient has EGFR mutation-positive disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TALTZ

Products Affected

- Taltz

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis AND 2) the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to one of the following products: Enbrel (etanercept), Humira (adalimumab), Otezla (apremilast), Skyrizi (risankizumab-rzaa), Stelara (ustekinumab). For active ankylosing spondylitis (new starts only): the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to one of the following products: Enbrel (etanercept), Humira (adalimumab), Rinvoq (upadacitinib), Xeljanz (tofacitinib)/Xeljanz XR (tofacitinib extended-release). For active psoriatic arthritis (PsA) (new starts only): the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to one of the following products: Enbrel (etanercept), Humira (adalimumab), Otezla (apremilast), Rinvoq (upadacitinib), Skyrizi (risankizumab-rzaa), Stelara (ustekinumab), Xeljanz (tofacitinib)/Xeljanz XR (tofacitinib extended-release). For active non-radiographic axial spondyloarthritis (new starts only): Patient meets any of the following: 1) has had an inadequate response to a non-steroidal anti-inflammatory drug (NSAID) trial or 2) has an intolerance or contraindication to NSAIDs.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TALZENNA

Products Affected

- Talzenna

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent germline breast cancer susceptibility gene (BRCA)-mutated breast cancer
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TARGRETIN TOPICAL

Products Affected

- Bexarotene GEL

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Stage 2 or higher mycosis fungoides/Sezary syndrome, chronic or smoldering adult T-cell leukemia/lymphoma, primary cutaneous marginal zone lymphoma, primary cutaneous follicle center lymphoma.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TASIGNA

Products Affected

- Tasigna

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL), gastrointestinal stromal tumor (GIST), myeloid and/or lymphoid neoplasms with eosinophilia and ABL1 rearrangement in the chronic phase or blast phase.
Exclusion Criteria	N/A
Required Medical Information	For chronic myeloid leukemia (CML) or acute lymphoblastic leukemia (ALL), diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene. For CML, including patients newly diagnosed with CML and patients who have received a hematopoietic stem cell transplant: patient has experienced resistance or intolerance to imatinib or dasatinib. If patient experienced resistance to an alternative tyrosine kinase inhibitor for CML, patient is negative for T315I, Y253H, E255K/V, and F359V/C/I mutations. For GIST, patient must have progressed on imatinib, sunitinib, and regorafenib.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TAZAROTENE

Products Affected

- Tazarotene CREA
- Tazarotene GEL
- Tazorac CREA 0.05%

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For plaque psoriasis: 1)The requested drug is being prescribed to treat less than or equal to 20 percent of the patient's body surface area (BSA) AND 2) the patient experienced an inadequate treatment response or intolerance to at least one topical corticosteroid OR has a contraindication that would prohibit a trial of topical corticosteroids.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TAZVERIK

Products Affected

- Tazverik

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Epithelioid sarcoma: 16 years of age or older, Follicular lymphoma: 18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TECFIDERA

Products Affected

- Tecfidera
- Tecfidera Starter Pack

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TECVAYLI

Products Affected

- Tecvayli

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TEMAZEPAM 30MG - 65

Products Affected

- Temazepam CAPS 30MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with the requested drug outweighs the potential risks to the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to doxepin (3 mg or 6 mg).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization requirement applies only to patients 70 years of age or older.

TEPMETKO

Products Affected

- Tepmetko

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent non-small cell lung cancer (NSCLC).
Exclusion Criteria	N/A
Required Medical Information	For recurrent, advanced, or metastatic NSCLC: Tumor is positive for mesenchymal-epithelial transition (MET) exon 14 skipping mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TESTOSTERONE ENANTHATE INJ

Products Affected

- Testosterone Enanthate INJ

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Gender Dysphoria
Exclusion Criteria	N/A
Required Medical Information	Primary or hypogonadotropic hypogonadism: 1) Request is for continuation of testosterone therapy and the patient had a confirmed low morning testosterone level according to current practice guidelines or your standard lab reference values before starting testosterone therapy [Note: Safety and efficacy of testosterone products in patients with "age-related hypogonadism" (also referred to as "late-onset hypogonadism") have not been established.] OR 2) Request is not for continuation of testosterone therapy and the patient has at least two confirmed low morning testosterone levels according to current practice guidelines or your standard lab reference values [Note: Safety and efficacy of testosterone products in patients with "age-related hypogonadism" (also referred to as "late-onset hypogonadism") have not been established.]. Gender dysphoria: The patient is able to make an informed decision to engage in hormone therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TETRABENAZINE

Products Affected

- Tetrabenazine

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Tic disorders, tardive dyskinesia, hemiballismus, chorea not associated with Huntington's disease.
Exclusion Criteria	N/A
Required Medical Information	For treatment of chorea associated with Huntington's disease: The patient must have a prior inadequate response or intolerable adverse event with deutetrabenazine therapy. For treatment of tardive dyskinesia: The patient must have a prior inadequate response or intolerable adverse event with deutetrabenazine or valbenazine therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

THALOMID

Products Affected

- Thalomid

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Myelofibrosis-related anemia, AIDS-related aphthous stomatitis, Kaposi sarcoma, chronic graft-versus-host disease, Crohn's disease, multicentric Castleman's disease., Rosai-Dorfman disease, Langerhans cell histiocytosis
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TIBSOVO

Products Affected

- Tibsovo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Conventional (grades 1-3) or dedifferentiated chondrosarcoma. Newly-diagnosed acute myeloid leukemia (AML) if 60-74 years of age and without comorbidities.
Exclusion Criteria	N/A
Required Medical Information	Patient has disease with a susceptible isocitrate dehydrogenase-1 (IDH1) mutation. For acute myeloid leukemia (AML): 1) patient has newly-diagnosed AML and meets one of the following: a) 75 years of age or older, b) patient has comorbidities that preclude use of intensive induction chemotherapy, or c) patient is 60 years of age or older and declines intensive induction chemotherapy, OR 2) patient is 60 years of age or older and the requested drug will be used as post-induction therapy following response to induction therapy with the requested drug, OR 3) patient has relapsed or refractory AML. For locally advanced, unresectable, or metastatic cholangiocarcinoma: the requested drug will be used as subsequent treatment for progression on or after systemic treatment.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TOBRAMYCIN

Products Affected

- Tobramycin NEBU 300MG/5ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Non-cystic fibrosis bronchiectasis
Exclusion Criteria	N/A
Required Medical Information	For cystic fibrosis and non-cystic fibrosis bronchiectasis, the patient must meet one of the following: 1) Pseudomonas aeruginosa is present in the patient's airway cultures, OR 2) the patient has a history of Pseudomonas aeruginosa infection or colonization in the airways.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

TOPICAL DOXEPIN

Products Affected

- Doxepin Hydrochloride CREA

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient has experienced an inadequate response to a topical corticosteroid or a topical calcineurin inhibitor.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 month
Other Criteria	N/A

TOPICAL LIDOCAINE

Products Affected

- Lidocaine OINT 5%

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1) The requested drug is being used for topical anesthesia, AND 2) If the requested drug will be used as part of a compounded product, then all the active ingredients in the compounded product are Food and Drug Administration (FDA) approved for topical use.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

TOPICAL TRETINOIN

Products Affected

- Tretinoin CREA
- Tretinoin GEL
- Tretinoin Microsphere GEL 0.04%, 0.1%
- Tretinoin Microsphere Pump

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TOREMIFENE

Products Affected

- Toremifene Citrate

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Congenital/acquired QT prolongation (long QT syndrome), uncorrected hypokalemia, or uncorrected hypomagnesemia.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TRIENTINE

Products Affected

- Trientine Hydrochloride

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TRIKAFTA

Products Affected

- Trikafta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For cystic fibrosis: The requested medication will not be used in combination with other medications containing ivacaftor.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TRULICITY AET

Products Affected

- Trulicity

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	10 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	The Prior Authorization only applies to patients whose claim is not submitted with an ICD-10 code indicating a diagnosis of type 2 diabetes mellitus OR to patients who do not have a history of an antidiabetic drug (EXCLUDING glucagon-like peptide receptor agonists [GLP-1 RAs] and combination glucose-dependent insulinotropic polypeptide [GIP] and GLP-1 RAs).

TRUSELTIQ

Products Affected

- Truseltiq

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TRUXIMA

Products Affected

- Truxima

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	<p>Non-Hodgkin's lymphoma subtypes [small lymphocytic lymphoma (SLL), mantle cell lymphoma, marginal zone lymphomas (nodal, splenic, gastric mucosa-associated lymphoid tissue [MALT], nongastric MALT), Burkitt lymphoma, primary cutaneous B-cell lymphoma, high-grade B-cell lymphoma with translocations of MYC and BCL2 and/or BCL6 (double/triple hit lymphoma), high-grade B-cell lymphoma not otherwise specified, histological transformation from follicular lymphoma to diffuse large B-cell lymphoma, histological transformation from nodal marginal zone lymphoma to diffuse large B-cell lymphoma, Castleman's disease, acquired immunodeficiency syndrome (AIDS)-related B-cell lymphoma, hairy cell leukemia, post-transplant lymphoproliferative disorder (PTLD), B-cell lymphoblastic lymphoma], refractory immune or idiopathic thrombocytopenic purpura (ITP), autoimmune hemolytic anemia, Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma, chronic graft-versus-host disease (GVHD), Sjogren syndrome, thrombotic thrombocytopenic purpura, refractory myasthenia gravis, Hodgkin's lymphoma (nodular lymphocyte-predominant), primary central nervous system (CNS) lymphoma, leptomeningeal metastases from lymphomas, acute lymphoblastic leukemia, prevention of Epstein-Barr virus (EBV)-related PTLD, multiple sclerosis, immune checkpoint inhibitor-related toxicities, pemphigus vulgaris, pediatric aggressive mature B-cell lymphomas, and Rosai-Dorfman disease, and pediatric mature B-cell acute leukemia.</p>
Exclusion Criteria	N/A

Required Medical Information	For moderately to severely active rheumatoid arthritis (new starts only): 1) patient meets ANY of the following: a) requested drug will be used in combination with methotrexate (MTX) OR b) patient has intolerance or contraindication to MTX, AND 2) patient meets ANY of the following: a) inadequate response, intolerance, or contraindication to MTX OR b) inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. Hematologic malignancies must be CD20-positive. For multiple sclerosis: 1) patient has a diagnosis of relapsing remitting multiple sclerosis and 2) patient has had an inadequate response to two or more disease-modifying drugs indicated for multiple sclerosis despite adequate duration of treatment.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Immune checkpoint inhibitor-related toxicities: 3 months, All other: Plan Year
Other Criteria	N/A

TUKYSA

Products Affected

- Tukysa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent human epidermal growth factor receptor 2 (HER2)-positive breast cancer
Exclusion Criteria	N/A
Required Medical Information	For colorectal cancer (including appendiceal adenocarcinoma): 1) the patient has advanced, unresectable, or metastatic disease AND 2) the patient has human epidermal growth factor receptor 2 (HER2)-positive disease AND 3) the patient has RAS wild-type disease AND 4) the requested drug will be used in combination with trastuzumab and 5) the patient has not previously been treated with a HER2 inhibitor.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TURALIO

Products Affected

- Turalio

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Langerhans Cell Histiocytosis, Erdheim-Chester Disease, Rosai-Dorfman Disease
Exclusion Criteria	N/A
Required Medical Information	For Langerhans Cell Histiocytosis: 1) disease has colony stimulating factor 1 receptor (CSF1R) mutation. For Erdheim-Chester Disease and Rosai-Dorfman Disease: 1) disease has CSF1R mutation AND patient has any of the following: a) symptomatic disease OR b) relapsed/refractory disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

UKONIQ

Products Affected

- Ukoniq

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VALCHLOR

Products Affected

- Valchlor

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Chronic or smoldering adult T-cell leukemia/lymphoma, Stage 2 or higher mycosis fungoides/Sezary syndrome, primary cutaneous marginal zone lymphoma, primary cutaneous follicle center lymphoma, lymphomatoid papulosis, unifocal Langerhans cell histiocytosis (LCH) with isolated skin disease.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VALTOCO

Products Affected

- Valtoco 10 Mg Dose
- Valtoco 15 Mg Dose
- Valtoco 20 Mg Dose
- Valtoco 5 Mg Dose

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	6 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VANFLYTA

Products Affected

- Vanflyta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VENCLEXTA

Products Affected

- Venclexta

- Venclexta Starting Pack

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Mantle cell lymphoma, blastic plasmacytoid dendritic cell neoplasm (BPDCN), multiple myeloma, relapsed or refractory acute myeloid leukemia (AML), Waldenstrom macroglobulinemia/lymphoplasmacytic lymphoma, relapsed or refractory systemic light chain amyloidosis with translocation t(11:14)
Exclusion Criteria	N/A
Required Medical Information	For acute myeloid leukemia (AML): 1) patient is 60 years of age or older OR 2) patient is less than 60 years of age with unfavorable risk genetics and TP53-mutation OR 3) patient has comorbidities that preclude use of intensive induction chemotherapy OR 4) patient has relapsed or refractory disease. For blastic plasmacytoid dendritic cell neoplasm (BPDCN): 1) patient has systemic disease being treated with palliative intent OR 2) patient has relapsed or refractory disease. For multiple myeloma: 1) the disease is relapsed or progressive AND 2) the requested drug will be used in combination with dexamethasone AND 3) patient has t(11:14) translocation. For Waldenstrom macroglobulinemia/lymphoplasmacytic lymphoma: 1) patient has previously treated disease that did not respond to primary therapy OR 2) patient has progressive or relapsed disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VENTAVIS

Products Affected

- Ventavis

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (World Health Organization [WHO] Group 1), the diagnosis was confirmed by right heart catheterization. For new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

VERSACLOZ

Products Affected

- Versacloz

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the treatment of a severely ill patient with schizophrenia who failed to respond adequately to standard antipsychotic treatment (i.e., treatment-resistant schizophrenia): 1) the patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following generic products: A) aripiprazole, B) asenapine, C) olanzapine, D) quetiapine, E) risperidone, F) ziprasidone AND 2) The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following brand products: A) Latuda, B) Rexulti, C) Secuado, D) Vraylar.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VERZENIO

Products Affected

- Verzenio

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer in combination with fulvestrant or an aromatase inhibitor, or as a single agent if progression on prior endocrine therapy and prior chemotherapy in the metastatic setting.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VICTOZA AET

Products Affected

- Victoza

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	The Prior Authorization only applies to patients whose claim is not submitted with an ICD-10 code indicating a diagnosis of type 2 diabetes mellitus OR to patients who do not have a history of an antidiabetic drug (EXCLUDING glucagon-like peptide receptor agonists [GLP-1 RAs] and combination glucose-dependent insulinotropic polypeptide [GIP] and GLP-1 RAs).

VIGABATRIN

Products Affected

- Vigabatrin
- Vigadrone

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For complex partial seizures (CPS): patient has experienced an inadequate treatment response to at least two antiepileptic drugs for CPS.
Age Restrictions	Infantile Spasms: 1 month to 2 years of age. CPS: 2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VITRAKVI

Products Affected

- Vitrakvi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Non-metastatic neurotrophic tyrosine receptor kinase (NTRK) gene fusion-positive solid tumors, first-line treatment of NTRK gene fusion-positive solid tumors.
Exclusion Criteria	N/A
Required Medical Information	For all neurotrophic tyrosine receptor kinase (NTRK) gene fusion-positive solid tumors, the disease is without a known acquired resistance mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VIZIMPRO

Products Affected

- Vizimpro

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent non-small cell lung cancer (NSCLC).
Exclusion Criteria	N/A
Required Medical Information	For non-small cell lung cancer (NSCLC): 1) the disease is recurrent, advanced or metastatic, and 2) the patient has sensitizing EGFR mutation-positive disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VONJO

Products Affected

- Vonjo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VORICONAZOLE

Products Affected

- Voriconazole INJ
- Voriconazole SUSR

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient will use the requested drug orally or intravenously.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

VOSEVI

Products Affected

- Vosevi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Decompensated cirrhosis/moderate or severe hepatic impairment (Child Turcotte Pugh class B or C)
Required Medical Information	For hepatitis C: Infection confirmed by presence of HCV RNA in the serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of HIV coinfection, presence or absence of resistance-associated substitutions where applicable, transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current American Association for the Study of Liver Diseases (AASLD) treatment guidelines.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Criteria will be applied consistent with current AASLD-IDSA guidance.
Other Criteria	N/A

VOTRIENT

Products Affected

- Votrient

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Thyroid carcinoma (follicular, papillary, Hurthle cell, or medullary), uterine sarcoma, chondrosarcoma, gastrointestinal stromal tumor.
Exclusion Criteria	N/A
Required Medical Information	For renal cell carcinoma: The disease is advanced, relapsed, or stage IV. For gastrointestinal stromal tumor (GIST): the disease is unresectable, recurrent, or metastatic AND the patient has failed on an FDA-approved therapy (e.g., imatinib, sunitinib, regorafenib, ripretinib). For soft tissue sarcoma (STS): The patient does not have an adipocytic soft tissue sarcoma. For uterine sarcoma: The disease is recurrent or metastatic.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VUMERITY

Products Affected

- Vumerity

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

WELIREG

Products Affected

- Welireg

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XALKORI

Products Affected

- Xalkori

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent non-small cell lung cancer (NSCLC), NSCLC with high-level MET amplification or MET exon 14 skipping mutation, symptomatic or relapsed/refractory anaplastic lymphoma kinase (ALK)-fusion positive Erdheim-Chester Disease, symptomatic or relapsed/refractory (ALK)-fusion positive Rosai-Dorfman Disease, (ALK)-fusion positive Langerhans Cell Histiocytosis.
Exclusion Criteria	N/A
Required Medical Information	For NSCLC, the requested drug is used in any of the following settings: 1) the patient has recurrent, advanced or metastatic ALK-positive NSCLC, 2) the patient has recurrent, advanced or metastatic ROS-1 positive NSCLC, or 3) the patient has NSCLC with high-level MET amplification or MET exon 14 skipping mutation. For IMT, the disease is ALK-positive. For ALCL, the disease is relapsed or refractory and ALK-positive.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XELJANZ

Products Affected

- Xeljanz

- Xeljanz Xr

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderately to severely active rheumatoid arthritis (new starts only): 1) patient has experienced an inadequate treatment response, intolerance or has a contraindication to methotrexate (MTX) AND at least one tumor necrosis factor (TNF) inhibitor (e.g., Enbrel [etanercept], Humira [adalimumab]). For active psoriatic arthritis (new starts only): 1) patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one TNF inhibitor (e.g., Enbrel [etanercept], Humira [adalimumab]) AND 2) the requested drug is used in combination with a nonbiologic DMARD. For active ankylosing spondylitis (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one tumor necrosis factor (TNF) inhibitor (e.g., Enbrel [etanercept], Humira [adalimumab]). For moderately to severely active ulcerative colitis (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one tumor necrosis factor (TNF) inhibitor (e.g., Humira [adalimumab]). For active polyarticular course juvenile idiopathic arthritis (pcJIA) (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one tumor necrosis factor (TNF) inhibitor (e.g., Enbrel [etanercept], Humira [adalimumab]).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XERMELO

Products Affected

- Xermelo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XGEVA

Products Affected

- Xgeva

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hypercalcemia of malignancy: condition is refractory to intravenous (IV) bisphosphonate therapy or there is a clinical reason to avoid IV bisphosphonate therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

XHANCE MAPD

Products Affected

- Xhance

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Patient has experienced an inadequate treatment response to generic fluticasone nasal spray.
Age Restrictions	18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XIFAXAN

Products Affected

- Xifaxan TABS 550MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For irritable bowel syndrome with diarrhea (IBS-D): 1) The patient has not previously received treatment with the requested drug OR 2) The patient has previously received treatment with the requested drug AND a) the patient is experiencing a recurrence of symptoms AND b) the patient has not already received an initial 14-day course of treatment and two additional 14-day courses of treatment with the requested drug.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Reduction in risk of overt HE recurrence: 6 Months, IBS-D: 14 Days
Other Criteria	N/A

XOLAIR

Products Affected

- Xolair

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For allergic asthma initial therapy: 1) Patient has positive skin test (or blood test) to at least 1 perennial aeroallergen, 2) Patient has baseline IgE level greater than or equal to 30 IU/mL, and 3) Patient has inadequate asthma control despite current treatment with both of the following medications: a) Inhaled corticosteroid, and b) Additional controller (long acting beta2-agonist, long-acting muscarinic antagonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies. For allergic asthma continuation therapy only: Patient's asthma control has improved on treatment with the requested drug since initiation of therapy. For chronic idiopathic urticaria (CIU) initial therapy: 1) Patient has been evaluated for other causes of urticaria, including bradykinin-related angioedema and IL-1-associated urticarial syndromes (auto-inflammatory disorders, urticarial vasculitis), and 2) Patient has experienced a spontaneous onset of wheals, angioedema, or both, for at least 6 weeks. For CIU continuation therapy: Patient has experienced a response (e.g., improved symptoms) since initiation of therapy. For nasal polyps: 1) the requested drug is used as add-on maintenance treatment, AND 2) the patient has experienced an inadequate treatment response to Xhance (fluticasone).
Age Restrictions	For CIU: 12 years of age or older. For allergic asthma: 6 years of age or older. For nasal polyps: 18 years of age or older.
Prescriber Restrictions	N/A
Coverage Duration	CIU initial: 6 months. All others: Plan Year.
Other Criteria	N/A

XOSPATA

Products Affected

- Xospata

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and FLT3 rearrangement
Exclusion Criteria	N/A
Required Medical Information	For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and FLT3 rearrangement: the disease is in chronic or blast phase.
Age Restrictions	18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XPOVIO

Products Affected

- Xpovio
- Xpovio 60 Mg Twice Weekly
- Xpovio 80 Mg Twice Weekly

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XTANDI

Products Affected

- Xtandi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XYREM

Products Affected

- Sodium Oxybate

- Xyrem

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The diagnosis has been confirmed by sleep lab evaluation. EDS: 1)The patient experienced an inadequate treatment response or intolerance to at least one central nervous system (CNS) stimulant drug (e.g., amphetamine, dextroamphetamine, or methylphenidate) OR has a contraindication that would prohibit a trial of central nervous system (CNS) stimulant drugs (e.g., amphetamine, dextroamphetamine, or methylphenidate) [Note: Coverage of amphetamines may require prior authorization.] AND 2) If the patient is 18 years of age or older, the patient experienced an inadequate treatment response or intolerance to at least one central nervous system (CNS) wakefulness promoting drug (e.g., armodafinil) OR has a contraindication that would prohibit a trial of central nervous system (CNS) wakefulness promoting drugs (e.g., armodafinil) [Note: coverage of armodafinil may require prior authorization.].
Age Restrictions	7 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with a sleep disorder specialist or neurologist.
Coverage Duration	Plan Year
Other Criteria	If the request is for a continuation of therapy, then the patient experienced a decrease in daytime sleepiness with narcolepsy or a decrease in cataplexy episodes with narcolepsy.

ZARXIO

Products Affected

- Zarxio

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Neutropenia in myelodysplastic syndromes (MDS), agranulocytosis, neutropenia in aplastic anemia, human immunodeficiency virus (HIV)-related neutropenia, neutropenia related to renal transplant.
Exclusion Criteria	Use of the requested product within 24 hours prior to or following chemotherapy.
Required Medical Information	For prophylaxis or treatment of myelosuppressive chemotherapy-induced febrile neutropenia (FN) patient must meet both of the following: 1) Patient has a solid tumor or non-myeloid cancer, and 2) Patient has received, is currently receiving, or will be receiving treatment with myelosuppressive anti-cancer therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

ZEJULA

Products Affected

- Zejula

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	In combination with bevacizumab for persistent or recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer for platinum-sensitive disease, uterine leiomyosarcoma
Exclusion Criteria	N/A
Required Medical Information	For uterine leiomyosarcoma: 1) the requested drug is used as second-line therapy AND 2) the patient has BRCA-altered disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZELBORAF

Products Affected

- Zelboraf

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Non-small cell lung cancer, hairy cell leukemia, thyroid carcinoma (i.e., papillary carcinoma, follicular carcinoma, and Hurthle cell carcinoma), central nervous system cancer (i.e., glioma, meningioma, astrocytoma), adjuvant systemic therapy for cutaneous melanoma, Langerhans cell histiocytosis.
Exclusion Criteria	N/A
Required Medical Information	For adjuvant treatment of melanoma, and central nervous system (CNS) cancer (i.e., glioma, meningioma, astrocytoma): 1) The tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K) and 2) The requested drug will be used in combination with cobimetinib. For unresectable or metastatic melanoma: 1) The tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K) and 2) the requested drug will be used as a single agent, or in combination with cobimetinib (with or without atezolizumab). For Erdheim-Chester Disease and Langerhans Cell Histiocytosis: Tumor is positive for BRAF V600 mutation. For non-small cell lung cancer: 1) Tumor is positive for the BRAF V600E mutation, and 2) The patient has recurrent, advanced, or metastatic disease. For thyroid carcinoma: 1) Tumor is positive for BRAF mutation, and 2) Patient has radioiodine refractory follicular, Hurthle cell, or papillary thyroid carcinoma. For hairy cell leukemia: The requested drug will be used for subsequent therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZIRABEV

Products Affected

- Zirabev

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Breast cancer, central nervous system (CNS) tumor types: adult low-grade (WHO Grade I or II) glioma, adult intracranial and spinal ependymoma, anaplastic gliomas, adult medulloblastoma, primary central nervous system lymphoma, meningiomas, limited and extensive brain metastases, and metastatic spine tumors, malignant pleural mesothelioma, ovarian cancer/fallopian tube cancer/primary peritoneal cancer types: carcinosarcoma (malignant mixed Mullerian tumors), clear cell carcinoma, mucinous carcinoma, grade 1 endometrioid carcinoma, low-grade serous carcinoma, ovarian borderline epithelial tumors (low malignant potential) with invasive implants, and malignant sex cord-stromal tumors, soft tissue sarcoma types: angiosarcoma and solitary fibrous tumor/hemangiopericytoma, uterine neoplasms, endometrial carcinoma, vulvar squamous cell carcinoma, and ophthalmic-related disorders: diabetic macular edema, neovascular (wet) age-related macular degeneration including polypoidal choroidopathy and retinal angiomatous proliferation subtypes, macular edema following retinal vein occlusion, proliferative diabetic retinopathy, choroidal neovascularization, neovascular glaucoma and retinopathy of prematurity, hepatocellular carcinoma, small bowel adenocarcinoma.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.
-----------------------	--

ZOLINZA

Products Affected

- Zolanza

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Mycosis fungoides, Sezary syndrome.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZONISADE

Products Affected

- Zonisade

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For adjunctive treatment of partial-onset seizures (i.e., focal-onset seizures): 1) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to a generic anticonvulsant AND the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to any of the following: Aptiom, Xcopri, Spritam OR 2) The patient has difficulty swallowing solid oral dosage forms (e.g., tablets, capsules).
Age Restrictions	16 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZTALMY

Products Affected

- Ztalmy

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZYDELIG

Products Affected

- Zydelig

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Refractory chronic lymphocytic leukemia (CLL), relapsed or refractory small lymphocytic lymphoma (SLL).
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZYKADIA

Products Affected

- Zykadia TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent ALK-positive non-small cell lung cancer (NSCLC), recurrent, advanced, or metastatic ROS1-positive NSCLC, inflammatory myofibroblastic tumor (IMT), brain metastases from NSCLC.
Exclusion Criteria	N/A
Required Medical Information	For NSCLC: the patient has recurrent, advanced, or metastatic ALK-positive or ROS1-positive disease. For inflammatory myofibroblastic tumor: the disease is ALK-positive. For brain metastases from NSCLC: the patient has ALK-positive NSCLC.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZYPREXA RELPREVV

Products Affected

- Zyprexa Relprevv

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Tolerability with oral olanzapine has been established.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PART B VERSUS PART D

Products Affected

- Abelcet
- Acetylcysteine INHALATION SOLN
- Acyclovir Sodium INJ 50MG/ML
- Albuterol Sulfate NEBU 0.083%, 0.63MG/3ML, 1.25MG/3ML, 2.5MG/0.5ML
- Amphotericin B INJ
- Amphotericin B Liposome
- Aprepitant CAPS
- Azathioprine INJ
- Azathioprine TABS 50MG
- Budesonide SUSP
- Clinimix 4.25%/dextrose 10%
- Clinimix 4.25%/dextrose 5%
- Clinimix 5%/dextrose 15%
- Clinimix 5%/dextrose 20%
- Clinimix 6/5
- Clinimix 8/10
- Clinimix 8/14
- Clinisol Sf 15%
- Clinolipid
- Cromolyn Sodium NEBU
- Cyclophosphamide CAPS
- Cyclophosphamide TABS
- Cyclosporine CAPS
- Cyclosporine INJ
- Cyclosporine Modified
- Dextrose 50%
- Dextrose 70%
- Emend SUSR
- Engerix-b
- Epoprostenol Sodium
- Everolimus TABS 0.25MG, 0.5MG, 0.75MG, 1MG
- Fiasp Pumpcart
- Freamine III INJ 89MEQ/L; 710MG/100ML; 950MG/100ML; 3MEQ/L; 24MG/100ML; 1400MG/100ML; 280MG/100ML; 690MG/100ML; 910MG/100ML; 730MG/100ML; 530MG/100ML; 560MG/100ML; 10MMOLE/L; 120MG/100ML; 1120MG/100ML; 590MG/100ML; 10MEQ/L; 400MG/100ML; 150MG/100ML; 660MG/100ML
- Gamastan
- Ganciclovir INJ 500MG, 500MG/10ML
- Gengraf CAPS 100MG, 25MG
- Gengraf SOLN
- Granisetron Hydrochloride TABS
- Hepatamine INJ 62MEQ/L; 770MG/100ML; 600MG/100ML; 3MEQ/L; 20MG/100ML; 900MG/100ML; 240MG/100ML; 900MG/100ML; 1100MG/100ML; 610MG/100ML; 100MG/100ML; 100MG/100ML; 115MG/100ML; 800MG/100ML; 500MG/100ML; 450MG/100ML; 66MG/100ML; 840MG/100ML
- Heplisav-b
- Humulin R U-500 (concentrated)
- Hydromorphone Hcl INJ 10MG/ML, 1MG/ML, 4MG/ML
- Hydromorphone Hydrochloride INJ 1MG/ML, 2MG/ML, 4MG/ML, 50MG/5ML
- Imovax Rabies (h.d.c.v.)
- Ipratropium Bromide INHALATION SOLN 0.02%
- Ipratropium Bromide/albuterol Sulfate
- Isolyte-s INJ 27MEQ/L; 98MEQ/L; 23MEQ/L; 3MEQ/L; 5MEQ/L; 140MEQ/L
- Isolyte-s Ph 7.4
- Jynneos
- Levalbuterol NEBU

- Levalbuterol Hcl NEBU
- Levalbuterol Hydrochloride NEBU 0.63MG/3ML
- Methylprednisolone TABS
- Methylprednisolone Acetate INJ 40MG/ML, 80MG/ML
- Methylprednisolone Sodium Succinate INJ 1000MG
- Methylprednisolone Sodiamsuccinate INJ 125MG, 40MG
- Morphine Sulfate INJ 0.5MG/ML, 10MG/ML, 1MG/ML, 2MG/ML, 4MG/ML, 50MG/ML, 5MG/ML, 8MG/ML
- Morphine Sulfate/sodium Chloride INJ 1MG/ML
- Mycophenolate Mofetil CAPS
- Mycophenolate Mofetil INJ
- Mycophenolate Mofetil SUSR
- Mycophenolate Mofetil TABS
- Mycophenolic Acid Dr
- Nulojix
- Nutrilipid
- Ondansetron Hcl SOLN
- Ondansetron Hcl TABS 24MG
- Ondansetron Hydrochloride TABS
- Ondansetron Odt
- Pentamidine Isethionate INHALATION SOLR
- Plenamine INJ 147.4MEQ/L; 2.17GM/100ML; 1.47GM/100ML; 434MG/100ML; 749MG/100ML; 1.04GM/100ML; 894MG/100ML; 749MG/100ML; 1.04GM/100ML; 1.18GM/100ML; 749MG/100ML; 1.04GM/100ML; 894MG/100ML; 592MG/100ML; 749MG/100ML; 250MG/100ML; 39MG/100ML; 960MG/100ML
- Prednisolone SOLN
- Prednisolone Sodium Phosphate ORAL SOLN 10MG/5ML, 15MG/5ML, 20MG/5ML, 25MG/5ML, 5MG/5ML
- Prednisone SOLN
- Prednisone TABS 10MG, 1MG, 2.5MG, 20MG, 50MG, 5MG
- Prednisone IntenSol
- Prehevbrio
- Premasol INJ 52MEQ/L; 1760MG/100ML; 880MG/100ML; 34MEQ/L; 1760MG/100ML; 372MG/100ML; 406MG/100ML; 526MG/100ML; 492MG/100ML; 492MG/100ML; 526MG/100ML; 356MG/100ML; 356MG/100ML; 390MG/100ML; 34MG/100ML; 152MG/100ML
- Procalamine
- Prograf PACK
- Prosol
- Rabavert
- Recombivax Hb
- Sandimmune SOLN
- Sirolimus SOLN
- Sirolimus TABS
- Tacrolimus CAPS
- Tpn Electrolytes INJ 29.5MEQ/20ML; 4.5MEQ/20ML; 35MEQ/20ML; 5MEQ/20ML; 20MEQ/20ML; 35MEQ/20ML
- Travasol INJ 52MEQ/L; 1760MG/100ML; 880MG/100ML; 34MEQ/L; 1760MG/100ML; 372MG/100ML; 406MG/100ML; 526MG/100ML; 492MG/100ML; 492MG/100ML; 526MG/100ML; 356MG/100ML; 500MG/100ML; 356MG/100ML; 390MG/100ML; 34MG/100ML; 152MG/100ML
- Trophamine INJ 0.54GM/100ML; 1.2GM/100ML; 0.32GM/100ML; 0; 0; 0.5GM/100ML; 0.36GM/100ML; 0.48GM/100ML; 0.82GM/100ML; 1.4GM/100ML; 1.2GM/100ML; 0.34GM/100ML; 0.48GM/100ML; 0.68GM/100ML; 0.38GM/100ML; 5MEQ/L; 0.025GM/100ML; 0.42GM/100ML; 0.2GM/100ML; 0.24GM/100ML; 0.78GM/100ML

Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

INDEX

A

Abelcet	322
Abiraterone.....	1
Abiraterone Acetate.....	1
Accutane.....	141
Acetylcysteine	322
Acitretin.....	2
Actimmune.....	3
Acyclovir Sodium	322
Adempas.....	4
Aimovig.....	5
Akeega.....	6
Albuterol Sulfate	322
Alecensa	7
Alosetron	8
Alosetron Hydrochloride.....	8
Alpha 1-proteinase Inhibitor	9
Alunbrig	10
Alyq.....	251
Ambrisentan	11
Amnesteem.....	141
Amphotericin B.....	322
Amphotericin B Liposome	322
Aprepitant.....	322
Arcalyst	12
Armodafinil.....	13
Asparlas.....	14
Atovaquone	175
Aubagio	15
Austedo	16
Austedo Xr	16
Austedo Xr Patient Titration Kit.....	16
Auvelity.....	17
Avonex	18
Avonex Pen	18
Ayvakit.....	19
Azathioprine	322

B

Balversa.....	20
Banzel	21
Benlysta.....	22
Benztropine Mesylate	113
Berinert	23
Besremi	24
Betaseron.....	25
Bexarotene	26, 257
Bivigam.....	144
Bosentan.....	27
Bosulif.....	28
Braftovi	29
Briviact.....	30, 31
Briviact Inj	31
Brukina	32
Budesonide.....	322
Buprenorphine.....	33, 34
Buprenorphine Hcl.....	33
Buprenorphine Patch.....	34
Bydureon Aet.....	35
Bydureon Beise.....	35
Byetta	36
Byetta Aet	36

C

Cabometyx	37
Calcipotriene.....	38
Calcitrene	38
Calcitriol	38
Calquence.....	39
Caprelsa.....	40
Carac	41
Carbaglu.....	42
Carbinoxamine Maleate	120
Carglumic Acid.....	42
Cayston	43

Cerdelga	44
Chantix	45
Chlordiazepoxide - 65	46
Chlordiazepoxide Hcl.....	46
Chlordiazepoxide Hydrochloride	46
Chlorzoxazone.....	123
Claravis.....	141
Clemastine Fumarate.....	121
Clinimix 4.25%/dextrose 10%	322
Clinimix 4.25%/dextrose 5%	322
Clinimix 5%/dextrose 15%	322
Clinimix 5%/dextrose 20%	322
Clinimix 6/5	322
Clinimix 8/10	322
Clinimix 8/14	322
Clinisol Sf 15%	322
Clinolipid.....	322
Clobazam.....	47
Clomipramine.....	48
Clomipramine Hydrochloride	48
Clorazepate.....	49
Clorazepate Dipotassium.....	49
Clozapine Odt.....	50
Colistimethate Sodium	51
Coly-mycin.....	51
Cometriq.....	52
Copaxone.....	102
Copiktra.....	53
Cotellic	54
Cromolyn Sodium	322
Cyclobenzaprine Hydrochloride	123
Cyclophosphamide	322
Cyclosporine	322
Cyclosporine Modified.....	322
Cyproheptadine Hcl.....	114
Cyproheptadine Hydrochloride	114
Cystagon.....	55
Cystaran.....	56

D

Dalfampridine.....	57
Dalfampridine Er.....	57
Daurismo	58
Deferasirox	59

Demser	60
Desvenlafaxine.....	61
Desvenlafaxine Er.....	61
Dextrose 50%.....	322
Dextrose 70%.....	322
Dhe Nasal.....	62
Diacomit.....	63
Diazepam	64
Diazepam Intensol	64
Dicyclomine Hydrochloride.....	109
Dihydroergotamine Mesylate.....	62, 109
Dipyridamole	109
Disopyramide Phosphate	109
Doptelet.....	65
Doxepin Hydrochloride	270
Doxycycline	196
Drizalma.....	66
Drizalma Sprinkle	66
Dronabinol	67
Droxidopa	184
Dupixent.....	68

E

Eligard.....	70
Emend	322
Emsam.....	71
Enbrel	72
Enbrel Mini	72
Enbrel Sureclick.....	72
Engerix-b.....	322
Enstilar	38
Epclusa.....	73
Epidiolex.....	74
Epo	75
Epoprostenol Sodium.....	322
Eprontia.....	77
Ergotamine.....	78
Ergotamine Tartrate/caffeine	78
Erivedge	79
Erleada	80
Erlotinib	81
Erlotinib Hydrochloride	81
Esbriet	82
Everolimus	83, 322

Exkivity 85

F

Fanapt 86

Fanapt Titration Pack 86

Farydak 87

Fasenra 88

Fasenra Pen 88

Fentanyl 89

Fentanyl Citrate Oral Transmucosal 197

Fentanyl Patch 89

Fetzima 90

Fetzima Titration Pack 90

Fiasp Pumpcart 322

Fingolimod 100

Fintepla 91

Flebogamma Dif 144

Fluoroplex 92

Fluorouracil 41, 92

Fortamet/glumetza 93

Forteo 94

Fotivda 96

Freamine III 322

Fycompa 97

G

Gamastan 322

Gammagard Liquid 144

Gammagard S/d Iga Less Than 1mcg/ml 144

Gammaked 144

Gammaplex 144

Gamunex-c 144

Ganciclovir 322

Gattex 98

Gavreto 99

Gefitinib 140

Gengraf 322

Genotropin 103

Genotropin Miniquick 103

Gilenya 100

Gilotrif 101

Glatiramer 102

Granisetron Hydrochloride 322

Growth Hormone 103

Guanfacine Er 109

Guanfacine Hydrochloride 109

H

Haegarda 105

Harvoni 106

Hepatamine 322

Heplisav-b 322

Hetlioz 107

Hetlioz Lq 108

High Risk Medication - 65 109

Hrm - Anticonvulsants - 65 110

Hrm - Hypnotics - 65 111

Hrm - Scopolamine - 65 112

Hrm-antiparkinson - 65 113

Hrm-cyproheptadine - 65 114

Hrm-hydroxyzine - 65 115

Hrm-hydroxyzine Inj - 65 117

Hrm-promethazine - 65 119

Hrms - Carbinoxamine 120

Hrms - Clemastine- 65 121

Hrms - Teas 122

Hrm-skeletal Muscle Relaxants - 65 123

Humira 124

Humira Pediatric Crohns Disease Starter Pack .. 124

Humira Pen 124

Humira Pen-cd/uc/hs Starter 124

Humira Pen-pediatric Uc Starter Pack 124

Humira Pen-ps/uv Starter 124

Humulin R U-500 (concentrated) 322

Hydrocodone Bitartrate Er 139

Hydromorphone Hcl 322

Hydromorphone Hydrochloride 322

Hydroxyzine Hcl 115, 117

Hydroxyzine Hydrochloride 115, 117

Hydroxyzine Pamoate 115

Hypnotic-benzodiazepines - 65 126

Hysingla Er 139

I

Ibrance 127

Icatibant 128

Icatibant Acetate 128

Iclusig 129

Idhifa	130
Imatinib	131
Imatinib Mesylate.....	131
Imbruvica	132
Imovax Rabies (h.d.c.v.)	322
Inbrija	134
Increlex.....	135
Inlyta.....	136
Inqovi	137
Inrebic.....	138
Ipratropium Bromide.....	322
Ipratropium Bromide/albuterol Sulfate	322
Ir Before Er.....	139
Iressa.....	140
Isolyte-s	322
Isolyte-s Ph 7.4.....	322
Isotretinoin	141
Itraconazole	142
Ivermectin.....	143
Ivermectin Tab	143
Ivig	144

J

Jakafi	146
Javygtor.....	156
Jaypirca.....	147
Jynneos	322

K

Kalydeco.....	148
Kesimpta.....	149
Ketoconazole.....	150
Ketorolac Tromethamine.....	109
Kevzara.....	151
Kisqali	152
Kisqali Femara 200 Dose	152
Kisqali Femara 400 Dose	152
Kisqali Femara 600 Dose	152
Korlym	153
Krazati.....	154
Kristalose.....	155
Kuvan	156

L

Lapatinib.....	157
----------------	-----

Lapatinib Ditosylate.....	157
Lenalidomide	222
Lenvima	158
Lenvima 10 Mg Daily Dose.....	158
Lenvima 12mg Daily Dose	158
Lenvima 14 Mg Daily Dose.....	158
Lenvima 18 Mg Daily Dose.....	158
Lenvima 20 Mg Daily Dose.....	158
Lenvima 24 Mg Daily Dose.....	158
Lenvima 4 Mg Daily Dose.....	158
Lenvima 8 Mg Daily Dose.....	158
Leuprolide	159
Leuprolide Acetate.....	159
Levalbuterol	322
Levalbuterol Hcl	323
Levalbuterol Hydrochloride.....	323
Lidocaine.....	160, 271
Lidocaine Patches	160
Linezolid	161
Lonsurf.....	162
Lorbrena.....	163
Lumakras.....	164
Lupron Depot (1-month).....	166
Lupron Depot (3-month).....	166
Lupron Depot-ped.....	165
Lupron Depot-ped (1-month).....	165
Lupron Depot-ped (3-month).....	165
Lupron Ped.....	165
Lupron-endometriosis	166
Lynparza	167
Lyrica Cr	168
Lytgobi.....	169

M

Mavyret.....	170
Mekinist	171
Mektovi.....	173
Memantine	174
Memantine Hcl Titration Pak.....	174
Memantine Hydrochloride	174
Memantine Hydrochloride Er	174
Mepron.....	175
Metformin Hydrochloride Er	93
Methadone Hcl.....	139

Methadone Hydrochloride	139
Methscopolamine Bromide	109
Methylprednisolone.....	323
Methylprednisolone Acetate	323
Methylprednisolone Sodium Succinate.....	323
Methylprednisolone Sodiumsuccinate	323
Metyrosine.....	60
Modafinil.....	176
Morphine Sulfate.....	323
Morphine Sulfate Er.....	139
Morphine Sulfate/sodium Chloride.....	323
Mycophenolate Mofetil.....	323
Mycophenolic Acid Dr.....	323

N

Natpara	177
Nayzilam	178
Nerlynx.....	179
Nexavar	180
Ninlaro.....	182
Nitisinone	183
Nothera.....	184
Nubeqa	185
Nuedexta.....	186
Nulojix.....	323
Nuplazid	187
Nurtec	188
Nutrilipid.....	323

O

Octagam	144
Octreotide.....	189
Octreotide Acetate.....	189
Odomzo	190
Ofev.....	191
Ojjaara	192
Oncaspar.....	193
Ondansetron Hcl.....	323
Ondansetron Hydrochloride	323
Ondansetron Odt	323
Onureg.....	194
Opsumit.....	195
Oracea.....	196
Oral-intranasal Fentanyl.....	197

Orgovyx	198
Orkambi	199
Orserdu.....	200
Otezla	201
Oxandrolone.....	202
Oxazepam	203
Oxazepam - 65	203
Ozempic	204
Ozempic Aet	204

P

Panzyga.....	144
Part B Versus Part D.....	322
Pegasys.....	205
Pemazyre.....	206
Pentamidine Isethionate	323
Perphenazine/amitriptyline	109
Phenobarbital	110
Phenobarbital Sodium.....	110
Phenylbutyrate	207
Piqray	208
Piqray 200mg Daily Dose.....	208
Piqray 250mg Daily Dose.....	208
Piqray 300mg Daily Dose.....	208
Pirfenidone.....	82
Plenammine.....	323
Pomalyst.....	209
Praluent	210
Prednisolone.....	323
Prednisolone Sodium Phosphate.....	323
Prednisone.....	323
Prednisone Intensol.....	323
Pregabalin	211
Pregabalin Er.....	168
Prehevbrio.....	323
Premasol.....	323
Pretomanid	212
Prevymis	213
Privigen.....	144
Procalamine.....	323
Procrit.....	75
Prograf.....	323
Prolastin-c	9
Promacta	214

Promethazine Hcl	119
Promethazine Hcl Plain.....	119
Promethazine Hydrochloride.....	119
Promethegan.....	119
Prosol.....	323
Protriptyline Hcl.....	122
Pulmozyme.....	216

Q

Qinlock	217
Quetiapine Fumarate Er	218
Quetiapine Xr	218
Quinine Sulfate.....	219

R

Rabavert	323
Recombivax Hb.....	323
Regranex.....	220
Retevmo	221
Revlimid.....	222
Rezlidhia.....	224
Rezurock.....	225
Rinvoq.....	226
Rozlytrek.....	228
Rubraca.....	229
Rufinamide	21
Rybelsus	230
Rybelsus Aet	230
Rydapt	231

S

Sajazir.....	128
Sandimmune.....	323
Sandostatin Lar.....	232
Sandostatin Lar Depot.....	232
Sapropterin Dihydrochloride.....	156
Scemblix.....	234
Scopolamine	112
Signifor.....	235
Sildenafil	236
Sildenafil Citrate	236
Sirolimus	323
Sirturo.....	237
Skyrizi	238, 239
Skyrizi Pen	238

Skyrizi-cd.....	239
Sodium Oxybate.....	310
Sodium Phenylbutyrate.....	207
Somatuline Depot.....	240
Somavert	241
Sorafenib Tosylate	180
Spritam.....	242
Sprycel	243
Stelara	244
Stivarga	245
Sunitinib Malate.....	246
Sutent	246
Symlin.....	247
Symlinpen 120	247
Symlinpen 60	247
Sympazan.....	248
Synribo.....	249

T

Tabrecta.....	250
Tacrolimus	323
Tadalafil	251
Tadalafil (pah).....	251
Tafinlar.....	252
Tagrisso.....	254
Taltz	255
Talzenna.....	256
Targretin Topical	257
Tasigna.....	258
Tasimelteon.....	107
Tazarotene.....	259
Tazorac.....	259
Tazverik	260
Tecfidera	261
Tecfidera Starter Pack.....	261
Tecvayli.....	262
Temazepam	126, 263
Temazepam 30mg - 65.....	263
Tepmetko	264
Testosterone Enanthate	265
Testosterone Enanthate Inj.....	265
Tetrabenazine.....	266
Thalomid.....	267
Thioridazine Hcl	109

Tibsovo.....	268
Tobramycin	269
Topical Doxepin.....	270
Topical Lidocaine.....	271
Topical Tretinoin.....	272
Toremifene	273
Toremifene Citrate	273
Tpn Electrolytes	323
Tracleer.....	27
Tramadol Hcl Er.....	139
Tramadol Hydrochloride Er	139
Travasol.....	323
Tretinoin	272
Tretinoin Microsphere.....	272
Tretinoin Microsphere Pump	272
Triazolam	126
Trientine	274
Trientine Hydrochloride.....	274
Trihexyphenidyl Hcl	113
Trihexyphenidyl Hydrochloride.....	113
Trikafta.....	275
Trimethobenzamide Hydrochloride	109
Trimipramine Maleate.....	122
Trophamine	323
Trulicity.....	276
Trulicity Aet	276
Truseltiq	277
Truxima	278
Tukysa	280
Turalio	281

U

Ukoniq.....	282
-------------	-----

V

Valchlor.....	283
Valtoco	284
Valtoco 10 Mg Dose	284
Valtoco 15 Mg Dose	284
Valtoco 20 Mg Dose	284
Valtoco 5 Mg Dose	284
Vanflyta.....	285
Varenicline Starting Month Box	45
Varenicline Tartrate.....	45

Venclexta	286
Venclexta Starting Pack.....	286
Ventavis	287
Versacloz.....	288
Verzenio.....	289
Victoza	290
Victoza Aet.....	290
Vigabatrin	291
Vigadrone.....	291
Vitrakvi	292
Vizimpro	293
Vonjo.....	294
Voriconazole	295
Vosevi	296
Votrient	297
Vumerity	298

W

Welireg.....	299
--------------	-----

X

Xalkori	300
Xeljanz	301
Xeljanz Xr.....	301
Xermelo.....	302
Xgeva	303
Xhance	304
Xhance Mapd.....	304
Xifaxan.....	305
Xolair	306
Xospata	307
Xpovio.....	308
Xpovio 60 Mg Twice Weekly.....	308
Xpovio 80 Mg Twice Weekly.....	308
Xtandi.....	309
Xyrem	310

Z

Zaleplon	111
Zarxio.....	311
Zejula	312
Zelboraf.....	313
Zenatane.....	141
Zirabev	314
Zolinza	316

Zolpidem Tartrate.....	111
Zonisade	317
Ztalmy	318
Zydelig	319

Zykadia	320
Zyprexa Relprevv.....	321
Zytiga	1