

Aetna Better Health of Virginia (HMO D-SNP), Aetna Medicare Advantra Cares (HMO D-SNP), Aetna Medicare Assure (HMO D-SNP), Aetna Medicare Assure 1 (HMO D-SNP), Aetna Medicare Assure Flex Plan (HMO D-SNP), Aetna Medicare Assure Plan (HMO D-SNP), Aetna Medicare Assure Plan (HMO-POS D-SNP), Aetna Medicare Assure Plus Plan (HMO-POS D-SNP), Aetna Medicare Assure Premier (HMO D-SNP), Aetna Medicare Assure Premier (PPO D-SNP), Aetna Medicare Assure Value (HMO D-SNP), Aetna Medicare Dual Choice (PPO D-SNP), Aetna Medicare Dual Complete Plan (HMO D-SNP), Aetna Medicare Dual Preferred (HMO D-SNP), Aetna Medicare Dual Preferred (PPO D-SNP), Aetna Medicare Dual Preferred Plan (HMO D-SNP), Aetna Medicare Dual Preferred Plan (PPO D-SNP), Aetna Medicare Dual Prime Plan (HMO D-SNP), Aetna Medicare Dual Select (HMO D-SNP), Aetna Medicare Dual Select Choice (PPO D-SNP), Aetna Medicare Dual Signature (HMO D-SNP), Aetna Medicare Dual Signature Choice (PPO D-SNP), Aetna Medicare Dual Signature Select (HMO D-SNP), Aetna Medicare Dual Signature Select (PPO D-SNP), and Aetna Medicare Preferred Plan (HMO D-SNP)

Last Updated: 04/01/2024

ABIRATERONE

Products Affected

- Abiraterone Acetate

- Zytiga TABS 500MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Node-positive (N1), non-metastatic (M0) prostate cancer and very-high-risk prostate cancer.
Exclusion Criteria	N/A
Required Medical Information	The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

NR_0009_3741 09/2014

Formulary ID: 24024: version 12

Other Criteria	N/A
-----------------------	-----

ACITRETIN

Products Affected

- Acitretin

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Prevention of non-melanoma skin cancers in high risk individuals, Lichen planus, Keratosis follicularis (Darier Disease)
Exclusion Criteria	N/A
Required Medical Information	Psoriasis: The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to methotrexate or cyclosporine.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ACTIMMUNE

Products Affected

- Actimmune

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Mycosis fungoides, Sezary syndrome.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ADEMPAS

Products Affected

- Adempas

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): PAH was confirmed by right heart catheterization. For PAH new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, AND 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units. For chronic thromboembolic pulmonary hypertension (CTEPH) (WHO Group 4): 1) Patient has persistent or recurrent CTEPH after pulmonary endarterectomy (PEA), OR 2) Patient has inoperable CTEPH with the diagnosis confirmed by right heart catheterization AND by computed tomography (CT), magnetic resonance imaging (MRI), or pulmonary angiography.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AIMOVIG

Products Affected

- Aimovig

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the preventive treatment of migraine, initial: 1) The patient experienced an inadequate treatment response with a 4-week trial of any one of the following: antiepileptic drugs (AEDs), beta-adrenergic blocking agents, antidepressants OR 2) The patient experienced an intolerance or has a contraindication that would prohibit a 4-week trial of any one of the following: antiepileptic drugs (AEDs), beta-adrenergic blocking agents, antidepressants. For preventive treatment of migraine, continuation: The patient received at least 3 months of treatment with the requested drug, and the patient had a reduction in migraine days per month from baseline.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial: 3 months, Continuation: Plan Year
Other Criteria	N/A

AKEEGA

Products Affected

- Akeega

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ALECENSA

Products Affected

- Alecensa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent ALK-positive non-small cell lung cancer (NSCLC), brain metastases from ALK-positive NSCLC, ALK-positive anaplastic large-cell lymphoma.
Exclusion Criteria	N/A
Required Medical Information	For non-small cell lung cancer (NSCLC): the disease is recurrent, advanced, or metastatic.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ALOSETRON

Products Affected

- Alosetron Hydrochloride

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For severe diarrhea-predominant irritable bowel syndrome (IBS): 1) The requested drug is being prescribed for a biological female or a person that self-identifies as a female, 2) chronic IBS symptoms lasting at least 6 months, 3) gastrointestinal tract abnormalities have been ruled out, AND 4) inadequate response to one conventional therapy (e.g., antispasmodics, antidepressants, antidiarrheals).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ALPHA1-PROTEINASE INHIBITOR

Products Affected

- Prolastin-c

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For alpha1-proteinase inhibitor deficiency: Patient must have 1) clinically evident emphysema, AND 2) pretreatment serum alpha 1-proteinase inhibitor level less than 11 micromol/L (80 mg/dL by radial immunodiffusion or 50 mg/dL by nephelometry).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ALUNBRIG

Products Affected

- Alunbrig

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent anaplastic lymphoma kinase (ALK)-positive non-small cell lung cancer (NSCLC), brain metastases from ALK-positive NSCLC, inflammatory myofibroblastic tumors (IMT) with ALK translocation.
Exclusion Criteria	N/A
Required Medical Information	For non-small cell lung cancer (NSCLC): 1) the disease is recurrent, advanced, or metastatic AND 2) the disease is anaplastic lymphoma kinase (ALK)-positive.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AMBRISENTAN

Products Affected

- Ambrisentan

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): PAH was confirmed by right heart catheterization. For PAH new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, AND 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ARCALYST

Products Affected

- Arcalyst

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Prevention of gout flares in patients initiating or continuing urate-lowering therapy.
Exclusion Criteria	N/A
Required Medical Information	For prevention of gout flares in patients initiating or continuing urate-lowering therapy (e.g., allopurinol) (new starts): 1) two or more gout flares within the previous 12 months, AND 2) inadequate response, intolerance, or contraindication to maximum tolerated doses of a non-steroidal anti-inflammatory drug (NSAID) and colchicine, AND 3) concurrent use with urate-lowering therapy. For prevention of gout flares in patients initiating or continuing urate-lowering therapy (e.g., allopurinol) (continuation): 1) patient must have achieved or maintained a clinical benefit (i.e., a fewer number of gout attacks or fewer flare days) compared to baseline, AND 2) continued use of urate-lowering therapy concurrently with the requested drug. For recurrent pericarditis: patient must have had an inadequate response, intolerance, or contraindication to maximum tolerated doses of an NSAID and colchicine.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ARMODAFINIL

Products Affected

- Armodafinil

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For excessive sleepiness associated with narcolepsy: The diagnosis has been confirmed by sleep lab evaluation. For excessive sleepiness associated with obstructive sleep apnea (OSA): The diagnosis has been confirmed by polysomnography.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ASPARLAS

Products Affected

- Asparlas

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	21 years of age or younger
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AUBAGIO

Products Affected

- Aubagio

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AUGTYRO

Products Affected

- Augtyro

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AUSTEDO

Products Affected

- Austedo
- Austedo Xr
- Austedo Xr Patient Titration Kit

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Tourette's syndrome
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AUVELITY

Products Affected

- Auvelity

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For Major Depressive Disorder (MDD): The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to two of the following: serotonin and norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), mirtazapine, bupropion.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AVONEX AET

Products Affected

- Avonex INJ 30MCG/0.5ML
- Avonex Pen

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

AYVAKIT

Products Affected

- Ayvakit

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Myeloid and lymphoid neoplasms with eosinophilia, gastrointestinal stromal tumor (GIST) for unresectable, recurrent, or metastatic disease without platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation.
Exclusion Criteria	N/A
Required Medical Information	For myeloid and lymphoid neoplasms with eosinophilia, the patient meets all of the following criteria: 1) The disease is FIP1L1- PDGFRA rearrangement-positive, AND 2) The disease harbors a PDGFRA D842A mutation, AND 3) The disease is resistant to imatinib. For GIST, the patient meets either of the following criteria: 1) The disease harbors PDGFRA exon 18 mutation, including PDGFRA D842V mutations, OR 2) The requested drug will be used after failure on at least two Food and Drug Administration (FDA)-approved therapies in unresectable, recurrent, or metastatic disease without PDGFRA exon 18 mutation. For systemic mastocytosis: 1) The patient has a diagnosis of indolent systemic mastocytosis or advanced systemic mastocytosis (including aggressive systemic mastocytosis [ASM], systemic mastocytosis with associated hematological neoplasm [SM-AHN], and mast cell leukemia [MCL]) AND 2) The patient has a platelet count of greater than or equal to 50,000/microliter (mcL).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BALVERSA

Products Affected

- Balversa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For urothelial carcinoma: 1) disease has susceptible fibroblast growth factor receptor 3 (FGFR3) or fibroblast growth factor receptor 2 (FGFR2) genetic alterations AND 2) the requested drug will be used as subsequent therapy for any of the following: a) locally advanced or metastatic urothelial carcinoma, b) recurrent primary carcinoma of the urethra, c) stage II-IV urothelial carcinoma of the bladder, d) urothelial carcinoma of the bladder with metastatic or local recurrence post cystectomy, or e) urothelial carcinoma of the bladder with muscle invasive local recurrence or persistent disease in a preserved bladder.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BANZEL

Products Affected

- Rufinamide

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	1 year of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BENLYSTA

Products Affected

- Benlysta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	For patients new to therapy: severe active central nervous system lupus.
Required Medical Information	For systemic lupus erythematosus (SLE): 1) patient is currently receiving a stable standard therapy regimen (e.g., corticosteroid, antimalarial, or NSAIDs) for SLE, OR 2) patient has experienced an intolerance or has a contraindication to standard therapy regimen for SLE. For lupus nephritis: 1) patient is currently receiving a stable standard therapy regimen (e.g., corticosteroid, cyclophosphamide, mycophenolate mofetil, or azathioprine) for lupus nephritis OR 2) patient has experienced an intolerance or has a contraindication to standard therapy regimen for lupus nephritis.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BERINERT

Products Affected

- Berinert

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For acute angioedema attacks due to hereditary angioedema (HAE): Patient meets either of the following: 1) the patient has HAE with C1 inhibitor deficiency or dysfunction confirmed by laboratory testing OR 2) the patient has HAE with normal C1 inhibitor confirmed by laboratory testing and one of the following: a) the patient tested positive for an F12, angiotensin-1, plasminogen, kininogen-1 (KNG1), heparan sulfate-glucosamine 3-O-sulfotransferase 6 (HS3ST6), or myoferlin (MYOF) gene mutation OR b) the patient has a family history of angioedema and the angioedema was refractory to a trial of high-dose antihistamine therapy for at least one month.
Age Restrictions	5 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
Coverage Duration	Plan Year
Other Criteria	N/A

BESREMI

Products Affected

- Besremi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BETASERON

Products Affected

- Betaseron

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BEXAROTENE

Products Affected

- Bexarotene CAPS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Mycosis fungoides (MF)/Sezary syndrome (SS), CD30-positive primary cutaneous anaplastic large cell lymphoma (ALCL), CD30-positive lymphomatoid papulosis (LyP)
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BOSENTAN

Products Affected

- Bosentan

- Tracleer TBSO

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): PAH was confirmed by right heart catheterization. For PAH new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, AND 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BOSULIF

Products Affected

- Bosulif

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Philadelphia chromosome positive B-cell acute lymphoblastic leukemia (Ph+ B-ALL), myeloid and/or lymphoid neoplasms with eosinophilia and ABL1 rearrangement in the chronic phase or blast phase
Exclusion Criteria	N/A
Required Medical Information	For chronic myeloid leukemia (CML), including patients newly diagnosed with CML and patients who have received a hematopoietic stem cell transplant: 1) Diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene, AND 2) If patient experienced resistance to an alternative tyrosine kinase inhibitor, patient is negative for all of the following mutations: T315I, G250E, V299L, and F317L, AND 3) patient has experienced resistance or intolerance to imatinib or dasatinib. For B-ALL including patient who have received hematopoietic stem cell transplant: 1) Diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene, and 2) If patient experienced resistance to an alternative tyrosine kinase inhibitor, patient is negative for all of the following mutations: T315I, G250E, V299L, and F317L.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BRAFTOVI

Products Affected

- Braftovi CAPS 75MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Adjuvant systemic therapy for cutaneous melanoma, appendiceal adenocarcinoma
Exclusion Criteria	N/A
Required Medical Information	For colorectal cancer (including appendiceal adenocarcinoma): 1) Tumor is positive for BRAF V600E mutation, AND 2) The requested drug will be used for either of the following: a) subsequent therapy for advanced or metastatic disease, b) primary treatment for unresectable metachronous metastases. For melanoma: 1) Tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K), AND 2) The requested drug will be used as a single agent or in combination with binimetinib, AND 3) The requested drug will be used for either of the following: a) unresectable, limited resectable, or metastatic disease, b) adjuvant systemic therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BRIVIACT

Products Affected

- Briviact ORAL SOLN
- Briviact TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For treatment of partial-onset seizures (i.e., focal-onset seizures): 1) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to a generic anticonvulsant AND 2) the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to any of the following: Aptiom (if 4 years of age or older), Xcopri (if 18 years of age or older), Spritam (if 4 years of age or older).
Age Restrictions	1 month of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BRIVIACT INJ

Products Affected

- Briviact INJ

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For treatment of partial-onset seizures (i.e., focal-onset seizures): 1) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to a generic anticonvulsant AND 2) the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to any of the following: Aptiom (if 4 years of age or older), Xcopri (if 18 years of age or older), Spritam (if 4 years of age or older).
Age Restrictions	1 month of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BRONCHITOL

Products Affected

- Bronchitol

- Bronchitol Tolerance Test

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BRUKINSA

Products Affected

- Brukinsa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BUPRENORPHINE

Products Affected

- Buprenorphine Hcl SUBL

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug is being prescribed for the treatment of opioid use disorder AND patient meets one of the following: 1) The patient is pregnant or breastfeeding, and the requested drug is being prescribed for induction therapy and/or subsequent maintenance therapy for treatment of opioid use disorder OR 2) The requested drug is being prescribed for induction therapy for transition from opioid use to treatment of opioid use disorder OR 3) The requested drug is being prescribed for maintenance therapy for treatment of opioid use disorder in a patient who is intolerant to naloxone.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BUPRENORPHINE PATCH

Products Affected

- Buprenorphine PTWK

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug is being prescribed for pain associated with cancer, sickle cell disease, a terminal condition, or pain being managed through palliative care OR the patient meets all of the following: 1) The requested drug is being prescribed for pain severe enough to require daily, around-the-clock, long-term treatment in a patient who has been taking an opioid AND 2) The patient can safely take the requested dose based on their history of opioid use [Note: This drug should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.] AND 3) The patient has been evaluated and the patient will be monitored for the development of opioid use disorder AND 4) This request is for continuation of therapy for a patient who has been receiving an extended-release opioid agent for at least 30 days OR the patient has taken an immediate-release opioid for at least one week.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

BYDUREON

Products Affected

- Bydureon Bcise

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	10 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	The Prior Authorization only applies to patients whose claim is not submitted with an ICD-10 code indicating a diagnosis of type 2 diabetes mellitus OR to patients who do not have a history of an antidiabetic drug (EXCLUDING glucagon-like peptide receptor agonists [GLP-1 RAs] and combination glucose-dependent insulinotropic polypeptide [GIP] and GLP-1 RAs).

BYETTA

Products Affected

- Byetta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	The Prior Authorization only applies to patients whose claim is not submitted with an ICD-10 code indicating a diagnosis of type 2 diabetes mellitus OR to patients who do not have a history of an antidiabetic drug (EXCLUDING glucagon-like peptide receptor agonists [GLP-1 RAs] and combination glucose-dependent insulinotropic polypeptide [GIP] and GLP-1 RAs).

CABOMETYX

Products Affected

- Cabometyx

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Non-small cell lung cancer, Ewing sarcoma, osteosarcoma, gastrointestinal stromal tumor, endometrial carcinoma
Exclusion Criteria	N/A
Required Medical Information	<p>For renal cell carcinoma: The disease is advanced, relapsed, or stage IV. For non-small cell lung cancer: 1) the disease is rearranged during transfection (RET) positive AND 2) the disease is recurrent, advanced, or metastatic. For hepatocellular carcinoma: the requested drug will be used as subsequent treatment. For gastrointestinal stromal tumor (GIST): The patient meets either of the following: 1) the disease is unresectable, recurrent/progressive, or metastatic AND the patient has failed a FDA-approved therapy (e.g., imatinib, sunitinib, regorafenib, ripretinib) OR 2) the requested drug will be used for palliation of symptoms if previously tolerated and effective. For Ewing sarcoma and osteosarcoma: the requested drug will be used as subsequent therapy. For differentiated thyroid cancer (DTC) (follicular, papillary, Hurthle cell): 1) The disease is locally advanced or metastatic disease, 2) the disease has progressed after a vascular endothelial growth factor receptor (VEGFR)- targeted therapy, AND 3) the patient is refractory to radioactive iodine therapy (RAI) or ineligible for RAI. For endometrial carcinoma: 1) the disease is recurrent or metastatic AND 2) the requested drug will be used as subsequent therapy.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CALCIPOTRIENE

Products Affected

- Calcipotriene CREA
- Calcipotriene OINT
- Calcipotriene SOLN
- Calcitrene
- Calcitriol OINT
- Enstilar

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For Treatment of Psoriasis: The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to a topical steroid.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CALQUENCE

Products Affected

- Calquence

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Waldenstrom macroglobulinemia (lymphoplasmacytic lymphoma), marginal zone lymphoma (including extranodal marginal zone lymphoma of the stomach, extranodal marginal zone lymphoma of nongastric sites, nodal marginal zone lymphoma, splenic marginal zone lymphoma)
Exclusion Criteria	N/A
Required Medical Information	For marginal zone lymphoma (including extranodal marginal zone lymphoma of the stomach, extranodal marginal zone lymphoma of nongastric sites, nodal marginal zone lymphoma, and splenic marginal zone lymphoma): the requested drug is being used for the treatment of relapsed, refractory, or progressive disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CAPRELSA

Products Affected

- Caprelsa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Differentiated thyroid carcinoma: papillary, follicular, and Hurthle cell.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CARAC

Products Affected

- Fluorouracil CREA 0.5%

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Patients who are pregnant or may become pregnant. Patients with dihydropyrimidine dehydrogenase (DPD) enzyme deficiency.
Required Medical Information	If being used as part of a compounded product, all active ingredients in the compounded product are Food and Drug Administration (FDA) approved for topical use.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A

CARBAGLU

Products Affected

- Carglumic Acid

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For N-acetylglutamate synthase (NAGS) deficiency: Diagnosis of NAGS deficiency was confirmed by enzymatic, biochemical, or genetic testing.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CAYSTON

Products Affected

- Cayston

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For treatment of respiratory symptoms in cystic fibrosis patients: 1) Pseudomonas aeruginosa is present in the patient's airway cultures, OR 2) The patient has a history of pseudomonas aeruginosa infection or colonization in the airways.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CERDELGA

Products Affected

- Cerdelga

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For type 1 Gaucher disease (GD1): 1) Diagnosis was confirmed by an enzyme assay demonstrating a deficiency of beta-glucocerebrosidase enzyme activity or by genetic testing, and 2) Patient's CYP2D6 metabolizer status has been established using an FDA-cleared test, and 3) Patient is a CYP2D6 extensive metabolizer, an intermediate metabolizer, or a poor metabolizer.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CHANTIX AET

Products Affected

- Varenicline Starting Month Box
- Varenicline Tartrate

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	This Prior Authorization requirement applies after the patient has received a cumulative 24 weeks of therapy in a Plan Year.

CHLORDIAZEPOXIDE - 65

Products Affected

- Chlordiazepoxide Hcl CAPS 10MG, 5MG
- Chlordiazepoxide Hydrochloride CAPS 25MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For all indications: the prescriber must acknowledge the benefit of therapy with the prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For the management of anxiety disorders: 1) the requested drug is being used concurrently with a selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the SSRI/SNRI becomes effective for the symptoms of anxiety, OR 2) the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to AT LEAST TWO agents from the following classes: a) selective serotonin reuptake inhibitors (SSRIs), or b) serotonin-norepinephrine reuptake inhibitors (SNRIs).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Short-term relief anxiety-preop apprehens and anx-1 mo, Anxiety Disorder-4 mo, Alc Withdrawal-PlanYR
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older.

CLOBAZAM

Products Affected

- Clobazam

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Seizures associated with Dravet syndrome
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Seizures associated with Lennox-Gastaut syndrome (LGS): 2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CLOMIPRAMINE

Products Affected

- Clomipramine Hydrochloride

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Depression, panic disorder
Exclusion Criteria	N/A
Required Medical Information	For obsessive-compulsive disorder (OCD) and panic disorder: The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to any of the following: a serotonin and norepinephrine reuptake inhibitor (SNRI), a selective serotonin reuptake inhibitor (SSRI). For depression: The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to two of the following: serotonin and norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), mirtazapine, bupropion.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CLORAZEPATE

Products Affected

- Clorazepate Dipotassium TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For all indications: The prescriber must acknowledge the benefit of therapy with this prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For the management of anxiety disorders: 1) The requested drug is being used concurrently with a selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the SSRI/SNRI becomes effective for the symptoms of anxiety, OR 2) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to AT LEAST TWO agents from the following classes: a) selective serotonin reuptake inhibitors (SSRIs), b) serotonin-norepinephrine reuptake inhibitors (SNRIs).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Short-term relief anxiety-1 month, Anxiety Disorders-4 months, All other Diagnoses-Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older.

CLOZAPINE ODT

Products Affected

- Clozapine Odt

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

COLY-MYCIN

Products Affected

- Colistimethate Sodium INJ

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug is being used intravenously (IV) or intramuscularly (IM). Administration will not be via nebulizer (Note: CMS endorsed compendia do not support inhalation/nebulization of colistimethate.). The infection is proven or strongly suspected to be caused by susceptible bacteria based on: 1) culture and susceptibility information, OR 2) local epidemiology and susceptibility patterns.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist
Coverage Duration	Initial approval: 3 months. Renewal: Plan Year
Other Criteria	N/A

COMETRIQ

Products Affected

- Cometriq

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Non-small cell lung cancer (NSCLC), differentiated thyroid carcinoma: papillary, follicular, and Hurthle cell.
Exclusion Criteria	N/A
Required Medical Information	For NSCLC: The requested medication is used for NSCLC when the patient's disease expresses rearranged during transfection (RET) gene rearrangements.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

COPIKTRA

Products Affected

- Copiktra

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Hepatosplenic T-Cell lymphoma, breast implant-associated anaplastic large cell lymphoma (ALCL), peripheral T-Cell lymphoma
Exclusion Criteria	N/A
Required Medical Information	For chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL), breast implant-associated anaplastic large cell lymphoma (ALCL), and peripheral T-Cell lymphoma: the patient has relapsed or refractory disease. For hepatosplenic T-Cell lymphoma: the patient has refractory disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

COTELLIC

Products Affected

- Cotellic

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Central nervous system (CNS) cancer (i.e., glioma, glioblastoma, astrocytoma, oligodendroglioma), adjuvant systemic therapy for cutaneous melanoma.
Exclusion Criteria	N/A
Required Medical Information	For central nervous system (CNS) cancer (i.e., glioma, glioblastoma, astrocytoma, oligodendroglioma): 1) The tumor is positive for BRAF V600E activating mutation, AND 2) The requested drug will be used in combination with vemurafenib. For melanoma: 1) The tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K), AND 2) The requested drug will be used in combination with vemurafenib, AND 3) The requested drug will be used for either of the following: a) unresectable, limited resectable, or metastatic disease, b) adjuvant systemic therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CYCLOPHOSPHAMIDE ORAL

Products Affected

- Cyclophosphamide CAPS
- Cyclophosphamide TABS

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

CYSTAGON

Products Affected

- Cystagon

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For nephropathic cystinosis: Diagnosis was confirmed by ANY of the following: 1) the presence of increased cystine concentration in leukocytes, OR 2) genetic testing, OR 3) demonstration of corneal cystine crystals by slit lamp examination.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

CYSTARAN

Products Affected

- Cystaran

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For cystinosis: 1) Diagnosis was confirmed by ANY of the following: a) the presence of increased cystine concentration in leukocytes, OR b) genetic testing, OR c) demonstration of corneal cystine crystals by slit lamp examination, AND 2) the patient has corneal cystine crystal accumulation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DALFAMPRIDINE

Products Affected

- Dalfampridine Er

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For multiple sclerosis, patient must meet the following: For new starts, prior to initiating therapy, patient demonstrates sustained walking impairment. For continuation of therapy: patient must have experienced an improvement in walking speed OR other objective measure of walking ability since starting the requested drug.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DAURISMO

Products Affected

- Daurismo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Post induction therapy following response to previous therapy with the same regimen for acute myeloid leukemia (AML). Relapsed/refractory AML as a component of repeating the initial successful induction regimen.
Exclusion Criteria	N/A
Required Medical Information	For acute myeloid leukemia: 1) the requested drug must be used in combination with cytarabine, 2) the patient is 75 years of age or older OR has comorbidities that preclude intensive chemotherapy, AND 3) the requested drug will be used as treatment for induction therapy, post-induction therapy, or relapsed or refractory disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DEFERASIROX

Products Affected

- Deferasirox

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For chronic iron overload due to blood transfusions: pretreatment serum ferritin level is greater than 1000 mcg/L.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DEMSEER

Products Affected

- Metyrosine

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to an alpha-adrenergic antagonist.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DESVENLAFAXINE

Products Affected

- Desvenlafaxine Er

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For Major Depressive Disorder (MDD): The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to TWO of the following: serotonin and norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), mirtazapine, bupropion.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DHE NASAL

Products Affected

- Dihydroergotamine Mesylate NASAL SOLN

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Coverage will be denied when used in conjunction with potent CYP3A4 inhibitors (e.g., ritonavir, nelfinavir, indinavir, erythromycin, clarithromycin).
Required Medical Information	The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one triptan 5-HT1 receptor agonist.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DIACOMIT

Products Affected

- Diacomit

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	6 months of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DIAZEPAM AET

Products Affected

- Diazepam CONC
- Diazepam INJ 5MG/ML
- Diazepam ORAL SOLN 5MG/5ML
- Diazepam TABS
- Diazepam Intensol

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For all indications: The prescriber must acknowledge the benefit of therapy with this prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For the management of anxiety disorders: 1) The requested drug is being used concurrently with a selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the SSRI/SNRI becomes effective for the symptoms of anxiety, OR 2) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to AT LEAST TWO agents from the following classes: a) selective serotonin reuptake inhibitors (SSRIs), b) serotonin-norepinephrine reuptake inhibitors (SNRIs).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Short-term relief anx-1 mo, skeletal muscle spasm-3 mo, Anx Disorders-4 mo, Other Diagnoses-PlanYR
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older.

DOPTELET

Products Affected

- Doptelet

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For thrombocytopenia in patients with chronic liver disease: Untransfused platelet count prior to a scheduled procedure is less than 50,000/mcL. For chronic immune thrombocytopenia (ITP): 1) For new starts: a) Patient has had an inadequate response or is intolerant to prior therapy such as corticosteroids or immunoglobulins, AND b) Untransfused platelet count at any point prior to the initiation of the requested medication is less than 30,000/mcL OR 30,000 to 50,000/mcL with symptomatic bleeding or risk factor(s) for bleeding (e.g., undergoing a medical or dental procedure where blood loss is anticipated, comorbidities such as peptic ulcer disease and hypertension, anticoagulation therapy, profession or lifestyle that predisposes patient to trauma). 2) For continuation of therapy, platelet count response to the requested drug: a) Current platelet count is less than or equal to 200,000/mcL OR b) Current platelet count is greater than 200,000/mcL and less than or equal to 400,000/mcL and dosing will be adjusted to a platelet count sufficient to avoid clinically important bleeding.
Age Restrictions	18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Chronic liver disease: 1 month, ITP initial: 6 months, ITP reauthorization: Plan Year
Other Criteria	N/A

DRIZALMA

Products Affected

- Drizalma Sprinkle

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Cancer pain, chemotherapy-induced neuropathic pain
Exclusion Criteria	N/A
Required Medical Information	1) The patient has tried duloxetine capsules OR 2) The patient is unable to take duloxetine capsules for any reason (e.g., difficulty swallowing capsules, requires nasogastric administration).
Age Restrictions	Generalized Anxiety Disorder - 7 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

DRONABINOL

Products Affected

- Dronabinol

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For nausea and vomiting associated with cancer chemotherapy: The patient has experienced an inadequate treatment response, intolerance, or contraindication to one oral 5-HT ₃ receptor antagonist.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

DUPIXENT

Products Affected

- Dupixent

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>For atopic dermatitis (AD), initial therapy: 1) Patient has moderate-to-severe disease, AND 2) Patient has had an inadequate treatment response to either a topical corticosteroid or a topical calcineurin inhibitor, OR topical corticosteroids and topical calcineurin inhibitors are not advisable for the patient. For AD, continuation of therapy: Patient achieved or maintained positive clinical response. For oral corticosteroid dependent asthma, initial therapy: Patient has inadequate asthma control despite current treatment with both of the following medications: 1) High-dose inhaled corticosteroid AND 2) Additional controller (i.e., long acting beta2-agonist, long-acting, muscarinic antagonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies. For moderate-to-severe asthma, initial therapy: Patient has a baseline blood eosinophil count of at least 150 cells per microliter and their asthma remains inadequately controlled despite current treatment with both of the following medications: 1) Medium-to-high-dose inhaled corticosteroid, AND 2) Additional controller (i.e., long acting beta2-agonist, long-acting muscarinic antagonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies. For asthma, continuation of therapy: Asthma control has improved on treatment with the requested drug, as demonstrated by a reduction in the frequency and/or severity of symptoms and exacerbations or a reduction in the daily maintenance oral corticosteroid dose. For chronic rhinosinusitis with nasal polyposis (CRSwNP): 1) The requested drug is used as add-on maintenance treatment, AND 2) Patient has experienced an inadequate treatment response to Xhance (fluticasone).</p>
Age Restrictions	Atopic Dermatitis: 6 months of age or older, Asthma: 6 years of age or older, Chronic Rhinosinusitis with Nasal Polyposis and Prurigo Nodularis: 18 years of age or older, Eosinophilic Esophagitis: 12 years of age or older

Prescriber Restrictions	N/A
Coverage Duration	AD, initial: 4 months, PN, initial: 6 months, All others: Plan Year
Other Criteria	For eosinophilic esophagitis (EoE), initial therapy: 1) Diagnosis has been confirmed by esophageal biopsy, AND 2) Patient weighs at least 40 kilograms, AND 3) Patient experienced an inadequate treatment response, intolerance, or patient has a contraindication to a topical corticosteroid (e.g., fluticasone propionate or budesonide). For EoE, continuation of therapy: Patient achieved or maintained a positive clinical response. For prurigo nodularis (PN), initial therapy: Patient has had an inadequate treatment response to a topical corticosteroid OR topical corticosteroids are not advisable for the patient. For PN, continuation of therapy: Patient achieved or maintained a positive clinical response.

ELIGARD

Products Affected

- Eligard

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent androgen receptor positive salivary gland tumors
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

EMSAM

Products Affected

- Emsam

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For Major Depressive Disorder (MDD): 1) The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to TWO of the following: serotonin and norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), mirtazapine, bupropion OR 2) The patient is unable to swallow oral formulations.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ENBREL

Products Affected

- Enbrel INJ 25MG/0.5ML, 50MG/ML
- Enbrel Mini
- Enbrel Sureclick

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Hidradenitis suppurativa, non-radiographic axial spondyloarthritis
Exclusion Criteria	N/A
Required Medical Information	For moderately to severely active rheumatoid arthritis (new starts only): 1) patient has experienced an inadequate treatment response, intolerance, or has a contraindication to methotrexate (MTX) OR 2) patient has experienced an inadequate treatment response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. For active ankylosing spondylitis and non-radiographic axial spondyloarthritis (new starts only): patient has experienced an inadequate treatment response or intolerance to a non-steroidal anti-inflammatory drug (NSAID) OR the patient has a contraindication that would prohibit a trial of NSAIDs. For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis AND 2) patient meets any of the following: a) the patient has experienced an inadequate treatment response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, c) patient has severe psoriasis that warrants a biologic as first-line therapy (i.e. at least 10% of the BSA or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected). For hidradenitis suppurativa (new starts only): patient has severe, refractory disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	N/A
-----------------------	-----

ENDARI

Products Affected

- Endari

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	5 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

EPCLUSA

Products Affected

- Epclusa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hepatitis C virus (HCV): Infection confirmed by presence of HCV RNA in the serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of human immunodeficiency virus (HIV) coinfection, presence or absence of resistance-associated substitutions where applicable, transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current American Association for the Study of Liver Diseases and Infectious Diseases Society of America (AASLD-IDSA) treatment guidelines.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Criteria will be applied consistent with current AASLD-IDSA guidance
Other Criteria	N/A

EPIDIOLEX

Products Affected

- Epidiolex

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	1 year of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

EPO

Products Affected

- Procrit

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Anemia due to myelodysplastic syndromes (MDS), anemia in rheumatoid arthritis (RA), anemia due to hepatitis C treatment (ribavirin in combination with either interferon alfa or peginterferon alfa)
Exclusion Criteria	Patients receiving chemotherapy with curative intent. Patients with myeloid cancer.
Required Medical Information	Requirements regarding hemoglobin (Hgb) values exclude values due to a recent transfusion. For initial approval: 1) for all uses except anemia due to chemotherapy or myelodysplastic syndrome (MDS): patient has adequate iron stores (for example, a transferrin saturation [TSAT] greater than or equal to 20%), AND 2) for all uses except surgery: pretreatment (no erythropoietin treatment in previous month) Hgb is less than 10 g/dL, AND 3) for MDS: pretreatment serum erythropoietin level is 500 international units/L or less. For reauthorizations (patient received erythropoietin treatment in previous month) in all uses except surgery: 1) patient has received at least 12 weeks of erythropoietin therapy, AND 2) patient responded to erythropoietin therapy, AND 3) current Hgb is less than 12 g/dL, AND 4) for all uses except anemia due to chemotherapy or MDS: patient has adequate iron stores (for example, a transferrin saturation [TSAT] greater than or equal to 20%).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	16 weeks
Other Criteria	Coverage includes use in anemia in patients whose religious beliefs forbid blood transfusions. Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual (e.g., used for treatment of anemia for a patient with chronic renal failure who is undergoing dialysis, or furnished from physician's supply incident to a physician service).

EPRONTIA

Products Affected

- Eprontia

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For treatment of partial-onset seizures (i.e., focal-onset seizures): 1)The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to a generic anticonvulsant AND 2) the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to any of the following: Aptiom (if 4 years of age or older), Xcopri (if 18 years of age or older), Spritam (if 4 years of age or older). For monotherapy treatment of primary generalized tonic-clonic seizures: 1) The patient has experienced an inadequate treatment response or intolerance to topiramate tablets or capsules, OR 2) The patient has difficulty swallowing solid oral dosage forms (e.g., tablets, capsules). For adjunctive treatment of primary generalized tonic-clonic seizures: 1) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to a generic anticonvulsant AND 2) If the patient is 6 years of age or older, the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to Spritam. For the preventative treatment of migraines: 1) The patient has experienced an inadequate treatment response or intolerance to topiramate tablets or capsules, OR 2) The patient has difficulty swallowing solid oral dosage forms (e.g., tablets, capsules).
Age Restrictions	Epilepsy: 2 years of age or older, Migraine: 12 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ERGOTAMINE

Products Affected

- Ergotamine Tartrate/caffeine

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Coverage will be denied when used in conjunction with potent CYP3A4 inhibitors (e.g., ritonavir, nelfinavir, indinavir, erythromycin, clarithromycin).
Required Medical Information	The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least ONE triptan 5-HT1 agonist.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ERIVEDGE

Products Affected

- Erivedge

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Adult medulloblastoma
Exclusion Criteria	N/A
Required Medical Information	For adult medulloblastoma: patient has received prior systemic therapy AND has tumor(s) with mutations in the sonic hedgehog pathway.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ERLEADA

Products Affected

- Erleada

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ERLOTINIB

Products Affected

- Erlotinib Hydrochloride TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent non-small cell lung cancer (NSCLC), recurrent chordoma, relapsed or stage IV renal cell carcinoma (RCC), brain metastases from non-small cell lung cancer (NSCLC), recurrent pancreatic cancer.
Exclusion Criteria	N/A
Required Medical Information	For NSCLC (including brain metastases from NSCLC): 1) the disease is recurrent, advanced, or metastatic and 2) the patient has sensitizing EGFR mutation-positive disease. For pancreatic cancer: the disease is locally advanced, unresectable, recurrent, or metastatic.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ESBRIET

Products Affected

- Pirfenidone

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For idiopathic pulmonary fibrosis (new starts only): 1) a high-resolution computed tomography (HRCT) study of the chest or a lung biopsy reveals the usual interstitial pneumonia (UIP) pattern, OR 2) HRCT study of the chest reveals a result other than the UIP pattern (e.g., probable UIP, indeterminate for UIP) and the diagnosis is supported either by a lung biopsy or by a multidisciplinary discussion between at least a radiologist and pulmonologist who are experienced in idiopathic pulmonary fibrosis if a lung biopsy has not been conducted.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

EVEROLIMUS

Products Affected

- Everolimus TABS 10MG, 2.5MG, 5MG, 7.5MG

- Everolimus TBSO

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Classic Hodgkin lymphoma, thymomas and thymic carcinomas, previously treated Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma, soft tissue sarcoma (perivascular epithelioid cell tumors (PEComa) and lymphangioliomyomatosis subtypes), gastrointestinal stromal tumors, neuroendocrine tumors of the thymus, well differentiated grade 3 neuroendocrine tumors, thyroid carcinoma (papillary, Hurthle cell, and follicular), endometrial carcinoma, histiocytic neoplasms (Rosai-Dorfman Disease, Erdheim-Chester Disease, Langerhans Cell Histiocytosis)
Exclusion Criteria	N/A
Required Medical Information	For breast cancer: 1) The disease is recurrent unresectable, advanced, or metastatic hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, AND 2) The requested drug is prescribed in combination with exemestane, fulvestrant, or tamoxifen, AND 3) The requested drug is used for subsequent treatment. For renal cell carcinoma: The disease is relapsed, advanced, or stage IV. For subependymal giant cell astrocytoma (SEGA): The requested drug is given as adjuvant treatment. For gastrointestinal stromal tumor: The disease is recurrent/progressive, unresectable, or metastatic AND the patient failed an FDA-approved therapy (e.g., imatinib, sunitinib, regorafenib, ripretinib). For symptomatic or relapsed/refractory Erdheim-Chester Disease (ECD), symptomatic or relapsed/refractory Rosai-Dorfman Disease, and Langerhans Cell Histiocytosis (LCH): the patient must have a phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha (PIK3CA) mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A

Coverage Duration	Plan Year
Other Criteria	N/A

EXKIVITY

Products Affected

- Exkivity

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FANAPT

Products Affected

- Fanapt

- Fanapt Titration Pack

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For treatment of schizophrenia: 1) The patient experienced an inadequate treatment response, intolerance, or has a contraindication to one of the following generic products: aripiprazole, asenapine, lurasidone, olanzapine, quetiapine, risperidone, ziprasidone, AND 2) The patient experienced an inadequate treatment response, intolerance, or has a contraindication to one of the following brand products: Caplyta, Rexulti, Secuado, Vraylar.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FASENRA

Products Affected

- Fasenra

- Fasenra Pen

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Severe asthma, initial therapy: 1) Either a) Patient has baseline blood eosinophil count of at least 150 cells per microliter OR b) Patient is dependent on systemic corticosteroids, and 2) Patient has a history of severe asthma despite current treatment with both of the following medications: a) medium-to-high-dose inhaled corticosteroid and b) additional controller (i.e., long-acting beta2-agonist, long-acting muscarinic antagonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies. Severe asthma, continuation of therapy: Asthma control has improved on treatment with the requested drug, as demonstrated by a reduction in the frequency and/or severity of symptoms and exacerbations or a reduction in the daily maintenance oral corticosteroid dose.
Age Restrictions	12 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FENTANYL PATCH

Products Affected

- Fentanyl PT72

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>The requested drug is being prescribed for pain associated with cancer, sickle cell disease, a terminal condition, or pain being managed through palliative care OR the patient meets all of the following: 1) The requested drug is being prescribed for pain severe enough to require daily, around-the-clock, long-term treatment in a patient who has been taking an opioid AND 2) The patient can safely take the requested dose based on their history of opioid use [Note: This drug should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.] AND 3) The patient has been evaluated and the patient will be monitored for the development of opioid use disorder AND 4) This request is for continuation of therapy for a patient who has been receiving an extended-release opioid agent for at least 30 days OR the patient has taken an immediate-release opioid for at least one week.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FETZIMA

Products Affected

- Fetzima

- Fetzima Titration Pack

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For major depressive disorder (MDD): The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to TWO of the following: serotonin and norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), mirtazapine, bupropion.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FINTEPLA

Products Affected

- Fintepla

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FIRMAGON

Products Affected

- Firmagon INJ 120MG/VIAL, 80MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FLUCYTOSINE

Products Affected

- Flucytosine CAPS 250MG, 500MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 weeks
Other Criteria	N/A

FLUOROURACIL

Products Affected

- Fluorouracil CREA 5%

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Patients who are pregnant or may become pregnant.
Required Medical Information	Applies to new starts only. If being used as part of a compounded product, all active ingredients in the compounded product are Food and Drug Administration (FDA) approved for topical use.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A

FORTEO

Products Affected

- Teriparatide INJ 620MCG/2.48ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>For postmenopausal osteoporosis: patient has ONE of the following: 1) history of fragility fracture, OR 2) pre-treatment T-score of less than or equal to -2.5 or pre-treatment T-score greater than -2.5 and less than -1 with a high pre-treatment Fracture Risk Assessment Tool (FRAX) fracture probability AND patient has ANY of the following: a) indicators for higher fracture risk (e.g., advanced age, frailty, glucocorticoid therapy, very low T-scores, or increased fall risk), OR b) patient has failed prior treatment with or is intolerant to a previous injectable osteoporosis therapy OR c) patient has had an oral bisphosphonate trial of at least 1-year duration or there is a clinical reason to avoid treatment with an oral bisphosphonate. For primary or hypogonadal osteoporosis in men: patient has ONE of the following: 1) history of osteoporotic vertebral or hip fracture, OR 2) pre-treatment T-score of less than or equal to -2.5, or pre-treatment T-score greater than -2.5 and less than -1 with a high pre-treatment FRAX fracture probability AND patient has ANY of the following: a) patient has failed prior treatment with or is intolerant to a previous injectable osteoporosis therapy, OR b) patient has had an oral bisphosphonate trial of at least 1-year duration or there is a clinical reason to avoid treatment with an oral bisphosphonate.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial: 24 months, Continuation: Plan Year

Other Criteria	For glucocorticoid-induced osteoporosis: Patient has had an oral bisphosphonate trial of at least 1-year duration unless patient has a contraindication or intolerance to an oral bisphosphonate, AND patient meets ANY of the following: 1) patient has a history of fragility fracture, OR 2) a pre-treatment T-score of less than or equal to -2.5, OR 3) pre-treatment T-score greater than -2.5 and less than -1 with a high pre-treatment FRAX fracture probability. Continuation of therapy: If the patient has received greater than or equal to 24 months of therapy with any parathyroid hormone analog: 1) The patient remains at or has returned to having a high risk for fracture, AND 2) The benefit of therapy with this prescribed medication outweighs the potential risks for this patient.
-----------------------	--

FOTIVDA

Products Affected

- Fotivda

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For advanced renal cell carcinoma: the following criteria must be met: 1) The disease is relapsed or refractory, 2) The requested drug must be used after at least two prior systemic therapies, and 3) The patient has experienced disease progression or an intolerable adverse event with a trial of Cabometyx (cabozantinib).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FRUZAQLA

Products Affected

- Fruzaqla

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

FYCOMPA

Products Affected

- Fycompa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For treatment of partial-onset seizures (i.e., focal-onset seizures): 1) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to a generic anticonvulsant AND 2) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to any of the following: Aptiom, Xcopri, Spritam. For adjunctive treatment of primary generalized tonic-clonic seizures: 1) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to a generic anticonvulsant AND 2) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to Spritam.
Age Restrictions	Partial-onset seizures (i.e., focal-onset seizures): 4 years of age or older. Primary generalized tonic-clonic seizures: 12 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

GATTEX

Products Affected

- Gattex

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For short bowel syndrome (SBS) initial therapy: 1) If the request is for an adult patient, the patient has been dependent on parenteral support for at least 12 months OR 2) If the request is for a pediatric patient, the patient is dependent on parenteral support. For SBS continuation: Requirement for parenteral support has decreased from baseline while on therapy with the requested drug.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, gastrointestinal surgeon, or nutritional support specialist.
Coverage Duration	Plan Year
Other Criteria	N/A

GAVRETO

Products Affected

- Gavreto

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent rearranged during transfection (RET) rearrangement-positive non-small cell lung cancer
Exclusion Criteria	N/A
Required Medical Information	For non-small cell lung cancer, patient must meet all of the following: 1) The disease is recurrent, advanced, or metastatic, and 2) The tumor is rearranged during transfection (RET) fusion-positive or RET rearrangement-positive.
Age Restrictions	Non-small cell lung cancer: 18 years of age or older. Medullary thyroid cancer and thyroid cancer: 12 years of age or older.
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

GILENYA

Products Affected

- Fingolimod

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

GILOTRIF

Products Affected

- Gilotrif

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For non-small cell lung cancer (NSCLC): Patient meets either of the following: 1) Patient has metastatic squamous NSCLC that progressed after platinum-based chemotherapy, OR 2) Patient has sensitizing epidermal growth factor receptor (EGFR) mutation-positive disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

GLATIRAMER

Products Affected

- Copaxone INJ 20MG/ML, 40MG/ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

GROWTH HORMONE

Products Affected

- Genotropin

- Genotropin Miniquick

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	Pediatric patients with closed epiphyses
Required Medical Information	Pediatric growth hormone deficiency (GHD): Patient (pt) is a neonate or was diagnosed with GHD as a neonate OR meets any of the following: 1) younger than 2.5 years old (yo) with pre-treatment (pre-tx) height (ht) more than 2 standard deviations (SD) below mean and slow growth velocity OR 2) 2.5 yo or older AND one of the following: a) pre-tx 1-year ht velocity more than 2 SD below mean OR b) pre-tx ht more than 2 SD below mean and 1-year ht velocity more than 1 SD below mean, AND patient meets any of the following: 1) failed 2 pre-tx growth hormone (GH) stimulation tests (peak below 10 ng/mL), OR 2) pituitary/central nervous system (CNS) disorder (e.g., genetic defects, acquired structural abnormalities, congenital structural abnormalities) and pre-tx insulin-like growth factor-1 (IGF-1) more than 2 SD below mean. Turner syndrome (TS): 1) Confirmed by karyotyping AND 2) pre-tx ht is less than the 5th percentile for age. Small for gestational age (SGA): 1) Birth weight (wt) less than 2500g at gestational age (GA) greater than 37 weeks, OR birth wt or length below 3rd percentile for GA or at least 2 SD below mean for GA, AND 2) did not manifest catch-up growth by age 2.
Age Restrictions	SGA: 2 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist, nephrologist, infectious disease specialist, gastroenterologist/nutritional support specialist, or geneticist.
Coverage Duration	Plan Year

Other Criteria	Adult GHD: Pt meets any of the following: 1) failed 2 pre-tx GH stimulation tests, OR 2) pre-tx IGF-1 more than 2 SD below mean AND failed 1 pre-tx GH stimulation test. (Note: Stimulation tests include: a) insulin tolerance test [ITT] [peak GH less than or equal to 5 ng/ml], or b) Macrilen-stimulation test [peak GH level less than 2.8ng/ml], or c) glucagon-stimulation test [GST] [peak GH level less than or equal to 3 ng/ml] for pt with a body mass index [BMI] 25-30 kg/m ² and high pretest probability of GHD [e.g., acquired structural abnormalities] or BMI less than 25 kg/m ² , or d) GST [peak GH level less than or equal to 1 ng/ml] in pt with BMI 25-30 kg/m ² and low pretest probability of GHD or BMI greater than 30 kg/m ²), OR 3) organic hypothalamic-pituitary disease (e.g., suprasellar mass with previous surgery and cranial irradiation) with 3 or more pituitary hormone deficiencies AND pre-tx IGF-1 more than 2 SD below mean, OR 4) genetic or structural hypothalamic-pituitary defects, OR 5) childhood-onset GHD with congenital (genetic or structural) abnormality of the hypothalamus/pituitary/CNS. Renewal for pediatric GHD, TS, SGA, and adult GHD: Patient is experiencing improvement.
-----------------------	--

HAEGARDA

Products Affected

- Haegarda

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the prevention of acute angioedema attacks due to hereditary angioedema (HAE): The patient meets either of the following: 1) the patient has hereditary angioedema (HAE) with C1 inhibitor deficiency or dysfunction confirmed by laboratory testing OR 2) the patient has hereditary angioedema with normal C1 inhibitor confirmed by laboratory testing and either of the following: a) patient tested positive for an F12, angiotensin-1, plasminogen, kininogen-1 (KNG1), heparan sulfate-glucosamine 3-O-sulfotransferase 6 (HS3ST6), or myoferlin (MYOF) gene mutation OR b) patient has a family history of angioedema and the angioedema was refractory to a trial of high-dose antihistamine therapy for at least one month.
Age Restrictions	6 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
Coverage Duration	Plan Year
Other Criteria	N/A

HARVONI

Products Affected

- Harvoni

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hepatitis C virus (HCV): Infection confirmed by presence of HCV RNA in the serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of human immunodeficiency virus (HIV) coinfection, presence or absence of resistance-associated substitutions where applicable, transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current American Association for the Study of Liver Diseases and Infectious Diseases Society of America (AASLD-IDSA) treatment guidelines.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Criteria applied consistent w/ current AASLD-IDSA guidance. Reminder for 8wk option if appropriate.
Other Criteria	N/A

HETLIOZ

Products Affected

- Tasimelteon

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For Non-24-Hour Sleep-Wake Disorder: 1) For initial therapy and continuation of therapy the patient must meet both of the following: a) diagnosis of total blindness in both eyes (e.g., nonfunctioning retinas) and b) unable to perceive light in either eye, AND 2) If currently on therapy with the requested drug, patient must meet at least one of the following: a) increased total nighttime sleep or b) decreased daytime nap duration. For nighttime sleep disturbances in Smith-Magenis Syndrome (SMS): 1) For initial therapy and continuation therapy, the patient has a confirmed diagnosis of SMS AND 2) If currently on therapy with the requested drug, the patient experienced improvement in the quality of sleep since starting therapy.
Age Restrictions	Non-24: 18 years of age or older. SMS: 16 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with a sleep disorder specialist, neurologist, or psychiatrist.
Coverage Duration	Initiation: 6 Months, Renewal: Plan Year
Other Criteria	N/A

HETLIOZ LQ

Products Affected

- HetlioZ Lq

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For nighttime sleep disturbances in Smith-Magenis Syndrome (SMS): 1) For initial therapy and continuation therapy, the patient has a confirmed diagnosis of SMS AND 2) If currently on therapy with the requested drug, the patient experienced improvement in the quality of sleep since starting therapy.
Age Restrictions	3 to 15 years of age
Prescriber Restrictions	Prescribed by or in consultation with a sleep disorder specialist or neurologist
Coverage Duration	Initiation: 6 Months, Renewal: Plan Year.
Other Criteria	N/A

HIGH RISK MEDICATION - 65

Products Affected

- Dicyclomine Hcl SOLN
- Dicyclomine Hydrochloride CAPS
- Dicyclomine Hydrochloride INJ
- Dicyclomine Hydrochloride TABS
- Dihydroergotamine Mesylate INJ
- Dipyridamole TABS
- Disopyramide Phosphate CAPS
- Guanfacine Er TB24 2MG
- Guanfacine Hydrochloride TABS
- Guanfacine Hydrochloride TB24 1MG, 3MG, 4MG
- Ketorolac Tromethamine TABS
- Methscopolamine Bromide TABS
- Perphenazine/amitriptyline
- Thioridazine Hcl TABS 100MG, 10MG, 25MG, 50MG
- Trimethobenzamide Hydrochloride

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM - ANTICONVULSANTS - 65

Products Affected

- Phenobarbital ELIX 20MG/5ML
- Phenobarbital TABS 100MG, 15MG, 16.2MG, 30MG, 32.4MG, 60MG, 64.8MG, 97.2MG
- Phenobarbital Sodium INJ 130MG/ML, 65MG/ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Epilepsy
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM - HYPNOTICS - 65

Products Affected

- Zaleplon

- Zolpidem Tartrate TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For insomnia: 1) The patient meets one of the following: a) the patient has a contraindication to the non-HRM (non-High Risk Medication) alternative drug doxepin (3 mg or 6 mg) OR b) The non-HRM (non-High Risk Medication) alternative drug doxepin (3 mg or 6 mg) has been tried AND the patient experienced an inadequate treatment response OR intolerance to the non-HRM (non-High Risk Medication) alternative drug doxepin (3 mg or 6 mg) AND 2) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient AND 3) If the patient is using two or more additional central nervous system (CNS) active medications (e.g., lorazepam, quetiapine, sertraline, clonazepam, escitalopram, alprazolam) with the requested drug, the prescriber has determined that taking multiple central nervous system (CNS) active medications is medically necessary for the patient [Note: Use of multiple central nervous system (CNS) active medications in older adults is associated with an increased risk of falls.].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prior authorization applies to greater than cumulative 90 days of therapy per year.

HRM - SCOPOLAMINE - 65

Products Affected

- Scopolamine PT72 1MG/3DAYS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Excessive salivation
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM-ANTIPARKINSON - 65

Products Affected

- Benztropine Mesylate TABS
- Trihexyphenidyl Hcl SOLN
- Trihexyphenidyl Hydrochloride

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. EPS (extrapyramidal symptoms): 1) The patient has not tried the non-HRM alternative drug amantadine AND 2) The patient has a contraindication to the non-HRM alternative drug amantadine OR 3) The patient has tried the non-HRM alternative drug amantadine AND 4) The patient experienced an inadequate treatment response OR intolerance to the non-HRM alternative drug amantadine. Parkinson's: 1) The patient has tried two of the following non-HRM alternative drugs: amantadine, carbidopa/levodopa, pramipexole, or ropinirole. AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: amantadine, carbidopa/levodopa, pramipexole, or ropinirole.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM-CYPROHEPTADINE - 65

Products Affected

- Cyproheptadine Hcl SYRP
- Cyproheptadine Hydrochloride TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Pruritus, spasticity due to spinal cord injury
Exclusion Criteria	N/A
Required Medical Information	The prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. For rhinitis: 1) The patient has tried two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM-DOXEPIN - 65

Products Affected

- Doxepin Hcl CAPS 75MG
- Doxepin Hcl CONC
- Doxepin Hydrochloride CAPS 100MG, 10MG, 150MG, 25MG, 50MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. For depression: 1) The patient tried two of the following alternative drugs: SSRIs (selective serotonin reuptake inhibitors), SNRIs (serotonin-norepinephrine reuptake inhibitors), bupropion, mirtazapine, or trazodone AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following alternative drugs: SSRIs (selective serotonin reuptake inhibitors), SNRIs (serotonin-norepinephrine reuptake inhibitors), bupropion, mirtazapine, or trazodone. For anxiety: 1) The patient has tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline, or venlafaxine extended-release AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline, or venlafaxine extended-release.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM-HYDROXYZINE - 65

Products Affected

- Hydroxyzine Hcl TABS 50MG
- Hydroxyzine Hydrochloride SYRP
- Hydroxyzine Hydrochloride TABS 10MG, 25MG
- Hydroxyzine Pamoate CAPS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For anxiety: 1) The patient has tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline, or venlafaxine extended-release AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline, or venlafaxine extended-release OR 3) The patient has not tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release AND 4) The patient has acute anxiety. For all indications: 1) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. AND 2) If the patient is taking one or more additional anticholinergic medications (e.g., oxybutynin, meclizine, paroxetine, amitriptyline, dicyclomine, cyclobenzaprine) with the requested drug, the prescriber has determined that taking multiple anticholinergic medications is medically necessary for the patient [Note: Use of multiple anticholinergic medications in older adults is associated with an increased risk of cognitive decline.].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)
-----------------------	--

HRM-HYDROXYZINE INJ - 65

Products Affected

- Hydroxyzine Hcl INJ 25MG/ML

- Hydroxyzine Hydrochloride INJ

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. For Alcohol Withdrawal Syndrome: 1) The patient has not tried one of the following alternative drugs: clorazepate or lorazepam AND 2) The patient has a contraindication to one of the following alternative drugs: clorazepate or lorazepam OR 3) The patient has tried one of the following alternative drugs: clorazepate or lorazepam AND 4) The patient experienced an inadequate treatment response OR intolerance to one of the following alternative drugs: clorazepate or lorazepam. For anxiety: 1) The patient has tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release OR 3) The patient has not tried two of the following alternative drugs: buspirone, duloxetine, escitalopram, sertraline or venlafaxine extended-release AND 4) The patient has acute anxiety.</p>
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	<p>This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)</p>

HRM-PROMETHAZINE - 65

Products Affected

- Promethazine Hcl INJ
- Promethazine Hcl SUPP 12.5MG, 25MG
- Promethazine Hcl TABS 12.5MG
- Promethazine Hydrochloride TABS 25MG, 50MG
- Promethazine Hydrochloride Plain
- Promethegan

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. For rhinitis: 1) The patient has tried two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRMS - ANTIDEPRESSANTS

Products Affected

- Amitriptyline Hcl TABS 100MG, 150MG, 25MG, 75MG
- Amitriptyline Hydrochloride TABS 100MG, 10MG, 50MG
- Desipramine Hydrochloride
- Imipramine Hcl TABS 25MG, 50MG
- Imipramine Hydrochloride TABS 10MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Neuropathic pain
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. For depression: 1) The patient tried two of the following alternative drugs: SSRIs (selective serotonin reuptake inhibitors), SNRIs (serotonin-norepinephrine reuptake inhibitors), bupropion, mirtazapine, or trazodone AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following alternative drugs: SSRIs (selective serotonin reuptake inhibitors), SNRIs (serotonin-norepinephrine reuptake inhibitors), bupropion, mirtazapine, or trazodone.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRMS - CARBINOXAMINE

Products Affected

- Carbinoxamine Maleate SOLN
- Carbinoxamine Maleate TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. For rhinitis: 1) The patient has tried two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRMS - CLEMASTINE- 65

Products Affected

- Clemastine Fumarate TABS 2.68MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient. For rhinitis: 1) The patient has tried two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following non-HRM alternative drugs: levocetirizine, azelastine nasal, fluticasone nasal, or flunisolide nasal.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRMS - TCAS

Products Affected

- Protriptyline Hcl

- Trimipramine Maleate CAPS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For depression: 1) The patient tried two of the following alternative drugs: SSRIs (selective serotonin reuptake inhibitors), SNRIs (serotonin-norepinephrine reuptake inhibitors), bupropion, mirtazapine, or trazodone AND 2) The patient experienced an inadequate treatment response OR intolerance to two of the following alternative drugs: SSRIs (selective serotonin reuptake inhibitors), SNRIs (serotonin-norepinephrine reuptake inhibitors), bupropion, mirtazapine, or trazodone AND 3) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)

HRM-SKELETAL MUSCLE RELAXANTS - 65

Products Affected

- Chlorzoxazone TABS 500MG
- Cyclobenzaprine Hydrochloride TABS 10MG, 5MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1) Prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for this patient AND 2) If the patient is using one or more additional anticholinergic medications (e.g., oxybutynin, meclizine, paroxetine, amitriptyline, dicyclomine, hydroxyzine) with the requested drug, the prescriber has determined that taking multiple anticholinergic medications is medically necessary for the patient [Note: Use of multiple anticholinergic medications in older adults is associated with an increased risk of cognitive decline.].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. (The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) Prior authorization applies to greater than cumulative 90 days of therapy per year.

HUMIRA

Products Affected

- Humira INJ 10MG/0.1ML, 20MG/0.2ML, 40MG/0.4ML, 40MG/0.8ML
- Humira Pediatric Crohns Disease Starter Pack INJ 0, 80MG/0.8ML
- Humira Pen
- Humira Pen-cd/uc/hs Starter INJ 80MG/0.8ML
- Humira Pen-pediatric Uc Starter Pack
- Humira Pen-ps/uv Starter

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderately to severely active rheumatoid arthritis (new starts only): 1) patient has experienced an inadequate treatment response, intolerance, or has a contraindication to methotrexate (MTX) OR 2) patient has experienced an inadequate treatment response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. For active ankylosing spondylitis and non-radiographic axial spondyloarthritis (new starts only): patient has experienced an inadequate treatment response or intolerance to a non-steroidal anti-inflammatory drug (NSAID) OR the patient has a contraindication that would prohibit a trial of NSAIDs. For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) the patient meets any of the following: a) the patient has experienced an inadequate treatment response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, c) the patient has severe psoriasis that warrants a biologic as first-line therapy (i.e., at least 10% of the BSA or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected).
Age Restrictions	N/A
Prescriber Restrictions	N/A

Coverage Duration	Plan Year
Other Criteria	For non-infectious intermediate, posterior and panuveitis (new starts only): 1) patient has experienced an inadequate treatment response or intolerance to a corticosteroid OR 2) the patient has a contraindication that would prohibit a trial of corticosteroids.

HYPNOTIC-BENZODIAZEPINES - 65

Products Affected

- Temazepam CAPS 15MG, 22.5MG, 7.5MG
- Triazolam TABS 0.125MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For short-term treatment of insomnia: 1) Prescriber must acknowledge the benefit of therapy with this prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) AND 2) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to doxepin (3 mg or 6 mg).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older. Prior authorization applies to greater than cumulative 90 days of therapy per year.

IBRANCE

Products Affected

- Ibrance

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Unresectable well-differentiated/dedifferentiated liposarcoma of the retroperitoneum, recurrent hormone receptor-positive human epidermal growth factor receptor 2 (HER2)-negative breast cancer
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ICATIBANT

Products Affected

- Icatibant Acetate

- Sajazir

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For acute angioedema attacks due to hereditary angioedema (HAE): Patient meets either of the following: 1) the patient has HAE with C1 inhibitor deficiency or dysfunction confirmed by laboratory testing OR 2) the patient has HAE with normal C1 inhibitor confirmed by laboratory testing and one of the following: a) the patient tested positive for an F12, angiotensin-1, plasminogen, kininogen-1 (KNG1), heparan sulfate-glucosamine 3-O-sulfotransferase 6 (HS3ST6), or myoferlin (MYOF) gene mutation OR b) the patient has a family history of angioedema and the angioedema was refractory to a trial of high-dose antihistamine therapy for at least one month.
Age Restrictions	18 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an immunologist, allergist, or rheumatologist
Coverage Duration	Plan Year
Other Criteria	N/A

ICLUSIG

Products Affected

- Iclusig

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Myeloid and/or lymphoid neoplasms with eosinophilia and FGFR1 or ABL1 rearrangement in the chronic phase or blast phase
Exclusion Criteria	N/A
Required Medical Information	For chronic myeloid leukemia (CML), including patients who have received a hematopoietic stem cell transplant: 1) patient has accelerated or blast phase CML and no other kinase inhibitor is indicated OR 2) patient has chronic phase CML and has experienced resistance or intolerance to at least 2 prior kinase inhibitors AND at least one of those was imatinib or dasatinib OR 3) patient is positive for the T315I mutation. For acute lymphoblastic leukemia (ALL), including patients who have received a hematopoietic stem cell transplant: diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

IDADACIO

Products Affected

- Adalimumab-aacf (2 Pen)
- Idacio (2 Pen)
- Idacio (2 Syringe)
- Idacio Starter Package For Crohns Disease
- Idacio Starter Package For Plaque Psoriasis

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderately to severely active rheumatoid arthritis (new starts only): 1) patient has experienced an inadequate treatment response, intolerance, or has a contraindication to methotrexate (MTX) OR 2) patient has experienced an inadequate treatment response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. For active ankylosing spondylitis and non-radiographic axial spondyloarthritis (new starts only): patient has experienced an inadequate treatment response or intolerance to a non-steroidal anti-inflammatory drug (NSAID) OR the patient has a contraindication that would prohibit a trial of NSAIDs. For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) the patient meets any of the following: a) the patient has experienced an inadequate treatment response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, c) the patient has severe psoriasis that warrants a biologic as first-line therapy (i.e., at least 10% of the BSA or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	For non-infectious intermediate, posterior and panuveitis (new starts only): 1) patient has experienced an inadequate treatment response or intolerance to a corticosteroid OR 2) the patient has a contraindication that would prohibit a trial of corticosteroids.
-----------------------	--

IDHIFA

Products Affected

- Idhifa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Newly-diagnosed acute myeloid leukemia
Exclusion Criteria	N/A
Required Medical Information	For acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation: 1) patient is 60 years of age or older with newly-diagnosed AML and meets one of the following: a) patient is not a candidate for intensive induction therapy, or b) patient declines intensive induction chemotherapy, OR 2) patient is 60 years of age or older and the requested drug will be used as post-induction therapy following response to induction therapy with the requested drug, OR 3) patient has relapsed or refractory AML.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

IMATINIB

Products Affected

- Imatinib Mesylate

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Desmoid tumors, pigmented villonodular synovitis/tenosynovial giant cell tumor (PVNS/TGCT), recurrent chordoma, melanoma, Kaposi sarcoma, chronic graft versus host disease (cGVHD), T-cell acute lymphoblastic leukemia with ABL-class translocation, aggressive systemic mastocytosis for well-differentiated systemic mastocytosis (WDSM) or when eosinophilia is present with FIP1L1-PDGFR A fusion gene, myeloid and/or lymphoid neoplasms with eosinophilia and ABL1, FIP1L1-PDGFR A, or PDGFR B rearrangement in the chronic phase or blast phase
Exclusion Criteria	N/A
Required Medical Information	For chronic myeloid leukemia (CML) or Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL), including patients who have received a hematopoietic stem cell transplant: diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene. For CML: patient did not fail (excluding failure due to intolerance) prior therapy with a tyrosine kinase inhibitor. For melanoma: c-Kit mutation is positive.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

IMBRUVICA

Products Affected

- Imbruvica

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Hairy cell leukemia, lymphoplasmacytic lymphoma, primary central nervous system (CNS) lymphoma, Human Immunodeficiency Virus (HIV) -related B-cell lymphoma, diffuse large B-cell lymphoma, post-transplant lymphoproliferative disorders, high-grade B-cell lymphoma, mantle cell lymphoma, marginal zone lymphoma (including extranodal marginal zone lymphoma of the stomach, extranodal marginal zone lymphoma of nongastric sites, nodal marginal zone lymphoma, splenic marginal zone lymphoma)
Exclusion Criteria	N/A
Required Medical Information	For mantle cell lymphoma: 1) the requested drug will be used as second-line or subsequent therapy, OR 2) the requested drug will be used in combination with rituximab as pretreatment to induction therapy with RHyperCVAD (rituximab, cyclophosphamide, vincristine, doxorubicin, and dexamethasone) regimen, OR 3) the requested drug will be used as aggressive induction therapy. For marginal zone lymphoma (including extranodal marginal zone lymphoma of the stomach, extranodal marginal zone lymphoma of nongastric sites, nodal marginal zone lymphoma, and splenic marginal zone lymphoma): the requested drug will be used as second-line or subsequent therapy. For hairy cell leukemia: the requested drug will be used as a single agent for disease progression. For primary CNS lymphoma: 1) the disease is relapsed or refractory, OR 2) the requested drug is used for induction therapy as a single agent. For diffuse large B-cell lymphoma and high-grade B-cell lymphoma: the requested drug will be used as second-line or subsequent therapy. For HIV-related B-cell lymphoma: the requested drug will be used as a single agent and as second-line or subsequent therapy for relapsed disease. For post-transplant lymphoproliferative disorders: the requested drug will be used in patients who have received prior chemoimmunotherapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A

Coverage Duration	Plan Year
Other Criteria	N/A

INBRIJA

Products Affected

- Inbrija

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For initial treatment of "off" episodes in Parkinson's disease: 1) The patient is currently being treated with oral carbidopa/levodopa, 2) The patient does not have any of the following: asthma, chronic obstructive pulmonary disease (COPD), or other chronic underlying lung disease. For continuation treatment of "off" episodes in Parkinson's disease: The patient is experiencing improvement on the requested drug.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

INCRELEX

Products Affected

- Increlex

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Pediatric patients with closed epiphyses
Required Medical Information	For growth failure due to severe primary insulin-like growth factor-1 (IGF-1) deficiency or growth hormone (GH) gene deletion in patients who have developed neutralizing antibodies to GH, patient meets all of the following prior to beginning therapy with the requested drug (new starts only): 1) height 3 or more standard deviations (SD) below the mean for children of the same age and gender AND 2) basal IGF-1 level 3 or more SD below the mean for children of the same age and gender AND 3) provocative growth hormone test showing a normal or elevated growth hormone level. For growth failure due to severe primary IGF-1 deficiency or GH gene deletion in patients who have developed neutralizing antibodies to GH, continuation of therapy: patient is experiencing improvement.
Age Restrictions	2 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	Plan Year
Other Criteria	N/A

INLYTA

Products Affected

- Inlyta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Thyroid carcinoma (papillary, Hurthle cell, or follicular), alveolar soft part sarcoma
Exclusion Criteria	N/A
Required Medical Information	For renal cell carcinoma: The disease is advanced, relapsed, or stage IV.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

INQOVI

Products Affected

- Inqovi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

INREBIC

Products Affected

- Inrebic

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and janus kinase 2 (JAK2) rearrangement, accelerated phase myelofibrosis, blast phase myelofibrosis/acute myeloid leukemia
Exclusion Criteria	N/A
Required Medical Information	For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and JAK2 rearrangement: the disease is in chronic or blast phase.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

IR BEFORE ER

Products Affected

- Hydrocodone Bitartrate Er T24A
- Hysingla Er
- Methadone Hcl INJ
- Methadone Hcl ORAL SOLN
- Methadone Hcl TABS
- Methadone Hydrochloride CONC
- Morphine Sulfate Er TBCR 100MG, 200MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug is being prescribed for pain associated with cancer, sickle cell disease, a terminal condition, or pain being managed through palliative care OR the patient meets all of the following: 1) The requested drug is being prescribed for pain severe enough to require daily, around-the-clock, long-term treatment in a patient who has been taking an opioid AND 2) The patient can safely take the requested dose based on their history of opioid use [Note: This drug should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.] AND 3) The patient has been evaluated and the patient will be monitored for the development of opioid use disorder AND 4) This request is for continuation of therapy for a patient who has been receiving an extended-release opioid agent for at least 30 days OR the patient has taken an immediate-release opioid for at least one week.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

IRESSA

Products Affected

- Gefitinib

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Sensitizing epidermal growth factor receptor (EGFR) mutation-positive recurrent non-small cell lung cancer (NSCLC).
Exclusion Criteria	N/A
Required Medical Information	For NSCLC: 1) disease must be metastatic, advanced, or recurrent and 2) patient must have a sensitizing EGFR mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ISOTRETINOIN

Products Affected

- Accutane
- Amnesteem
- Claravis
- Isotretinoin CAPS
- Zenatane

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Refractory acne vulgaris, severe refractory rosacea, neuroblastoma, cutaneous T-cell lymphoma (CTCL) (e.g., mycosis fungoides, Sezary syndrome), high risk for developing skin cancer (squamous cell cancers), transient acantholytic dermatosis (Grover's Disease), keratosis follicularis (Darier Disease), lamellar ichthyosis, pityriasis rubra pilaris.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ITRACONAZOLE

Products Affected

- Itraconazole CAPS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Coccidioidomycosis, Coccidioidomycosis prophylaxis in HIV infection,, Cryptococcosis, Microsporidiosis, Talaromycosis (formerly Penicilliosis), Histoplasmosis prophylaxis in HIV infection, Invasive fungal infection prophylaxis in liver transplant, chronic granulomatous disease (CGD), and hematologic malignancy, Sporotrichosis, Pityriasis versicolor, Tinea versicolor, Tinea corporis, Tinea cruris, Tinea capitis, Tinea manuum, Tinea pedis, primary treatment for allergic bronchopulmonary aspergillosis, primary treatment for chronic cavitary or subacute invasive (necrotizing) pulmonary aspergillosis
Exclusion Criteria	N/A
Required Medical Information	The requested drug will be used orally. For the treatment of onychomycosis due to dermatophytes (Tinea unguium), the diagnosis has been confirmed by a fungal diagnostic test (e.g., potassium hydroxide [KOH] preparation, fungal culture, or nail biopsy). For primary treatment of allergic bronchopulmonary aspergillosis, the requested drug is initiated in combination with systemic corticosteroids.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Disseminated/CNS histo, histo/CM/CGD ppx, chronic cavitary/necrotizing PA: 12 mths. Others: 6 mths
Other Criteria	N/A

IVERMECTIN TAB

Products Affected

- Ivermectin TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Ascariasis, Cutaneous larva migrans, Mansonelliasis, Scabies, Gnathostomiasis, Pediculosis
Exclusion Criteria	N/A
Required Medical Information	The requested drug is not being prescribed for the prevention or treatment of coronavirus disease 2019 (COVID-19).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 month
Other Criteria	N/A

IVIG

Products Affected

- Gammaked INJ 10GM/100ML, 1GM/10ML, 20GM/200ML, 5GM/50ML
- Gamunex-c
- Octagam INJ 10GM/100ML, 10GM/200ML, 1GM/20ML, 2.5GM/50ML, 20GM/200ML, 2GM/20ML, 30GM/300ML, 5GM/100ML, 5GM/50ML
- Priviligen

PA Criteria	Criteria Details
Indications	All Medically-accepted Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For B-cell chronic lymphocytic leukemia (CLL): 1) serum IgG less than 500 mg/dL OR 2) a history of recurrent bacterial infections. For bone marrow transplant/hematopoietic stem cell transplant (BMT/HSCT): 1) IVIG is requested within the first 100 days post-transplant OR 2) serum IgG less than 400 mg/dL. For pediatric human immunodeficiency virus (HIV) infection: 1) serum IgG less than 400 mg/dL OR 2) history of recurrent bacterial infections. For dermatomyositis and polymyositis: 1) at least one standard first-line treatment (corticosteroid or immunosuppressant) has been tried but was unsuccessful or not tolerated OR 2) patient is unable to receive standard therapy because of a contraindication or other clinical reason. For pure red cell aplasia (PRCA): PRCA is secondary to parvovirus B19 infection.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

IWILFIN

Products Affected

- Iwilfin

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

JAKAFI

Products Affected

- Jakafi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Lower-risk myelofibrosis, accelerated phase myelofibrosis, blast phase myelofibrosis/acute myeloid leukemia, acute lymphoblastic leukemia (ALL), chronic myelomonocytic leukemia (CMML)-2, myelodysplastic syndrome/myeloproliferative neoplasm (MDS/MPN) with neutrophilia, essential thrombocythemia, and myeloid, lymphoid or mixed lineage neoplasms with eosinophilia and JAK2 rearrangement
Exclusion Criteria	N/A
Required Medical Information	For polycythemia vera: patient had an inadequate response or intolerance to interferon therapy or hydroxyurea. For acute lymphoblastic leukemia: patient has a cytokine receptor-like factor 2 (CRLF2) mutation or a mutation associated with activation of the Janus kinase/signal transducers and activators of transcription (JAK/STAT) pathway. For CMML-2: the requested drug is used in combination with a hypomethylating agent. For myelodysplastic syndrome/myeloproliferative neoplasm (MDS/MPN) with neutrophilia: the requested drug is used as a single agent or in combination with a hypomethylating agent. For essential thrombocythemia: patient had an inadequate response or loss of response to hydroxyurea, interferon therapy, or anagrelide. For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and JAK2 rearrangement: the disease is in chronic or blast phase.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

JAYPIRCA

Products Affected

- Jaypirca

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL): The patient meets both of the following: 1) The patient has received prior treatment with one of the following: Imbruvica (ibrutinib), Brukinsa (zanubrutinib), or Calquence (acalabrutinib), AND 2) The patient has received prior treatment with a B-cell lymphoma 2 (BCL-2) inhibitor.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KALYDECO

Products Affected

- Kalydeco

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For cystic fibrosis (CF): The requested medication will not be used in combination with other medications containing ivacaftor.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KESIMPTA

Products Affected

- Kesimpta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KETOCONAZOLE

Products Affected

- Ketoconazole TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Cushing's syndrome
Exclusion Criteria	Acute or chronic liver disease. Concurrent use with drugs that are contraindicated with ketoconazole tablets: dofetilide, quinidine, pimozide, cisapride, methadone, disopyramide, dronedarone, ranolazine, ergot alkaloids, irinotecan, lurasidone, oral midazolam, alprazolam, triazolam, felodipine, nisoldipine, tolvaptan, eplerenone, lovastatin, simvastatin, or colchicine.
Required Medical Information	The potential benefits outweigh the risks of treatment with oral ketoconazole. For systemic fungal infections, the patient has any of the following diagnoses: blastomycosis, coccidioidomycosis, histoplasmosis, chromomycosis, or paracoccidioidomycosis. For Cushing's syndrome: the requested drug is being prescribed for a patient who cannot tolerate surgery or where surgery has not been curative.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

KEVZARA

Products Affected

- Kevzara

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderately to severely active rheumatoid arthritis (new starts only): 1) patient has had an inadequate response, intolerance or contraindication to methotrexate (MTX) OR 2) patient has had an inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. For polymyalgia rheumatica (PMR) (new starts only): 1) The patient has experienced an inadequate treatment response to corticosteroids OR 2) The patient has experienced a disease flare while attempting to taper corticosteroids.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KISQALI

Products Affected

- Kisqali
- Kisqali Femara 200 Dose
- Kisqali Femara 400 Dose
- Kisqali Femara 600 Dose

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent hormone receptor-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer, in combination with an aromatase inhibitor, or fulvestrant.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KORLYM

Products Affected

- Korlym

- Mifepristone TABS 300MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	Plan Year
Other Criteria	N/A

KOSELUGO

Products Affected

- Koselugo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	BRAF fusion or BRAF V600E activating mutation-positive recurrent or progressive pilocytic astrocytoma
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	For neurofibromatosis type 1: 2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KRAZATI

Products Affected

- Krazati

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

KRISTALOSE

Products Affected

- Kristalose

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For constipation: 1) The patient has experienced an inadequate treatment response to a one month trial of generic lactulose solution, OR 2) The patient has experienced an intolerance that would prohibit a one month trial of generic lactulose solution, OR 3) the patient has a contraindication to an inactive ingredient which is not contained in the requested drug.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LAPATINIB

Products Affected

- Lapatinib Ditosylate

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Brain metastases from human epidermal growth factor receptor 2 (HER2)-positive breast cancer, recurrent HER2-positive breast cancer, recurrent epidermal growth factor receptor (EGFR)-positive chordoma, HER2-amplified and RAS and BRAF wild-type colorectal cancer (including appendiceal adenocarcinoma).
Exclusion Criteria	N/A
Required Medical Information	For breast cancer, the patient meets all the following: a) the disease is recurrent, advanced, or metastatic (including brain metastases), b) the disease is human epidermal growth factor receptor 2 (HER2)-positive, c) the requested drug will be used in combination with any of the following: 1) aromatase inhibitor, 2) capecitabine, OR 3) trastuzumab. For colorectal cancer: 1) requested drug will be used in combination with trastuzumab and 2) patient has not had previous treatment with a HER2 inhibitor.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LENVIMA

Products Affected

- Lenvima 10 Mg Daily Dose
- Lenvima 12mg Daily Dose
- Lenvima 14 Mg Daily Dose
- Lenvima 18 Mg Daily Dose
- Lenvima 20 Mg Daily Dose
- Lenvima 24 Mg Daily Dose
- Lenvima 4 Mg Daily Dose
- Lenvima 8 Mg Daily Dose

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Medullary thyroid carcinoma, recurrent endometrial carcinoma, thymic carcinoma
Exclusion Criteria	N/A
Required Medical Information	For differentiated thyroid cancer (follicular, papillary, or Hurthle cell): disease is not amenable to radioactive iodine therapy and unresectable, locally recurrent, persistent, or metastatic. For hepatocellular carcinoma: disease is unresectable or inoperable, local, metastatic or with extensive liver tumor burden. For renal cell carcinoma, the disease is advanced, relapsed, or stage IV. For endometrial carcinoma, the patient meets ALL of the following: 1) The disease is advanced, recurrent, or metastatic, 2) The requested drug will be used in combination with pembrolizumab, 3) The patient experienced disease progression following prior systemic therapy, AND 4) The patient is not a candidate for curative surgery or radiation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LEUPROLIDE

Products Affected

- Leuprolide Acetate INJ 1MG/0.2ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Use in combination with growth hormone for children with growth failure and advancing puberty, recurrent androgen receptor positive salivary gland tumors, central precocious puberty.
Exclusion Criteria	N/A
Required Medical Information	For central precocious puberty (CPP): Patients not currently receiving therapy must meet all of the following criteria: 1) Diagnosis of CPP was confirmed by a pubertal response to a gonadotropin releasing hormone (GnRH) agonist test OR a pubertal level of a third generation luteinizing hormone (LH) assay, AND 2) Assessment of bone age versus chronological age supports the diagnosis of CPP, AND 3) The onset of secondary sexual characteristics occurred prior to 8 years of age for female patients OR prior to 9 years of age for male patients.
Age Restrictions	CPP: Patient must be less than 12 years old if female and less than 13 years old if male
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LIDOCAINE PATCHES

Products Affected

- Lidocaine PTCH 5%
- Lidocan III

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Pain associated with diabetic neuropathy, pain associated with cancer-related neuropathy (including treatment-related neuropathy [e.g., neuropathy associated with radiation treatment or chemotherapy]).
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LINEZOLID

Products Affected

- Linezolid

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The infection is proven or strongly suspected to be caused by susceptible bacteria based on: 1) culture and susceptibility information, OR 2) local epidemiology and susceptibility patterns. The patient will be using the requested drug orally or intravenously.
Age Restrictions	N/A
Prescriber Restrictions	Prescribed or directed by an Infectious Disease specialist when being converted from intravenous (IV) linezolid (Zyvox)
Coverage Duration	28 days
Other Criteria	N/A

LONSURF

Products Affected

- Lonsurf

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For colorectal cancer (including appendiceal adenocarcinoma): The disease is advanced or metastatic. For gastric or gastroesophageal junction adenocarcinoma, all of the following criteria must be met: 1) The disease is unresectable locally advanced, recurrent, or metastatic, and 2) The patient has been previously treated with at least two prior lines of chemotherapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LORBRENA

Products Affected

- Lorbrena

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Anaplastic lymphoma kinase (ALK)-positive recurrent non-small cell lung cancer (NSCLC). Repressor of silencing (ROS)-1 rearrangement-positive recurrent, advanced, or metastatic NSCLC following progression on crizotinib, entrectinib, or ceritinib. Symptomatic or relapsed/refractory ALK-positive Erdheim-Chester Disease. Inflammatory myofibroblastic tumor (IMT) with ALK translocation.
Exclusion Criteria	N/A
Required Medical Information	For recurrent, advanced, or metastatic NSCLC: Patient has ALK-positive disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LUMAKRAS

Products Affected

- Lumakras

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent KRAS G12C-positive non-small cell lung cancer (NSCLC)
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LUPRON PED

Products Affected

- Lupron Depot-ped (1-month)
- Lupron Depot-ped (3-month)
- Lupron Depot-ped (6-month)

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For central precocious puberty (CPP): Patients not currently receiving therapy must meet all of the following criteria: 1) Diagnosis of CPP was confirmed by a pubertal response to a gonadotropin releasing hormone (GnRH) agonist test OR a pubertal level of a third generation luteinizing hormone (LH) assay, AND 2) Assessment of bone age versus chronological age supports the diagnosis of CPP, AND 3) The onset of secondary sexual characteristics occurred prior to 8 years of age for female patients OR prior to 9 years of age for male patients.
Age Restrictions	CPP: Patient must be less than 12 years old if female and less than 13 years old if male
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LUPRON-ENDOMETRIOSIS

Products Affected

- Lupron Depot (1-month) INJ 3.75MG
- Lupron Depot (3-month) INJ 11.25MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Breast cancer, epithelial ovarian cancer/fallopian tube cancer/primary peritoneal cancer, androgen receptor positive recurrent salivary gland tumor
Exclusion Criteria	N/A
Required Medical Information	For retreatment of endometriosis, the requested drug is used in combination with norethindrone acetate. For uterine fibroids, patient must meet one of the following: 1) Diagnosis of anemia (e.g., hematocrit less than or equal to 30 percent and/or hemoglobin less than or equal to 10g/dL), OR 2) the requested medication will be used prior to surgery for uterine fibroids. For breast cancer, the requested drug is used for hormone receptor (HR)-positive disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Fibroids: 3 months (mo), max 6 mo total. Endometriosis: 6 mo, max 12 mo total. Others: Plan Year
Other Criteria	N/A

LYNPARZA

Products Affected

- Lynparza TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent HER2-negative, BRCA 1/2-germline mutated breast cancer, recurrent or metastatic HER2-positive, BRCA 1/2-germline mutated breast cancer, uterine leiomyosarcoma.
Exclusion Criteria	N/A
Required Medical Information	For recurrent or metastatic breast cancer: the disease is BRCA 1/2-germline mutated. For prostate cancer: 1) The patient has a BRCA mutation and the requested drug will be used in combination with abiraterone and either prednisone or prednisolone OR 2) The patient has progressed on prior treatment with an androgen receptor-directed therapy. For epithelial ovarian, fallopian tube, or primary peritoneal cancer: The requested drug is used for maintenance therapy for stage II-IV or recurrent disease who are in complete or partial response to chemotherapy. For uterine leiomyosarcoma: 1) the patient has had at least one prior therapy AND 2) the patient has BRCA-altered disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LYRICA CR

Products Affected

- Pregabalin Er

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to gabapentin.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

LYTGOBI

Products Affected

- Lytgobi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Extrahepatic cholangiocarcinoma
Exclusion Criteria	N/A
Required Medical Information	For cholangiocarcinoma: 1) patient has a diagnosis of unresectable, locally advanced or metastatic cholangiocarcinoma, 2) patient has received a previous treatment, AND 3) patient has a disease that has a fibroblast growth factor receptor 2 (FGFR2) gene fusion or other rearrangement.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

MAVYRET

Products Affected

- Mavyret

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Decompensated cirrhosis/moderate or severe hepatic impairment (Child Turcotte Pugh [CTP] class B or C).
Required Medical Information	For hepatitis C virus (HCV): Infection confirmed by presence of HCV RNA in the serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [CTP class B or C]), presence or absence of human immunodeficiency virus (HIV) coinfection, presence or absence of resistance-associated substitutions where applicable, transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current American Association for the Study of Liver Diseases and Infectious Diseases Society of America (AASLD-IDSA) treatment guidelines.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Criteria will be applied consistent with current AASLD-IDSA guidance
Other Criteria	N/A

MEKINIST

Products Affected

- Mekinist

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Langerhans cell histiocytosis, Erdheim-Chester disease, Rosai-Dorfman disease.
Exclusion Criteria	N/A
Required Medical Information	For melanoma: 1) The tumor is positive for a BRAF V600 activating mutation (e.g., V600E or V600K), AND 2) The requested drug will be used as a single agent or in combination with dabrafenib, AND 3) The requested drug will be used for either of the following: a) unresectable, limited resectable, or metastatic disease, b) adjuvant systemic therapy. For central nervous system (CNS) cancer (i.e., glioma, oligodendroglioma, astrocytoma, glioblastoma), non-small cell lung cancer, solid tumors, and anaplastic thyroid cancer: 1) The tumor is positive for a BRAF V600E mutation, AND 2) The requested drug will be used in combination with dabrafenib. For uveal melanoma: The requested drug will be used as a single agent. For ovarian cancer, fallopian tube cancer, and primary peritoneal cancer: The requested drug will be used to treat persistent or recurrent disease. For gallbladder cancer, intrahepatic cholangiocarcinoma, and extrahepatic cholangiocarcinoma: 1) The tumor is positive for a BRAF V600E mutation, AND 2) The disease is unresectable or metastatic, AND 3) The requested drug will be used in combination with dabrafenib. For papillary, follicular, and hurthle cell thyroid carcinoma: 1) The disease is positive for BRAF V600E mutation, AND 2) The disease is not amenable to radioactive iodine (RAI) therapy, AND 3) The requested drug will be used in combination with dabrafenib.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

MEKTOVI

Products Affected

- Mektovi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Adjuvant systemic therapy for cutaneous melanoma, Langerhans Cell Histiocytosis
Exclusion Criteria	N/A
Required Medical Information	For melanoma: 1) The tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K), AND 2) The requested drug will be used in combination with encorafenib, AND 3) The requested drug will be used for either of the following: a) unresectable, limited resectable, or metastatic disease, b) adjuvant systemic therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

MEMANTINE

Products Affected

- Memantine Hcl Titration Pak
- Memantine Hydrochloride SOLN 2MG/ML
- Memantine Hydrochloride TABS
- Memantine Hydrochloride Er

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This prior authorization only applies to patients less than 30 years of age.

MEPRON AET

Products Affected

- Atovaquone SUSP

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Babesiosis, Toxoplasmosis, Pneumocystis jirovecii pneumonia prophylaxis in pediatric patients, mild-to-moderate Pneumocystis jirovecii pneumonia treatment in pediatric patients.
Exclusion Criteria	N/A
Required Medical Information	For the treatment of mild-to-moderate Pneumocystis jirovecii pneumonia (PCP): the patient had an intolerance or has a contraindication to sulfamethoxazole/trimethoprim (SMX-TMP). For the prevention of PCP and primary toxoplasmosis prophylaxis indications: A) the patient had an intolerance or has a contraindication to SMX-TMP, AND B) the patient has had a CD4 cell count of less than 200 cells per cubic millimeter within the past 3 months. For secondary toxoplasmosis prophylaxis: the patient has had a CD4 cell count of less than 200 cells per cubic millimeter within the past 6 months. For babesiosis treatment: the requested drug is used concurrently with azithromycin.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Secondary toxoplasmosis prophylaxis: 6 months, All other indications: 3 months
Other Criteria	N/A

METFORMIN ER

Products Affected

- Metformin Hydrochloride Er TB24 500MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient has experienced an intolerance with a 4-week trial of generic Glucophage XR.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

METHYLTESTOSTERONE

Products Affected

- Methyltestosterone CAPS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to alternative testosterone products (e.g., topical testosterone, transdermal testosterone, injectable testosterone). For primary hypogonadism or hypogonadotropic hypogonadism, initial therapy: The patient has at least two confirmed low morning serum total testosterone concentrations based on the reference laboratory range or current practice guidelines [Note: Safety and efficacy of testosterone products in patients with "age-related hypogonadism" (also referred to as "late-onset hypogonadism") have not been established.]. For primary hypogonadism or hypogonadotropic hypogonadism, continuation of therapy: The patient had a confirmed low morning serum total testosterone concentration based on the reference laboratory range or current practice guidelines before starting testosterone therapy [Note: Safety and efficacy of testosterone products in patients with "age-related hypogonadism" (also referred to as "late-onset hypogonadism") have not been established.].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

MODAFINIL

Products Affected

- Modafinil TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For excessive sleepiness associated with narcolepsy: The diagnosis has been confirmed by sleep lab evaluation. For excessive sleepiness associated with obstructive sleep apnea (OSA): The diagnosis has been confirmed by polysomnography.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

MOUNJARO

Products Affected

- Mounjaro

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	The Prior Authorization only applies to patients whose claim is not submitted with an ICD-10 code indicating a diagnosis of type 2 diabetes mellitus OR to patients who do not have a history of an antidiabetic drug (EXCLUDING glucagon-like peptide receptor agonists [GLP-1 RAs] and combination glucose-dependent insulinotropic polypeptide [GIP] and GLP-1 RAs).

NATPARA

Products Affected

- Natpara

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Acute postsurgical hypoparathyroidism (within 6 months of surgery) and expected recovery from hypoparathyroidism.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NAYZILAM

Products Affected

- Nayzilam

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	12 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NERLYNX

Products Affected

- Nerlynx

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent human epidermal growth factor receptor 2 (HER2)-positive breast cancer, brain metastases from HER2-positive breast cancer.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NEXAVAR

Products Affected

- Nexavar

- Sorafenib Tosylate TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Acute myeloid leukemia, soft tissue sarcoma (angiosarcoma, desmoid tumors/aggressive fibromatosis, and solitary fibrous tumor subtypes), gastrointestinal stromal tumor, medullary thyroid carcinoma, osteosarcoma, recurrent chordoma, epithelial ovarian cancer, fallopian tube cancer, primary peritoneal cancer, lymphoid, myeloid, or mixed lineage neoplasms with eosinophilia
Exclusion Criteria	N/A
Required Medical Information	For acute myeloid leukemia: the disease is FMS-like tyrosine kinase 3-internal tandem duplication (FLT3-ITD) mutation-positive AND either of the following is met (1 OR 2): 1) the requested drug will be used as maintenance therapy after hematopoietic stem cell transplant, OR 2) the requested drug is used in combination with azacitidine or decitabine for low-intensity treatment induction or post-induction therapy AND either a) the patient has is 60 years of age or older or b) the disease is relapsed/refractory. For thyroid carcinoma: histology is follicular, papillary, Hurthle cell or medullary. For gastrointestinal stromal tumor (GIST): the patient meets either of the following: 1) the disease is unresectable, recurrent/progressive, or metastatic AND the patient has failed on an FDA-approved therapy (e.g., imatinib, sunitinib, regorafenib, ripretinib) OR 2) the requested drug is being used for palliation of symptoms if previously tolerated and effective. For renal cell carcinoma: the disease is advanced. For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia: 1) the disease has a FLT3 rearrangement AND 2) the disease is in chronic or blast phase.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	N/A
-----------------------	-----

NINLARO

Products Affected

- Ninlaro

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Relapsed/refractory systemic light chain amyloidosis, Waldenstrom macroglobulinemia, lymphoplasmacytic lymphoma
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NITISINONE

Products Affected

- Nitisinone

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hereditary tyrosinemia type 1 (HT-1): Diagnosis of HT-1 is confirmed by one of the following: 1) biochemical testing (e.g., detection of succinylacetone in urine) OR 2) DNA testing (mutation analysis).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NORTHERA

Products Affected

- Droxidopa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For neurogenic orthostatic hypotension (nOH): Prior to initial therapy, patient has a persistent, consistent decrease in systolic blood pressure of at least 20 mmHg OR decrease in diastolic blood pressure of at least 10 mmHg within 3 minutes of standing or head-up tilt test. For continuation of therapy for nOH, patient must experience a sustained reduction in symptoms of nOH (i.e., decrease in dizziness, lightheadedness, or feeling faint). For both initial and continuation of therapy for nOH, the requested drug will be used for patients with neurogenic orthostatic hypotension associated with one of the following diagnoses: 1) primary autonomic failure due to Parkinson's disease, multiple system atrophy, or pure autonomic failure, OR 2) dopamine beta-hydroxylase deficiency, OR 3) non-diabetic autonomic neuropathy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	N/A

NUBEQA

Products Affected

- Nubeqa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NUEDEXTA

Products Affected

- Nuedexta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NUPLAZID

Products Affected

- Nuplazid CAPS
- Nuplazid TABS 10MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hallucinations and delusions associated with Parkinson's disease psychosis, the diagnosis of Parkinson's disease must be made prior to the onset of psychotic symptoms.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

NURTEC

Products Affected

- Nurtec

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Acute migraine treatment: The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to one triptan 5-HT1 receptor agonist . Preventive treatment of migraine, initial: The patient meets either of the following: 1) The patient experienced an inadequate treatment response with a 4-week trial of any one of the following: Antiepileptic drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants OR 2) The patient experienced an intolerance or has a contraindication that would prohibit a 4-week trial of any one of the following: Antiepileptic drugs (AEDs), Beta-adrenergic blocking agents, Antidepressants. Preventive treatment of migraine, continuation: The patient received at least 3 months of treatment with the requested drug, and the patient had a reduction in migraine days per month from baseline.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Preventive treatment of migraine - initial: 3 months, All other indications: Plan Year
Other Criteria	N/A

OCTREOTIDE

Products Affected

- Octreotide Acetate INJ 1000MCG/ML, 100MCG/ML, 200MCG/ML, 500MCG/ML, 50MCG/ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Tumor control of thymomas and thymic carcinomas.
Exclusion Criteria	N/A
Required Medical Information	For acromegaly, initial: 1) Patient has a high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range, AND 2) Patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For acromegaly, continuation of therapy: Patient's IGF-1 level has decreased or normalized since initiation of therapy. For tumor control of thymomas and thymic carcinomas: The requested drug will be used for any of the following: 1) locally advanced or metastatic disease, 2) postoperatively following tumor resection.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ODOMZO

Products Affected

- Odomzo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

OFEV

Products Affected

- Ofev

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For idiopathic pulmonary fibrosis (new starts only): 1) a high-resolution computed tomography (HRCT) study of the chest or a lung biopsy reveals the usual interstitial pneumonia (UIP) pattern, OR 2) HRCT study of the chest reveals a result other than the UIP pattern (e.g., probable UIP, indeterminate for UIP) and the diagnosis is supported either by a lung biopsy or by a multidisciplinary discussion between at least a radiologist and pulmonologist who are experienced in idiopathic pulmonary fibrosis if a lung biopsy has not been conducted.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

OGSIVEO

Products Affected

- Ogsiveo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

OJJAARA

Products Affected

- Ojjaara

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

OMEGA-3

Products Affected

- Omega-3-acid Ethyl Esters CAPS
375MG; 465MG; 1GM

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hypertriglyceridemia: Prior to the start of treatment with a triglyceride lowering drug, the patient has/had a pretreatment triglyceride level greater than or equal to 500 mg/dL.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ONCASPAR

Products Affected

- Oncaspar

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Extranodal natural killer/T-cell lymphoma, hepatosplenic T-cell lymphoma
Exclusion Criteria	N/A
Required Medical Information	For extranodal natural killer/T-cell lymphoma, acute lymphoblastic leukemia, and hepatosplenic T-cell lymphoma: the requested drug must be used in conjunction with multi-agent chemotherapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ONUREG

Products Affected

- Onureg

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

OPSUMIT

Products Affected

- Opsumit

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): PAH was confirmed by right heart catheterization. For PAH new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, AND 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) Pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ORACEA AET

Products Affected

- Doxycycline CPDR
- Oracea

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to a formulary generic topical metronidazole.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ORAL-INTRANASAL FENTANYL

Products Affected

- Fentanyl Citrate Oral Transmucosal

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1) The requested drug is indicated for the treatment of breakthrough cancer-related pain only. The requested drug is being prescribed for the management of breakthrough pain in a cancer patient with underlying cancer pain AND 2) The International Classification of Diseases (ICD) diagnosis code provided supports the cancer-related diagnosis. [Note: For drug coverage approval, ICD diagnosis code provided MUST support the cancer-related diagnosis.] AND 3) The patient is currently receiving, and will continue to receive, around-the-clock opioid therapy for underlying cancer pain AND 4) The requested drug is intended only for use in opioid tolerant patients. The patient can safely take the requested dose based on their current opioid use history. [Note: Patients considered opioid tolerant are those who are taking around-the-clock medicine consisting of at least 60 mg of oral morphine per day, at least 25 mcg per hour of transdermal fentanyl, at least 30 mg of oral oxycodone per day, at least 60 mg of oral hydrocodone per day, at least 8 mg of oral hydromorphone per day, at least 25 mg of oral oxymorphone per day, or an equianalgesic dose of another opioid medication daily for one week or longer.].
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ORGOVYX

Products Affected

- Orgovyx

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ORKAMBI

Products Affected

- Orkambi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For cystic fibrosis (CF): The requested medication will not be used in combination with other medications containing ivacaftor.
Age Restrictions	1 year of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ORSERDU

Products Affected

- Orserdu

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent hormone receptor positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer
Exclusion Criteria	N/A
Required Medical Information	Breast cancer: 1) the disease is estrogen receptor (ER) positive, human epidermal growth factor receptor 2 (HER2)-negative, and ESR1 mutated AND 2) the patient meets either of the following: a) the disease is advanced, recurrent, or metastatic AND the patient has disease progression following at least one line of endocrine therapy OR b) the disease had no response to preoperative systemic therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

OTEZLA

Products Affected

- Otezla

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For plaque psoriasis (new starts only): Patient meets either of the following: 1) Inadequate treatment response or intolerance to ANY of the following: a) a topical therapy (e.g., topical corticosteroids, calcineurin inhibitors, vitamin D analogs), b) phototherapy (e.g., UVB, PUVA), or c) pharmacologic treatment with methotrexate, cyclosporine, or acitretin, OR 2) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

OXANDROLONE

Products Affected

- Oxandrolone TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Cachexia associated with AIDS (HIV wasting). To enhance growth in patients with Turners Syndrome
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Turners Syndrome: Plan Year, All other diagnoses: 6 months
Other Criteria	Coverage will be denied if request is for an indication excluded from Medicare Part D.

OXAZEPAM - 65

Products Affected

- Oxazepam

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For all indications: The prescriber must acknowledge the benefit of therapy with this prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For the management of anxiety disorders, anxiety associated with depression, and the management of anxiety, tension, agitation and irritability in older patients: 1) The requested drug is being used concurrently with a selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the SSRI/SNRI becomes effective for the symptoms of anxiety, OR 2) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to AT LEAST TWO agents from the following classes: a) selective serotonin reuptake inhibitors (SSRIs), b) serotonin-norepinephrine reuptake inhibitors (SNRIs).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Short-term relief anxiety-1 month, Anxiety Disorders-4 months, Alcohol Withdrawal-Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older.

OZEMPIC

Products Affected

- Ozempic

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	The Prior Authorization only applies to patients whose claim is not submitted with an ICD-10 code indicating a diagnosis of type 2 diabetes mellitus OR to patients who do not have a history of an antidiabetic drug (EXCLUDING glucagon-like peptide receptor agonists [GLP-1 RAs] and combination glucose-dependent insulintropic polypeptide [GIP] and GLP-1 RAs).

PANRETIN

Products Affected

- Panretin

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Topical treatment of cutaneous lesions in patients with non-AIDS-related Kaposi sarcoma
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PEGASYS

Products Affected

- Pegasys

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Myeloproliferative neoplasm (essential thrombocythemia, polycythemia vera, symptomatic lower-risk myelofibrosis), systemic mastocytosis, adult T-cell leukemia/lymphoma, mycosis fungoides/sezary syndrome, primary cutaneous CD30+ T-cell lymphoproliferative disorders, hairy cell leukemia, Erdheim-Chester disease, initial treatment during pregnancy for chronic myeloid leukemia.
Exclusion Criteria	N/A
Required Medical Information	For chronic hepatitis C: Hepatitis C virus (HCV) confirmed by presence of hepatitis C virus HCV RNA in serum prior to starting treatment and the planned treatment regimen.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	HCV: 12-48wks. Criteria applied consistent w/current AASLD/IDSA guidance. HBV: 48wks. Other: Plan Yr
Other Criteria	N/A

PEMAZYRE

Products Affected

- Pemazyre

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PHENYL BUTYRATE

Products Affected

- Sodium Phenylbutyrate POWD 3GM/TSP
- Sodium Phenylbutyrate TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For urea cycle disorders (UCD): Diagnosis of UCD was confirmed by enzymatic, biochemical, or genetic testing.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PIQRAY

Products Affected

- Piqray 200mg Daily Dose
- Piqray 250mg Daily Dose
- Piqray 300mg Daily Dose

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, PIK3CA-mutated breast cancer in combination with fulvestrant.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

POMALYST

Products Affected

- Pomalyst

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Relapsed/refractory systemic light chain amyloidosis, primary central nervous system (CNS) lymphoma, POEMS (polyneuropathy, organomegaly, endocrinopathy, monoclonal protein, skin changes) syndrome.
Exclusion Criteria	N/A
Required Medical Information	For multiple myeloma, patient has previously received at least two prior therapies for multiple myeloma, including an immunomodulatory agent AND a proteasome inhibitor.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

POSACONAZOLE

Products Affected

- Posaconazole Dr

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The requested drug will be used orally. For prophylaxis of invasive Aspergillus and Candida infections: patient weighs greater than 40 kilograms.
Age Restrictions	Treatment of Invasive Aspergillosis: 13 years of age or older, Prophylaxis of Invasive Aspergillus and Candida Infections: 2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

PREGABALIN

Products Affected

- Pregabalin CAPS

- Pregabalin SOLN

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Cancer-related neuropathic pain, cancer treatment-related neuropathic pain
Exclusion Criteria	N/A
Required Medical Information	For the management of postherpetic neuralgia, the management of neuropathic pain associated with diabetic peripheral neuropathy, cancer-related neuropathic pain, and cancer treatment-related neuropathic pain: The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to gabapentin.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PRETOMANID AET

Products Affected

- Pretomanid

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For tuberculosis (TB), patient meets ALL of the following: 1) the requested drug is being used for treatment of pulmonary extensively drug resistant (XDR), treatment intolerant or nonresponsive multidrug resistant (MDR) tuberculosis, AND 2) the requested drug is used in combination with bedaquiline and linezolid.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PREVYMIS

Products Affected

- Prevymis TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For prophylaxis of cytomegalovirus (CMV) infection or disease in hematopoietic stem cell transplant (HSCT): 1) the patient is CMV-seropositive, AND 2) the patient is a recipient of an allogeneic HSCT. For prophylaxis of CMV disease in kidney transplant: 1) the patient is CMV-seronegative, AND 2) the patient is a high risk recipient of kidney transplant.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	7 months
Other Criteria	N/A

PROMACTA

Products Affected

- Promacta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>For chronic or persistent immune thrombocytopenia (ITP): 1) For new starts: a) Patient (pt) has had an inadequate response or is intolerant to a prior therapy such as corticosteroids or immunoglobulins AND b) Untransfused platelet (plt) count at any point prior to the initiation of the requested medication is less than 30,000/mcL OR 30,000-50,000/mcL with symptomatic bleeding or risk factor(s) for bleeding (e.g., undergoing a medical or dental procedure where blood loss is anticipated, comorbidities such as peptic ulcer disease and hypertension, anticoagulation therapy, profession or lifestyle that predisposes pt to trauma) AND c) For chronic ITP only: pt has had an inadequate response or intolerance to Doptelet (avatrombopag). 2) For continuation of therapy, plt count response to the requested drug: a) Current plt count is less than or equal to 200,000/mcL, OR b) Current plt count is greater than 200,000/mcL to less than or equal to 400,000/mcL and dosing will be adjusted to a plt count sufficient to avoid clinically important bleeding. For thrombocytopenia associated with chronic hepatitis C: 1) For new starts: the requested drug is used for initiation and maintenance of interferon-based therapy. 2) For continuation of therapy: pt is receiving interferon-based therapy. For severe aplastic anemia (AA): 1) For new starts: a) Pt will use the requested drug with standard immunosuppressive therapy for first line treatment OR b) the pt had an insufficient response to immunosuppressive therapy. 2) For continuation of therapy: 1) Current plt count is 50,000-200,000/mcL, OR 2) Current plt count is less than 50,000/mcL and pt has not received appropriately titrated therapy for at least 16 weeks, OR 3) Current plt count is less than 50,000/mcL and pt is transfusion-independent, OR 4) Current plt count is greater than 200,000/mcL to less than or equal to 400,000/mcL and dosing will be adjusted to achieve and maintain an appropriate target plt count.</p>
Age Restrictions	N/A

Prescriber Restrictions	N/A
Coverage Duration	HCV: 6mo, ITP/AA initial: 6mo, ITP reauth: Plan Year, AA reauth: APR-Plan Year, IPR-16 wks
Other Criteria	APR: adequate platelet response (greater than 50,000/mcL), IPR: inadequate platelet response (less than 50,000/mcL).

PULMOZYME

Products Affected

- Pulmozyme SOLN 2.5MG/2.5ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

QINLOCK

Products Affected

- Qinlock

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent/progressive or unresectable gastrointestinal stromal tumor (GIST)
Exclusion Criteria	N/A
Required Medical Information	For unresectable, recurrent/progressive, advanced, or metastatic gastrointestinal stromal tumor (GIST), the patient meets either of the following: 1) patient has received prior treatment with 3 or more kinase inhibitors, including imatinib OR 2) patient has experienced disease progression following treatment with avapritinib and dasatinib.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

QUETIAPINE XR

Products Affected

- Quetiapine Fumarate Er

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Maintenance monotherapy treatment in bipolar I disorder, monotherapy treatment of generalized anxiety disorder, monotherapy treatment of major depressive disorder
Exclusion Criteria	N/A
Required Medical Information	<p>For all indications: If the patient is 65 years of age or older AND is using two or more additional central nervous system (CNS) active medications (e.g., lorazepam, sertraline, clonazepam, escitalopram, alprazolam, zolpidem) with the requested drug, the prescriber determined that taking multiple central nervous system (CNS) active medications is medically necessary. [Note: Use of multiple central nervous system (CNS) active medications in older adults is associated with an increased risk of falls].</p> <p>For treatment of schizophrenia: The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following generic products: aripiprazole, asenapine, lurasidone, olanzapine, quetiapine immediate-release, risperidone, ziprasidone. For acute treatment of manic or mixed episodes associated with bipolar I disorder or maintenance treatment of bipolar I disorder: The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following generic products: aripiprazole, asenapine, olanzapine, quetiapine immediate-release, risperidone, ziprasidone. For acute treatment of depressive episodes associated with bipolar I disorder: The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following generic products: lurasidone, olanzapine, quetiapine immediate-release. For acute treatment of depressive episodes associated with bipolar II disorder: The patient experienced an inadequate treatment response or intolerance to generic quetiapine immediate-release. For adjunctive treatment of major depressive disorder (MDD): The patient experienced an inadequate treatment response, intolerance, or contraindication to one of the following generic products: aripiprazole, olanzapine, quetiapine immediate-release.</p>
Age Restrictions	N/A

Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

QUININE SULFATE

Products Affected

- Quinine Sulfate CAPS 324MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Babesiosis, uncomplicated Plasmodium vivax malaria.
Exclusion Criteria	N/A
Required Medical Information	For babesiosis: the requested drug is used in combination with clindamycin.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 month
Other Criteria	N/A

QULIPTA

Products Affected

- Qulipta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Preventive treatment of migraine, initial: 1) The patient experienced an inadequate treatment response with a 4-week trial of any one of the following: antiepileptic drugs (AEDs), beta-adrenergic blocking agents, antidepressants OR 2) The patient experienced an intolerance or has a contraindication that would prohibit a 4-week trial of any one of the following: antiepileptic drugs (AEDs), beta-adrenergic blocking agents, antidepressants. Preventive treatment of migraine, continuation: The patient received at least 3 months of treatment with the requested drug and the patient had a reduction in migraine days per month from baseline.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial: 3 months, Continuation: Plan Year
Other Criteria	N/A

REGRANEX

Products Affected

- Regranex

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	20 weeks
Other Criteria	N/A

REPATHA

Products Affected

- Repatha
- Repatha Pushtronex System
- Repatha Sureclick

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

RETEVMO

Products Affected

- Retevmo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent rearranged during transfection (RET)-rearrangement positive non-small cell lung cancer, Langerhans Cell Histiocytosis with a RET gene fusion, symptomatic or relapsed/refractory Erdheim-Chester Disease with a RET gene fusion, symptomatic or relapsed/refractory Rosai-Dorfman Disease with a RET gene fusion, RET-fusion positive recurrent or persistent thyroid carcinoma (papillary carcinoma, follicular carcinoma, and Hurthle cell carcinoma), RET-fusion positive anaplastic thyroid carcinoma.
Exclusion Criteria	N/A
Required Medical Information	For non-small cell lung cancer, patient must meet all of the following: 1) The disease is recurrent, advanced or metastatic, and 2) Tumor is RET fusion-positive or RET rearrangement-positive.
Age Restrictions	Medullary thyroid cancer and thyroid cancer: 12 years of age or older.
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

REVLIMID

Products Affected

- Lenalidomide

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Systemic light chain amyloidosis, classical Hodgkin lymphoma, myelodysplastic syndrome without the 5q deletion cytogenetic abnormality, myelofibrosis-associated anemia, POEMS (polyneuropathy, organomegaly, endocrinopathy, monoclonal protein, skin changes) syndrome, myeloproliferative neoplasms, Kaposi Sarcoma, Langerhans cell histiocytosis, peripheral T-Cell lymphomas not otherwise specified, angioimmunoblastic T-cell lymphoma (AITL), enteropathy-associated T-cell lymphoma, monomorphic epitheliotropic intestinal T-cell lymphoma, nodal peripheral T-cell lymphoma, adult T-cell leukemia/lymphoma, hepatosplenic T-cell lymphoma, primary central nervous system (CNS) lymphoma, chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL), acquired immunodeficiency syndrome (AIDS)-related B-cell lymphoma, monomorphic post-transplant lymphoproliferative disorder, diffuse large B-cell lymphoma, multicentric Castleman's disease, high-grade B-cell lymphomas, histologic transformation of indolent lymphoma to diffuse large B-cell lymphoma.
Exclusion Criteria	N/A
Required Medical Information	For myelodysplastic syndrome (MDS): patient has lower risk MDS with symptomatic anemia per the Revised International Prognostic Scoring System (IPSS-R), International Prognostic Scoring System (IPSS), or World Health organization (WHO) classification-based Prognostic Scoring System (WPSS).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

REZLIDHIA

Products Affected

- Rezlidhia

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

REZUROCK

Products Affected

- Rezurock

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	12 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

RINVOQ

Products Affected

- Rinvoq

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>For moderately to severely active rheumatoid arthritis (new starts only): patient has experienced an inadequate treatment response, intolerance or has a contraindication to at least one tumor necrosis factor (TNF) inhibitor (e.g., adalimumab-aacf, Enbrel [etanercept], Humira [adalimumab], Idacio [adalimumab-aacf]). For active psoriatic arthritis (new starts only): patient has experienced an inadequate treatment response, intolerance or has a contraindication to at least one TNF inhibitor (e.g., adalimumab-aacf, Enbrel [etanercept], Humira [adalimumab], Idacio [adalimumab-aacf]). For moderately to severely active ulcerative colitis (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one TNF inhibitor (e.g., adalimumab-aacf, Humira [adalimumab], Idacio [adalimumab-aacf]). For moderately to severely active Crohn's disease (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one TNF inhibitor (e.g., adalimumab-aacf, Humira [adalimumab], Idacio [adalimumab-aacf]). For atopic dermatitis (new starts only): 1) patient has refractory, moderate to severe disease, AND 2) patient has had an inadequate response to treatment with other systemic drug products, including biologics, or use of these therapies are inadvisable. For atopic dermatitis (continuation of therapy): the patient achieved or maintained positive clinical response. For active ankylosing spondylitis (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one TNF inhibitor (e.g., adalimumab-aacf, Enbrel [etanercept], Humira [adalimumab], Idacio [adalimumab-aacf]). For non-radiographic axial spondyloarthritis (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one TNF inhibitor.</p>
Age Restrictions	Atopic dermatitis: 12 years of age or older

Prescriber Restrictions	N/A
Coverage Duration	Atopic dermatitis (initial): 4 months, All others: Plan Year
Other Criteria	N/A

ROZLYTREK

Products Affected

- Rozlytrek

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent ROS1-positive non-small cell lung cancer (NSCLC), Non-metastatic neurotrophic tyrosine receptor kinase (NTRK) gene fusion-positive solid tumors, first-line treatment of NTRK gene fusion-positive solid tumors.
Exclusion Criteria	N/A
Required Medical Information	For all neurotrophic tyrosine receptor kinase (NTRK) gene fusion-positive solid tumors, the disease is without a known acquired resistance mutation. For ROS1-positive non-small cell lung cancer, the patient has recurrent, advanced, or metastatic disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

RUBRACA

Products Affected

- Rubraca

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Uterine leiomyosarcoma, pancreatic adenocarcinoma, advanced (stage II-IV) epithelial ovarian, fallopian tube, or primary peritoneal cancer
Exclusion Criteria	N/A
Required Medical Information	For metastatic castration-resistant prostate cancer with a deleterious breast cancer susceptibility gene (BRCA) mutation (germline and/or somatic): 1) patient has been treated with androgen receptor-directed therapy, AND 2) patient has been treated with a taxane-based chemotherapy or the patient is not fit for chemotherapy, AND 3) the requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy. For maintenance treatment of BRCA mutated epithelial ovarian, fallopian tube, primary peritoneal cancer: 1) the patient has advanced (stage II-IV) disease and is in complete or partial response to primary therapy, OR 2) the patient has recurrent disease and is in complete or partial response to platinum-based chemotherapy. For uterine leiomyosarcoma: 1) the requested drug is used as second-line therapy, AND 2) the patient has BRCA-altered disease. For pancreatic adenocarcinoma: 1) the patient has metastatic disease, AND 2) the patient has somatic or germline BRCA or PALB-2 mutations.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

RYBELSUS

Products Affected

- Rybelsus

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	The Prior Authorization only applies to patients whose claim is not submitted with an ICD-10 code indicating a diagnosis of type 2 diabetes mellitus OR to patients who do not have a history of an antidiabetic drug (EXCLUDING glucagon-like peptide receptor agonists [GLP-1 RAs] and combination glucose-dependent insulinotropic polypeptide [GIP] and GLP-1 RAs).

RYDAPT

Products Affected

- Rydapt

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Relapsed or refractory acute myeloid leukemia (AML), myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and FGFR1 or FLT3 rearrangements, post-induction therapy for AML, re-induction in residual disease for AML
Exclusion Criteria	N/A
Required Medical Information	For acute myeloid leukemia (AML): AML is FMS-like tyrosine kinase 3 (FLT3) mutation-positive. For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and Fibroblast growth factor receptor type 1 (FGFR1) or FLT3 rearrangements: the disease is in chronic or blast phase.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SANDOSTATIN LAR

Products Affected

- Sandostatin Lar Depot

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Tumor control of thymomas and thymic carcinomas, tumor control of neuroendocrine tumors (NETs) of the pancreas, gastrointestinal tract, lung, thymus, unresected primary gastrinoma, well-differentiated grade 3 NETs, pheochromocytoma/paraganglioma.
Exclusion Criteria	N/A
Required Medical Information	For acromegaly, initial: 1) Patient has a high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range, AND 2) Patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For acromegaly, continuation of therapy: Patient's IGF-1 level has decreased or normalized since initiation of therapy. The requested drug will be used for tumor control for any of the following: 1) Neuroendocrine tumor (NET) of the gastrointestinal tract or pancreas in patients with recurrent, locoregional advanced disease and/or distant metastatic disease, OR 2) NET of the thymus or lung in patients with locoregional unresectable disease and/or distant metastatic disease, OR 3) Unresected primary gastrinoma, OR 4) Well-differentiated grade 3 unresectable locally advanced or metastatic NET with favorable biology (e.g., relatively low Ki-67 [less than 55%] and positive somatostatin receptor [SSTR]-based positron emission tomography [PET] imaging), OR 5) Thymomas or thymic carcinomas when the following criteria are met: a) Locally advanced or metastatic disease OR b) Postoperatively following tumor resection, OR 6) Pheochromocytomas or paragangliomas when the following criteria is met: a) Locally unresectable OR b) Distant metastatic disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	N/A
-----------------------	-----

SAPROPTERIN

Products Affected

- Javygtor

- Sapropterin Dihydrochloride

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For phenylketonuria (PKU): For patients who have not yet received a therapeutic trial of the requested drug, the patient's pretreatment (including before dietary management) phenylalanine level is greater than 6 mg/dL (360 micromol/L). For patients who completed a therapeutic trial of the requested drug, the patient must have experienced improvement (e.g., reduction in blood phenylalanine levels, improvement in neuropsychiatric symptoms).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Initial: 2 months, All others: Plan Year
Other Criteria	N/A

SCSEMBLIX

Products Affected

- Scemblix

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For chronic myeloid leukemia (CML) in the chronic phase: 1) the diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene AND 2) the patient meets either of the following: A) the patient has previously been treated with 2 or more tyrosine kinase inhibitors (TKIs) AND at least one of those was imatinib or dasatinib, OR B) the patient is positive for the T315I mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SIGNIFOR

Products Affected

- Signifor

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	Plan Year
Other Criteria	N/A

SILDENAFIL

Products Affected

- Sildenafil Citrate TABS 20MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): PAH was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, AND 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) If the request is for an adult, pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SILDENAFIL INJ

Products Affected

- Sildenafil INJ

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): PAH was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, AND 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) If the request is for an adult, pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 month
Other Criteria	Patient was previously receiving oral Revatio or sildenafil but is now temporarily unable to take oral medications.

SIRTURO

Products Affected

- Sirturo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist.
Coverage Duration	Plan Year
Other Criteria	N/A

SKYRIZI

Products Affected

- Skyrizi INJ 150MG/ML, 180MG/1.2ML, 360MG/2.4ML, 600MG/10ML
- Skyrizi Pen

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) patient meets any of the following: a) patient has experienced an inadequate treatment response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, c) patient has severe psoriasis that warrants a biologic as first-line therapy (i.e., at least 10% of the body surface area or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SOMATULINE DEPOT

Products Affected

- Somatuline Depot

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Tumor control of neuroendocrine tumors (NETs) of the lung, thymus or unresected primary gastrinoma, well-differentiated grade 3 neuroendocrine tumors not of gastroenteropancreatic origin, pheochromocytoma/paraganglioma.
Exclusion Criteria	N/A
Required Medical Information	For acromegaly, initial: 1) Patient has a high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range, AND 2) Patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For acromegaly, continuation of therapy: Patient's IGF-1 level has decreased or normalized since initiation of therapy. For tumor control of neuroendocrine tumors (NETs) of the thymus or lung: Patient has locoregional unresectable, recurrent, and/or distant metastatic disease. For tumor control of well-differentiated grade 3 unresectable locally advanced or metastatic NETs (not of gastroenteropancreatic origin): Patient has favorable biology (e.g., relatively low Ki-67 [less than 55%] and positive somatostatin receptor [SSTR]-based positron emission tomography [PET] imaging). For tumor control of pheochromocytomas or paragangliomas: Patient has locally unresectable or distant metastatic disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SOMAVERT

Products Affected

- Somavert

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For acromegaly, initial: 1) Patient has a high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range, AND 2) Patient had an inadequate or partial response to surgery or radiotherapy OR there is a clinical reason for why the patient has not had surgery or radiotherapy. For acromegaly, continuation of therapy: Patient's IGF-1 level has decreased or normalized since initiation of therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SPRITAM

Products Affected

- Spritam

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Partial-Onset Seizures (i.e., focal-onset seizures): 4 years of age or older. Myoclonic Seizures: 12 years of age or older. Primary Generalized Tonic-Clonic Seizures: 6 years of age or older.
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SPRYCEL

Products Affected

- Sprycel

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Gastrointestinal stromal tumor (GIST), metastatic chondrosarcoma, recurrent chordoma, T-cell acute lymphoblastic leukemia (ALL), and Philadelphia (Ph)-like B-ALL, myeloid and/or lymphoid neoplasms with eosinophilia and ABL1 rearrangement in the chronic phase or blast phase
Exclusion Criteria	N/A
Required Medical Information	For chronic myeloid leukemia (CML), including patients who have received a hematopoietic stem cell transplant: 1) Diagnosis was confirmed by detection of the Philadelphia (Ph) chromosome or BCR-ABL gene, and 2) If patient experienced resistance to an alternative tyrosine kinase inhibitor, patient is negative for all of the following mutations: T315I/A, F317L/V/I/C, and V299L. For acute lymphoblastic leukemia (ALL), the patient has a diagnosis of one of the following: 1) Philadelphia chromosome positive ALL, including patients who have received a hematopoietic stem cell transplant: diagnosis that has been confirmed by detection of the Ph chromosome or BCR-ABL gene, and if patient experienced resistance to an alternative tyrosine kinase inhibitor, patient is negative for all of the following mutations: T315I/A, F317L/V/I/C, and V299L, OR 2) Ph-like B-ALL with ABL-class kinase fusion, OR 3) relapsed or refractory T-cell ALL with ABL-class kinase fusion. For GIST, 1) the patient meets all of the following: A) the disease is unresectable, recurrent/progressive, or metastatic, B) the patient has received prior therapy with imatinib or avapritinib AND C) patients is positive for PDGFRA exon 18 mutations, OR 2) the requested drug is being used for palliation of symptoms.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

STELARA

Products Affected

- Stelara

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderate to severe plaque psoriasis (new starts): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis AND 2) patient meets any of the following: a) the patient has experienced an inadequate treatment response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, c) patient has severe psoriasis that warrants a biologic as first-line therapy (i.e. at least 10% of the BSA or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

STIVARGA

Products Affected

- Stivarga

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Progressive gastrointestinal stromal tumors (GIST), osteosarcoma, glioblastoma, angiosarcoma, retroperitoneal/intra-abdominal soft tissue sarcoma, rhabdomyosarcoma, soft tissue sarcomas of the extremities, body wall, head and neck.
Exclusion Criteria	N/A
Required Medical Information	For gastrointestinal stromal tumors: The disease is progressive, locally advanced, unresectable, or metastatic. For colorectal cancer: The disease is advanced or metastatic.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SUTENT

Products Affected

- Sunitinib Malate

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Thyroid carcinoma (follicular, medullary, papillary, and Hurthle cell), soft tissue sarcoma (angiosarcoma, solitary fibrous tumor, and alveolar soft part sarcoma subtypes), recurrent chordoma, thymic carcinoma, lymphoid, myeloid, or mixed lineage neoplasms with eosinophilia, pheochromocytoma, paraganglioma, gastrointestinal stromal tumor (GIST) (unresectable, recurrent/progressive, or metastatic disease after progression on approved therapies, unresectable succinate dehydrogenase (SDH)-deficient GISTs and use for palliation of symptoms if previously tolerated and effective).
Exclusion Criteria	N/A
Required Medical Information	For renal cell carcinoma (RCC): the patient meets either of the following: 1) the disease is relapsed, advanced, or stage IV OR 2) the requested drug is being used as adjuvant treatment for patients that are at high risk of recurrent RCC following nephrectomy. For gastrointestinal stromal tumor (GIST): the patient meets one of the following: 1) the requested drug will be used after disease progression on or intolerance to imatinib, 2) the disease is unresectable, recurrent/progressive, or metastatic AND the patient has failed on an FDA-approved therapy (e.g., imatinib, sunitinib, regorafenib, ripretinib), 3) the requested drug will be used for unresectable succinate dehydrogenase (SDH)-deficient GIST, OR 4) the requested drug will be used for the palliation of symptoms if previously tolerated and effective. For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia: 1) the disease has a FLT3 rearrangement AND 2) the disease is in chronic or blast phase.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SYMLIN

Products Affected

- Symlinpen 120
- Symlinpen 60

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

SYMPAZAN

Products Affected

- Sympazan

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Seizures associated with Dravet syndrome
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Seizures associated with Lennox-Gastaut syndrome (LGS): 2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TABRECTA

Products Affected

- Tabrecta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent non-small cell lung cancer (NSCLC).
Exclusion Criteria	N/A
Required Medical Information	For recurrent, advanced, or metastatic NSCLC: Tumor is positive for mesenchymal-epithelial transition (MET) exon 14 skipping mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TADALAFIL (PAH)

Products Affected

- Alyq
- Tadalafil TABS 20MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group 1): PAH was confirmed by right heart catheterization. For PAH new starts only: 1) Pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, AND 2) Pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) Pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TAFINLAR

Products Affected

- Tafinlar

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Thyroid carcinoma (papillary carcinoma, follicular carcinoma, and Hurthle cell carcinoma), central nervous system (CNS) cancer (i.e., oligodendroglioma, astrocytoma, glioblastoma), gallbladder cancer, extrahepatic cholangiocarcinoma, intrahepatic cholangiocarcinoma, Langerhans cell histiocytosis, Erdheim-Chester disease, ovarian cancer, fallopian tube cancer, and primary peritoneal cancer.
Exclusion Criteria	N/A
Required Medical Information	For central nervous system (CNS) cancer (i.e., glioma, oligodendroglioma, astrocytoma, glioblastoma): 1) The tumor is positive for a BRAF V600E mutation AND 2) The requested drug will be used in combination with trametinib. For melanoma: 1) The tumor is positive for a BRAF V600 activating mutation (e.g., V600E or V600K), AND 2) The requested drug will be used as a single agent or in combination with trametinib, AND 3) The requested drug will be used for either of the following: a) unresectable, limited resectable, or metastatic disease, b) adjuvant systemic therapy. For non-small cell lung cancer: 1) The tumor is positive for a BRAF V600E mutation, AND 2) The requested drug will be used as a single agent or in combination with trametinib. For papillary, follicular, and Hurthle cell thyroid carcinoma: 1) The tumor is BRAF-positive, AND 2) The disease is not amenable to radioactive iodine (RAI) therapy. For Langerhans Cell Histiocytosis and Erdheim-Chester Disease: The disease is positive for a BRAF V600E mutation. For gallbladder cancer, extrahepatic cholangiocarcinoma, and intrahepatic cholangiocarcinoma: 1) The disease is positive for a BRAF V600E mutation, AND 2) The disease is unresectable or metastatic, AND 3) The requested drug will be used in combination with trametinib. For solid tumors: 1) The tumor is positive for a BRAF V600E mutation, AND 2) The requested drug will be used in combination with trametinib. For ovarian cancer, fallopian tube cancer, and primary peritoneal cancer: 1) The disease is positive for BRAF V600E mutation, AND 2) The disease is persistent or recurrent, AND 3) The requested drug will be used in combination with trametinib.
Age Restrictions	N/A

Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TAGRISSEO

Products Affected

- Tagrisso

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Sensitizing epidermal growth factor receptor (EGFR) mutation-positive recurrent non-small cell lung cancer (NSCLC), brain metastases from sensitizing EGFR mutation-positive NSCLC, leptomeningeal metastases from EGFR mutation-positive NSCLC.
Exclusion Criteria	N/A
Required Medical Information	For NSCLC, the requested drug is used in any of the following settings: 1) The patient meets both of the following: a) patient has metastatic, advanced, or recurrent NSCLC (including brain and/or leptomeningeal metastases from NSCLC) and b) patient has a sensitizing EGFR mutation OR 2) The patient meets both of the following: a) request is for adjuvant treatment of NSCLC following tumor resection and b) patient has EGFR mutation-positive disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TALTZ

Products Affected

- Taltz

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis AND 2) the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to one of the following products: adalimumab-aacf, Enbrel (etanercept), Humira (adalimumab), Idacio (adalimumab-aacf), Otezla (apremilast), Skyrizi (risankizumab-rzaa), Stelara (ustekinumab). For active ankylosing spondylitis (new starts only): the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to one of the following products: adalimumab-aacf, Enbrel (etanercept), Humira (adalimumab), Idacio (adalimumab-aacf), Rinvoq (upadacitinib), Xeljanz (tofacitinib)/Xeljanz XR (tofacitinib extended-release). For active psoriatic arthritis (PsA) (new starts only): the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to one of the following products: adalimumab-aacf, Enbrel (etanercept), Humira (adalimumab), Idacio (adalimumab-aacf), Otezla (apremilast), Rinvoq (upadacitinib), Skyrizi (risankizumab-rzaa), Stelara (ustekinumab), Xeljanz (tofacitinib)/Xeljanz XR (tofacitinib extended-release). For active non-radiographic axial spondyloarthritis (new starts only): patient meets any of the following: 1) patient has experienced an inadequate treatment response to a non-steroidal anti-inflammatory drug (NSAID) OR 2) patient has experienced an intolerance or has a contraindication to NSAIDs.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	N/A
-----------------------	-----

TALZENNA

Products Affected

- Talzenna

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent germline breast cancer susceptibility gene (BRCA)-mutated breast cancer
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TARGRETIN TOPICAL

Products Affected

- Bexarotene GEL

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Stage 2 or higher mycosis fungoides (MF)/Sezary syndrome (SS), chronic or smoldering adult T-cell leukemia/lymphoma (ATLL), primary cutaneous marginal zone lymphoma, primary cutaneous follicle center lymphoma
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TASIGNA

Products Affected

- Tassigna

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL), gastrointestinal stromal tumor (GIST), myeloid and/or lymphoid neoplasms with eosinophilia and ABL1 rearrangement in the chronic phase or blast phase, pigmented villonodular synovitis/tenosynovial giant cell tumor
Exclusion Criteria	N/A
Required Medical Information	For chronic myeloid leukemia (CML), including patients newly diagnosed with CML and patients who have received a hematopoietic stem cell transplant, 1) Diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene, 2) patient has experienced resistance or intolerance to imatinib or dasatinib, AND 3) If patient experienced resistance to an alternative tyrosine kinase inhibitor for CML, patient is negative for T315I, Y253H, E255K/V, and F359V/C/I mutations. For acute lymphoblastic leukemia (ALL), including patients who have received a hematopoietic stem cell transplant: 1) Diagnosis was confirmed by detection of the Philadelphia chromosome or BCR-ABL gene, AND 2) if the patient has experienced resistance to an alternative tyrosine kinase inhibitor for ALL, patient is negative for T315I, Y253H, E255K/V, F359V/C/I and G250E. For gastrointestinal stromal tumor (GIST), the patients meets either of the following: 1) the disease is unresectable, recurrent/progressive, or metastatic AND the disease has progressed on at least 2 approved therapies (e.g. imatinib, sunitinib, dasatinib, regorafenib, ripretinib) OR 2) the requested drug is being prescribed for palliation of symptoms.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TAZAROTENE

Products Affected

- Tazarotene CREA
- Tazarotene GEL
- Tazorac CREA 0.05%

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For plaque psoriasis, the patient meets the following criteria: 1) the patient has less than or equal to 20 percent of affected body surface area (BSA), AND 2) the patient experienced an inadequate treatment response or intolerance to at least one topical corticosteroid OR has a contraindication that would prohibit a trial of topical corticosteroids.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TAZVERIK

Products Affected

- Tazverik

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	Epithelioid sarcoma: 16 years of age or older, Follicular lymphoma: 18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TECFIDERA

Products Affected

- Tecfidera
- Tecfidera Starter Pack

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TECVAYLI

Products Affected

- Tecvayli

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TEMAZEPAM 30MG - 65

Products Affected

- Temazepam CAPS 30MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For short-term treatment of insomnia: 1) The prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) AND 2) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to doxepin (3 mg or 6 mg).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	This Prior Authorization only applies to patients 65 years of age or older.

TEPMETKO

Products Affected

- Tepmetko

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent non-small cell lung cancer (NSCLC).
Exclusion Criteria	N/A
Required Medical Information	For recurrent, advanced, or metastatic NSCLC: Tumor is positive for mesenchymal-epithelial transition (MET) exon 14 skipping mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TESTOSTERONE ENANTHATE INJ

Products Affected

- Testosterone Enanthate INJ

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Gender Dysphoria
Exclusion Criteria	N/A
Required Medical Information	For primary hypogonadism or hypogonadotropic hypogonadism, initial therapy: The patient has at least two confirmed low morning serum total testosterone concentrations based on the reference laboratory range or current practice guidelines [Note: Safety and efficacy of testosterone products in patients with "age-related hypogonadism" (also referred to as "late-onset hypogonadism") have not been established.]. For primary hypogonadism or hypogonadotropic hypogonadism, continuation of therapy: The patient had a confirmed low morning serum total testosterone concentration based on the reference laboratory range or current practice guidelines before starting testosterone therapy [Note: Safety and efficacy of testosterone products in patients with "age-related hypogonadism" (also referred to as "late-onset hypogonadism") have not been established.]. For gender dysphoria: The patient is able to make an informed decision to engage in hormone therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TETRABENAZINE

Products Affected

- Tetrabenazine

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Tic disorders, tardive dyskinesia, hemiballismus, chorea not associated with Huntington's disease.
Exclusion Criteria	N/A
Required Medical Information	For treatment of tardive dyskinesia and treatment of chorea associated with Huntington's disease: The patient has experienced an inadequate treatment response or intolerable adverse event to deutetrabenazine.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

THALOMID

Products Affected

- Thalomid

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Myelofibrosis-associated anemia, AIDS-related aphthous stomatitis, Kaposi sarcoma, chronic graft-versus-host disease, Crohn's disease, multicentric Castleman's disease, Rosai-Dorfman disease, Langerhans cell histiocytosis
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TIBSOVO

Products Affected

- Tibsovo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Conventional (grades 1-3) or dedifferentiated chondrosarcoma. Newly-diagnosed acute myeloid leukemia (AML) if 60-74 years of age and without comorbidities.
Exclusion Criteria	N/A
Required Medical Information	Patient has disease with a susceptible isocitrate dehydrogenase-1 (IDH1) mutation. For acute myeloid leukemia (AML): 1) patient has newly-diagnosed AML and meets one of the following: a) 75 years of age or older, b) patient has comorbidities that preclude use of intensive induction chemotherapy, or c) patient is 60 years of age or older and declines intensive induction chemotherapy, OR 2) patient is 60 years of age or older and the requested drug will be used as post-induction therapy following response to induction therapy with the requested drug, OR 3) patient has relapsed or refractory AML. For locally advanced, unresectable, or metastatic cholangiocarcinoma: the requested drug will be used as subsequent treatment for progression on or after systemic treatment.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TOBRAMYCIN

Products Affected

- Tobramycin NEBU 300MG/5ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Non-cystic fibrosis bronchiectasis
Exclusion Criteria	N/A
Required Medical Information	For cystic fibrosis and non-cystic fibrosis bronchiectasis: 1) Pseudomonas aeruginosa is present in the patient's airway cultures, OR 2) The patient has a history of Pseudomonas aeruginosa infection or colonization in the airways.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

TOPICAL DOXEPIN

Products Affected

- Doxepin Hydrochloride CREA

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to a topical corticosteroid or a topical calcineurin inhibitor.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 month
Other Criteria	N/A

TOPICAL LIDOCAINE

Products Affected

- Lidocaine OINT 5%

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	1) The requested drug is being used for topical anesthesia, AND 2) If the requested drug will be used as part of a compounded product, then all the active ingredients in the compounded product are Food and Drug Administration (FDA) approved for topical use.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	3 months
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

TOPICAL TRETINOIN

Products Affected

- Tretinoin CREA
- Tretinoin GEL
- Tretinoin Microsphere GEL 0.04%, 0.1%
- Tretinoin Microsphere Pump

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TOREMIFENE

Products Affected

- Toremifene Citrate

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Congenital/acquired QT prolongation (long QT syndrome), uncorrected hypokalemia, or uncorrected hypomagnesemia.
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TRIENTINE

Products Affected

- Trientine Hydrochloride

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TRIKAFTA

Products Affected

- Trikafta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For cystic fibrosis: The requested medication will not be used in combination with other medications containing ivacaftor.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TRULICITY

Products Affected

- Trulicity

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	The Prior Authorization only applies to patients whose claim is not submitted with an ICD-10 code indicating a diagnosis of type 2 diabetes mellitus OR to patients who do not have a history of an antidiabetic drug (EXCLUDING glucagon-like peptide receptor agonists [GLP-1 RAs] and combination glucose-dependent insulinotropic polypeptide [GIP] and GLP-1 RAs).

TRUQAP

Products Affected

- Truqap

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TRUSELTIQ

Products Affected

- Truseltiq

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TRUXIMA

Products Affected

- Truxima

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	<p>Non-Hodgkin's lymphoma subtypes [small lymphocytic lymphoma (SLL), mantle cell lymphoma, marginal zone lymphomas (nodal, splenic, extranodal marginal zone lymphoma), Burkitt lymphoma, primary cutaneous B-cell lymphoma, high-grade B-cell lymphoma, histological transformation from indolent lymphomas to diffuse large B-cell lymphoma, histological transformation chronic lymphocytic leukemia (CLL)/SLL to diffuse large B-cell lymphoma, Castleman's disease, human immunodeficiency virus (HIV)-related B-cell lymphoma, hairy cell leukemia, post-transplant lymphoproliferative disorder (PTLD), B-cell lymphoblastic lymphoma], refractory immune or idiopathic thrombocytopenic purpura (ITP), autoimmune hemolytic anemia, Waldenstrom's macroglobulinemia/lymphoplasmacytic lymphoma, chronic graft-versus-host disease (GVHD), Sjogren syndrome, thrombotic thrombocytopenic purpura, refractory myasthenia gravis, Hodgkin's lymphoma (nodular lymphocyte-predominant), primary central nervous system (CNS) lymphoma, leptomeningeal metastases from lymphomas, acute lymphoblastic leukemia, prevention of Epstein-Barr virus (EBV)-related PTL, multiple sclerosis, immune checkpoint inhibitor-related toxicities, pemphigus vulgaris, pediatric aggressive mature B-cell lymphomas, Rosai-Dorfman disease, and pediatric mature B-cell acute leukemia.</p>
Exclusion Criteria	N/A
Required Medical Information	<p>For moderately to severely active rheumatoid arthritis (new starts only): 1) patient meets ANY of the following: a) requested drug will be used in combination with methotrexate (MTX) OR b) patient has intolerance or contraindication to MTX, AND 2) patient meets ANY of the following: a) inadequate response, intolerance, or contraindication to MTX OR b) inadequate response or intolerance to a prior biologic disease-modifying antirheumatic drug (DMARD) or a targeted synthetic DMARD. Hematologic malignancies must be CD20-positive. For multiple sclerosis: 1) patient has a diagnosis of relapsing remitting multiple sclerosis, AND 2) patient has had an inadequate response to two or more disease-modifying drugs indicated for multiple sclerosis despite adequate duration of treatment.</p>

Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Immune checkpoint inhibitor-related toxicities: 3 months, All other: Plan Year
Other Criteria	N/A

TUKYSA

Products Affected

- Tukysa

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent human epidermal growth factor receptor 2 (HER2)-positive breast cancer
Exclusion Criteria	N/A
Required Medical Information	For colorectal cancer (including appendiceal adenocarcinoma): 1) the patient has advanced, unresectable, or metastatic disease AND 2) the patient has human epidermal growth factor receptor 2 (HER2)-positive disease AND 3) the patient has RAS wild-type disease AND 4) the requested drug will be used in combination with trastuzumab and 5) the patient has not previously been treated with a HER2 inhibitor.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

TURALIO

Products Affected

- Turalio

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Langerhans cell histiocytosis, Erdheim-Chester disease, Rosai-Dorfman disease
Exclusion Criteria	N/A
Required Medical Information	For Langerhans cell histiocytosis: 1) disease has colony stimulating factor 1 receptor (CSF1R) mutation. For Erdheim-Chester disease and Rosai-Dorfman disease: 1) disease has CSF1R mutation AND patient has any of the following: a) symptomatic disease OR b) relapsed/refractory disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

UBRELVY

Products Affected

- Ubrelyvy

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For acute treatment of migraine: The patient has experienced an inadequate treatment response, intolerance, or the patient has a contraindication to at least one triptan 5-HT ₁ receptor agonist.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VALCHLOR

Products Affected

- Valchlor

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Chronic or smoldering adult T-cell leukemia/lymphoma (ATLL), Stage 2 or higher mycosis fungoides (MF)/Sezary syndrome (SS), primary cutaneous marginal zone lymphoma, primary cutaneous follicle center lymphoma, CD30-positive lymphomatoid papulosis (LyP), unifocal Langerhans cell histiocytosis (LCH) with isolated skin disease
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VALTOCO

Products Affected

- Valtoco 10 Mg Dose
- Valtoco 15 Mg Dose
- Valtoco 20 Mg Dose
- Valtoco 5 Mg Dose

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	6 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VANFLYTA

Products Affected

- Vanflyta

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VENCLEXTA

Products Affected

- Venclexta

- Venclexta Starting Pack

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Mantle cell lymphoma, blastic plasmacytoid dendritic cell neoplasm (BPDCN), multiple myeloma, relapsed or refractory acute myeloid leukemia (AML), Waldenstrom macroglobulinemia/lymphoplasmacytic lymphoma, relapsed or refractory systemic light chain amyloidosis with translocation t(11:14), myelodysplastic syndrome
Exclusion Criteria	N/A
Required Medical Information	For acute myeloid leukemia (AML): 1) patient is 60 years of age or older, OR 2) patient is less than 60 years of age with unfavorable risk genetics and TP53-mutation, OR 3) patient has comorbidities that preclude use of intensive induction chemotherapy, OR 4) patient has relapsed or refractory disease. For blastic plasmacytoid dendritic cell neoplasm (BPDCN): 1) patient has systemic disease being treated with palliative intent, OR 2) patient has relapsed or refractory disease. For multiple myeloma: 1) the disease is relapsed or progressive, AND 2) the requested drug will be used in combination with dexamethasone, AND 3) patient has t(11:14) translocation. For Waldenstrom macroglobulinemia/lymphoplasmacytic lymphoma: 1) patient has previously treated disease that did not respond to primary therapy, OR 2) patient has progressive or relapsed disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VENTAVIS

Products Affected

- Ventavis

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For pulmonary arterial hypertension (World Health Organization [WHO] Group 1): PAH was confirmed by right heart catheterization. For new starts only: 1) pretreatment mean pulmonary arterial pressure is greater than 20 mmHg, AND 2) pretreatment pulmonary capillary wedge pressure is less than or equal to 15 mmHg, AND 3) pretreatment pulmonary vascular resistance is greater than or equal to 3 Wood units.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

VERQUVO

Products Affected

- Verquvo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For symptomatic chronic heart failure: the patient has a left ventricular ejection fraction (LVEF) less than 45 percent. For initial therapy, the patient meets ANY of the following: 1) hospitalization for heart failure within the past 6 months OR 2) use of outpatient intravenous diuretics for heart failure within the past 3 months.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VERSACLOZ

Products Affected

- Versacloz

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the treatment of a severely ill patient with schizophrenia who failed to respond adequately to standard antipsychotic treatment (i.e., treatment-resistant schizophrenia): 1) the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to one of the following generic products: aripiprazole, asenapine, lurasidone, olanzapine, quetiapine, risperidone, ziprasidone, AND 2) the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to one of the following brand products: Caplyta, Rexulti, Secuado, Vraylar.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VERZENIO

Products Affected

- Verzenio

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer in combination with fulvestrant or an aromatase inhibitor, or as a single agent if progression on prior endocrine therapy and prior chemotherapy in the metastatic setting.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VIGABATRIN

Products Affected

- Vigabatrin
- Vigadrone
- Vigpoder

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For complex partial seizures (i.e., focal impaired awareness seizures): patient has experienced an inadequate treatment response to at least two antiepileptic drugs for complex partial seizures (i.e., focal impaired awareness seizures).
Age Restrictions	Infantile Spasms: 1 month to 2 years of age. Complex partial seizures (i.e., focal impaired awareness seizures): 2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VITRAKVI

Products Affected

- Vitrakvi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Non-metastatic neurotrophic tyrosine receptor kinase (NTRK) gene fusion-positive solid tumors, first-line treatment of NTRK gene fusion-positive solid tumors.
Exclusion Criteria	N/A
Required Medical Information	For all neurotrophic tyrosine receptor kinase (NTRK) gene fusion-positive solid tumors, the disease is without a known acquired resistance mutation.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VIZIMPRO

Products Affected

- Vizimpro

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent non-small cell lung cancer (NSCLC).
Exclusion Criteria	N/A
Required Medical Information	For non-small cell lung cancer (NSCLC): 1) the disease is recurrent, advanced or metastatic, and 2) the patient has sensitizing EGFR mutation-positive disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VONJO

Products Affected

- Vonjo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VORICONAZOLE

Products Affected

- Voriconazole INJ
- Voriconazole SUSR

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	The patient will use the requested drug orally or intravenously.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

VOSEVI

Products Affected

- Vosevi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	Decompensated cirrhosis/moderate or severe hepatic impairment (Child Turcotte Pugh class B or C)
Required Medical Information	For hepatitis C: Infection confirmed by presence of HCV RNA in the serum prior to starting treatment. Planned treatment regimen, genotype, prior treatment history, presence or absence of cirrhosis (compensated or decompensated [Child Turcotte Pugh class B or C]), presence or absence of HIV coinfection, presence or absence of resistance-associated substitutions where applicable, transplantation status if applicable. Coverage conditions and specific durations of approval will be based on current American Association for the Study of Liver Diseases and Infectious Diseases Society of America (AASLD-IDSA) treatment guidelines.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Criteria will be applied consistent with current AASLD-IDSA guidance.
Other Criteria	N/A

VOTRIENT

Products Affected

- Pazopanib Hydrochloride

- Votrient

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Thyroid carcinoma (follicular, papillary, Hurthle cell, or medullary), uterine sarcoma, chondrosarcoma, gastrointestinal stromal tumor
Exclusion Criteria	N/A
Required Medical Information	For renal cell carcinoma: 1) The disease is advanced, relapsed, or stage IV, OR 2) the requested drug will be used for von Hippel-Lindau (VHL)-associated renal cell carcinoma. For gastrointestinal stromal tumor (GIST): the patients meets one of the following: 1) the disease is unresectable, recurrent/progressive, or metastatic AND the patient has failed an FDA-approved therapy (e.g., imatinib, sunitinib, regorafenib, ripretinib), 2) the requested drug will be used for unresectable succinate dehydrogenase (SDH)-deficient GIST, OR 3) the requested drug will be used for the palliation of symptoms if previously tolerated and effective. For soft tissue sarcoma (STS): The patient does not have an adipocytic soft tissue sarcoma. For uterine sarcoma: The disease is recurrent or metastatic.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

VUMERITY

Products Affected

- Vumerity

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

WELIREG

Products Affected

- Welireg

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XALKORI

Products Affected

- Xalkori

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent non-small cell lung cancer (NSCLC), NSCLC with high-level MET amplification or MET exon 14 skipping mutation, symptomatic or relapsed/refractory anaplastic lymphoma kinase (ALK)-fusion positive Erdheim-Chester Disease, symptomatic or relapsed/refractory (ALK)-fusion positive Rosai-Dorfman Disease, (ALK)-fusion positive Langerhans Cell Histiocytosis.
Exclusion Criteria	N/A
Required Medical Information	For NSCLC, the requested drug is used in any of the following settings: 1) the patient has recurrent, advanced or metastatic ALK-positive NSCLC, OR 2) the patient has recurrent, advanced or metastatic ROS-1 positive NSCLC, OR 3) the patient has NSCLC with high-level MET amplification or MET exon 14 skipping mutation. For IMT, the disease is ALK-positive. For ALCL, the disease is relapsed or refractory and ALK-positive.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XELJANZ

Products Affected

- Xeljanz

- Xeljanz Xr

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For moderately to severely active rheumatoid arthritis (new starts only): patient has experienced an inadequate treatment response, intolerance or has a contraindication to at least one tumor necrosis factor (TNF) inhibitor (e.g., adalimumab-aacf, Enbrel [etanercept], Humira [adalimumab], Idacio [adalimumab-aacf]). For active psoriatic arthritis (new starts only): 1) patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one TNF inhibitor (e.g., adalimumab-aacf, Enbrel [etanercept], Humira [adalimumab], Idacio [adalimumab-aacf]) AND 2) the requested drug is used in combination with a nonbiologic DMARD. For active ankylosing spondylitis (new starts only): Patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one tumor necrosis factor (TNF) inhibitor (e.g., adalimumab-aacf, Enbrel [etanercept], Humira [adalimumab], Idacio [adalimumab-aacf]). For moderately to severely active ulcerative colitis (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one tumor necrosis factor (TNF) inhibitor (e.g., adalimumab-aacf, Humira [adalimumab], Idacio [adalimumab-aacf]). For active polyarticular course juvenile idiopathic arthritis (new starts only): patient has experienced an inadequate treatment response, intolerance, or has a contraindication to at least one tumor necrosis factor (TNF) inhibitor (e.g., adalimumab-aacf, Enbrel [etanercept], Humira [adalimumab], Idacio [adalimumab-aacf]).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year

Other Criteria	N/A
-----------------------	-----

XERMELO

Products Affected

- Xermelo

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XGEVA

Products Affected

- Xgeva

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For hypercalcemia of malignancy: condition is refractory to intravenous (IV) bisphosphonate therapy or there is a clinical reason to avoid IV bisphosphonate therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

XHANCE MAPD

Products Affected

- Xhance

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Patient has experienced an inadequate treatment response to generic fluticasone nasal spray.
Age Restrictions	18 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XIFAXAN

Products Affected

- Xifaxan TABS 550MG

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For irritable bowel syndrome with diarrhea (IBS-D): 1) The patient has not previously received treatment with the requested drug OR 2) The patient has previously received treatment with the requested drug AND a) the patient is experiencing a recurrence of symptoms AND b) the patient has not already received an initial 14-day course of treatment and two additional 14-day courses of treatment with the requested drug.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Reduction in risk of overt HE recurrence: 6 Months, IBS-D: 14 Days
Other Criteria	N/A

XOLAIR

Products Affected

- Xolair INJ 150MG, 150MG/ML, 75MG/0.5ML

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	<p>For moderate to severe persistent asthma, initial therapy: 1) Patient has a positive skin test (or blood test) to at least one perennial aeroallergen, 2) Patient has baseline IgE level greater than or equal to 30 IU/mL, AND 3) Patient has inadequate asthma control despite current treatment with both of the following medications: a) Medium-to-high-dose inhaled corticosteroid, AND b) Additional controller (i.e., long acting beta2-agonist, long-acting muscarinic antagonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies. For moderate to severe persistent asthma, continuation of therapy: Asthma control has improved on treatment with the requested drug, as demonstrated by a reduction in the frequency and/or severity of symptoms and exacerbations or a reduction in the daily maintenance oral corticosteroid dose. For chronic spontaneous urticaria (CSU), initial therapy: 1) Patient has been evaluated for other causes of urticaria, including bradykinin-related angioedema and IL-1-associated urticarial syndromes (e.g., auto-inflammatory disorders, urticarial vasculitis), 2) Patient has experienced a spontaneous onset of wheals, angioedema, or both, for at least 6 weeks, AND 3) Patient remains symptomatic despite H1 antihistamine treatment. For CSU, continuation of therapy: Patient has experienced a benefit (e.g., improved symptoms) since initiation of therapy. For chronic rhinosinusitis with nasal polyps (CRSwNP): 1) The requested drug is used as add-on maintenance treatment, AND 2) Patient has experienced inadequate treatment response to Xhance (fluticasone).</p>
Age Restrictions	CSU: 12 years of age or older. Asthma: 6 years of age or older. CRSwNP: 18 years of age or older
Prescriber Restrictions	N/A

Coverage Duration	CSU initial: 6 months, All others: Plan Year
Other Criteria	N/A

XOSPATA

Products Affected

- Xospata

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and FLT3 rearrangement
Exclusion Criteria	N/A
Required Medical Information	For myeloid, lymphoid, or mixed lineage neoplasms with eosinophilia and FMS-like tyrosine kinase 3 (FLT3) rearrangement: the disease is in chronic or blast phase.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XPOVIO

Products Affected

- Xpovio
- Xpovio 60 Mg Twice Weekly
- Xpovio 80 Mg Twice Weekly

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Histologic transformation of indolent lymphomas to diffuse large B-cell lymphoma, acquired immunodeficiency syndrome (AIDS)-related B-cell lymphoma, high-grade B-cell lymphoma
Exclusion Criteria	N/A
Required Medical Information	For multiple myeloma: Patient must have been treated with at least one prior therapy. For B-cell lymphomas: Patient must have been treated with at least two lines of systemic therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XTANDI

Products Affected

- Xtandi

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the treatment of castration-resistant prostate cancer or metastatic castration-sensitive prostate cancer: The requested drug will be used in combination with a gonadotropin-releasing hormone (GnRH) analog or after bilateral orchiectomy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

XYREM

Products Affected

- Sodium Oxybate

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the treatment of excessive daytime sleepiness in a patient with narcolepsy, initial request: 1) The diagnosis has been confirmed by sleep lab evaluation, AND 2) The patient meets one of the following criteria: a) if the patient is 17 years of age or younger, the patient has experienced an inadequate treatment response or intolerance to at least one central nervous system (CNS) stimulant drug (e.g., amphetamine, dextroamphetamine, methylphenidate), OR has a contraindication that would prohibit a trial of central nervous system (CNS) stimulant drugs (e.g., amphetamine, dextroamphetamine, methylphenidate), b) If the patient is 18 years of age or older, the patient has experienced an inadequate treatment response or intolerance to at least one central nervous system (CNS) wakefulness promoting drug (e.g., armodafinil, modafinil), OR has a contraindication that would prohibit a trial of central nervous system (CNS) wakefulness promoting drugs (e.g., armodafinil, modafinil). For the treatment of cataplexy in a patient with narcolepsy, initial request: The diagnosis has been confirmed by sleep lab evaluation. If the request is for a continuation of therapy, then the patient experienced a decrease in daytime sleepiness with narcolepsy or a decrease in cataplexy episodes with narcolepsy.
Age Restrictions	7 years of age or older
Prescriber Restrictions	Prescribed by or in consultation with a sleep disorder specialist or neurologist
Coverage Duration	Plan Year
Other Criteria	N/A

ZARXIO

Products Affected

- Zarxio

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Neutropenia in myelodysplastic syndromes (MDS), agranulocytosis, neutropenia in aplastic anemia, human immunodeficiency virus (HIV)-related neutropenia, neutropenia related to renal transplant, hematopoietic syndrome of acute radiation syndrome
Exclusion Criteria	Use of the requested product within 24 hours prior to or following chemotherapy.
Required Medical Information	For prophylaxis or treatment of myelosuppressive chemotherapy-induced febrile neutropenia (FN) patient must meet both of the following: 1) Patient has a solid tumor or non-myeloid cancer, and 2) Patient has received, is currently receiving, or will be receiving treatment with myelosuppressive anti-cancer therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	6 months
Other Criteria	N/A

ZEJULA

Products Affected

- Zejula

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Uterine leiomyosarcoma
Exclusion Criteria	N/A
Required Medical Information	For uterine leiomyosarcoma: 1) the requested drug is used as second-line therapy AND 2) the patient has BRCA-altered disease.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZELBORAF

Products Affected

- Zelboraf

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Non-small cell lung cancer, hairy cell leukemia, thyroid carcinoma (i.e., papillary carcinoma, follicular carcinoma, and Hurthle cell carcinoma), central nervous system cancer (i.e., glioma, astrocytoma, glioblastoma, pediatric diffuse high-grade glioma), adjuvant systemic therapy for cutaneous melanoma, Langerhans cell histiocytosis.
Exclusion Criteria	N/A
Required Medical Information	For central nervous system (CNS) cancer (i.e., glioma, astrocytoma, glioblastoma, pediatric diffuse high-grade glioma): 1) The tumor is positive for BRAF V600E mutation, AND 2) The requested drug will be used in combination with cobimetinib OR the requested drug is being used for the treatment of pediatric diffuse high-grade glioma. For melanoma: 1) The tumor is positive for BRAF V600 activating mutation (e.g., V600E or V600K), AND 2) the requested drug will be used as a single agent, or in combination with cobimetinib, AND 3) The requested drug will be used for either of the following: a) unresectable, limited resectable, or metastatic disease, or b) adjuvant systemic therapy. For Erdheim-Chester Disease and Langerhans Cell Histiocytosis: Tumor is positive for BRAF V600 mutation. For non-small cell lung cancer: 1) The tumor is positive for the BRAF V600E mutation, AND 2) The patient has recurrent, advanced, or metastatic disease. For papillary, follicular, and hurthle cell thyroid carcinoma: 1) The tumor is positive for BRAF mutation, AND 2) The disease is not amenable to radioactive iodine (RAI) therapy.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZIRABEV

Products Affected

- Zirabev

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Ampullary adenocarcinoma, breast cancer, central nervous system (CNS) cancers, malignant pleural mesothelioma, malignant peritoneal mesothelioma, pericardial mesothelioma, tunica vaginalis testis mesothelioma, soft tissue sarcomas, uterine neoplasms, endometrial carcinoma, vulvar cancers, small bowel adenocarcinoma, and ophthalmic-related disorders: diabetic macular edema, neovascular (wet) age-related macular degeneration including polypoidal choroidopathy and retinal angiomatous proliferation subtypes, macular edema following retinal vein occlusion, proliferative diabetic retinopathy, choroidal neovascularization, neovascular glaucoma and retinopathy of prematurity.
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	Coverage under Part D will be denied if coverage is available under Part A or Part B as the medication is prescribed and dispensed or administered for the individual.

ZOLINZA

Products Affected

- Zolanza

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Mycosis fungoides (MF)/Sezary syndrome (SS)
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZONISADE

Products Affected

- Zonisade

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For adjunctive treatment of partial-onset seizures (i.e., focal-onset seizures): 1) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to a generic anticonvulsant AND the patient has experienced an inadequate treatment response, intolerance, or has a contraindication to any of the following: Aptiom, Xcopri, Spritam OR 2) The patient has difficulty swallowing solid oral dosage forms (e.g., tablets, capsules).
Age Restrictions	16 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZTALMY

Products Affected

- Ztalmy

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	N/A
Age Restrictions	2 years of age or older
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZURZUVAE

Products Affected

- Zurzuvae

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	For the treatment of postpartum depression (PPD): diagnosis was confirmed using standardized rating scales that reliably measure depressive symptoms (e.g., Hamilton Depression Rating Scale [HDRS], Edinburgh Postnatal Depression Scale [EPDS], Patient Health Questionnaire 9 [PHQ9], Montgomery-Asberg Depression Rating Scale [MADRS], Beck's Depression Inventory [BDI], etc.).
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	1 month
Other Criteria	N/A

ZYDELIG

Products Affected

- Zydelig

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Small lymphocytic lymphoma (SLL)
Exclusion Criteria	N/A
Required Medical Information	For CLL/SLL: the requested drug is used as second-line or subsequent therapy
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZYKADIA

Products Affected

- Zykadia TABS

PA Criteria	Criteria Details
Indications	All FDA-approved Indications, Some Medically accepted Indications.
Off-Label Uses	Recurrent ALK-positive non-small cell lung cancer (NSCLC), recurrent, advanced, or metastatic ROS1-positive NSCLC, inflammatory myofibroblastic tumor (IMT), brain metastases from NSCLC.
Exclusion Criteria	N/A
Required Medical Information	For NSCLC: the patient has recurrent, advanced, or metastatic ALK-positive or ROS1-positive disease. For inflammatory myofibroblastic tumor: the disease is ALK-positive. For brain metastases from NSCLC: the patient has ALK-positive NSCLC.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

ZYPREXA RELPREVV

Products Affected

- Zyprexa Relprevv

PA Criteria	Criteria Details
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Exclusion Criteria	N/A
Required Medical Information	Tolerability with oral olanzapine has been established.
Age Restrictions	N/A
Prescriber Restrictions	N/A
Coverage Duration	Plan Year
Other Criteria	N/A

PART B VERSUS PART D

Products Affected

- Abelcet
- Acetylcysteine INHALATION SOLN
- Acyclovir Sodium INJ 50MG/ML
- Albuterol Sulfate NEBU 0.083%, 0.63MG/3ML, 1.25MG/3ML, 2.5MG/0.5ML
- Amphotericin B INJ
- Amphotericin B Liposome
- Aprepitant CAPS
- Astagraf XL
- Azathioprine INJ
- Azathioprine TABS 50MG
- Budesonide SUSP
- Clinimix 4.25%/dextrose 10%
- Clinimix 4.25%/dextrose 5%
- Clinimix 5%/dextrose 15%
- Clinimix 5%/dextrose 20%
- Clinimix 6/5
- Clinimix 8/10
- Clinimix 8/14
- Clinisol Sf 15%
- Clinolipid
- Cromolyn Sodium NEBU
- Cyclosporine CAPS
- Cyclosporine INJ
- Cyclosporine Modified
- Dextrose 50%
- Dextrose 70%
- Emend SUSR
- Engerix-b
- Epoprostenol Sodium
- Everolimus TABS 0.25MG, 0.5MG, 0.75MG, 1MG
- Fiasp Pumpcart
- Gamastan
- Ganciclovir INJ 500MG, 500MG/10ML
- Gengraf CAPS 100MG, 25MG
- Gengraf SOLN
- Granisetron Hydrochloride TABS
- Hepatamine INJ 62MEQ/L; 770MG/100ML; 600MG/100ML; 3MEQ/L; 20MG/100ML; 900MG/100ML; 240MG/100ML; 900MG/100ML; 1100MG/100ML; 610MG/100ML; 100MG/100ML; 100MG/100ML; 115MG/100ML; 800MG/100ML; 500MG/100ML; 450MG/100ML; 66MG/100ML; 840MG/100ML
- Heplisav-b
- Humulin R U-500 (concentrated)
- Hydromorphone Hcl INJ 10MG/ML, 1MG/ML, 4MG/ML
- Hydromorphone Hydrochloride INJ 0.25MG/0.5ML, 1MG/ML, 2MG/ML, 4MG/ML, 50MG/5ML
- Imovax Rabies (h.d.c.v.)
- Ipratropium Bromide INHALATION SOLN 0.02%
- Ipratropium Bromide/albuterol Sulfate
- Isolyte-s INJ 27MEQ/L; 98MEQ/L; 23MEQ/L; 3MEQ/L; 5MEQ/L; 140MEQ/L
- Isolyte-s Ph 7.4
- Jynneos
- Levalbuterol NEBU
- Levalbuterol Hcl NEBU
- Levalbuterol Hydrochloride NEBU 0.63MG/3ML
- Methylprednisolone TABS
- Methylprednisolone Acetate INJ 40MG/ML, 80MG/ML
- Methylprednisolone Sodium Succinate INJ 1000MG, 125MG
- Methylprednisolone Sodiamsuccinate INJ 40MG
- Morphine Sulfate INJ 0.5MG/ML, 10MG/ML, 1MG/ML, 2MG/ML, 4MG/ML, 50MG/ML, 5MG/ML, 8MG/ML

- Morphine Sulfate/sodium Chloride INJ 1MG/ML
- Mycophenolate Mofetil CAPS
- Mycophenolate Mofetil INJ
- Mycophenolate Mofetil SUSR
- Mycophenolate Mofetil TABS
- Mycophenolic Acid Dr
- Nulojix
- Nutrilipid
- Ondansetron Hcl SOLN
- Ondansetron Hcl TABS 24MG
- Ondansetron Hydrochloride TABS
- Ondansetron Odt
- Pentamidine Isethionate INHALATION SOLR
- Plenamine INJ 147.4MEQ/L; 2.17GM/100ML; 1.47GM/100ML; 434MG/100ML; 749MG/100ML; 1.04GM/100ML; 894MG/100ML; 749MG/100ML; 1.04GM/100ML; 1.18GM/100ML; 749MG/100ML; 1.04GM/100ML; 894MG/100ML; 592MG/100ML; 749MG/100ML; 250MG/100ML; 39MG/100ML; 960MG/100ML
- Prednisolone SOLN
- Prednisolone Sodium Phosphate ORAL SOLN 10MG/5ML, 15MG/5ML, 20MG/5ML, 25MG/5ML, 5MG/5ML
- Prednisone SOLN
- Prednisone TABS 10MG, 1MG, 2.5MG, 20MG, 50MG, 5MG
- Prednisone Intensol
- Prehevbrio
- Premasol INJ 52MEQ/L; 1760MG/100ML; 880MG/100ML; 34MEQ/L; 1760MG/100ML; 372MG/100ML; 406MG/100ML; 526MG/100ML; 492MG/100ML; 492MG/100ML; 526MG/100ML; 356MG/100ML; 356MG/100ML; 390MG/100ML; 34MG/100ML; 152MG/100ML
- Prograf PACK
- Prosol
- Rabavert
- Recombivax Hb
- Sandimmune SOLN
- Sirolimus SOLN
- Sirolimus TABS
- Tacrolimus CAPS
- Tpn Electrolytes INJ 29.5MEQ/20ML; 4.5MEQ/20ML; 35MEQ/20ML; 5MEQ/20ML; 20MEQ/20ML; 35MEQ/20ML
- Travasol INJ 52MEQ/L; 1760MG/100ML; 880MG/100ML; 34MEQ/L; 1760MG/100ML; 372MG/100ML; 406MG/100ML; 526MG/100ML; 492MG/100ML; 492MG/100ML; 526MG/100ML; 356MG/100ML; 500MG/100ML; 356MG/100ML; 390MG/100ML; 34MG/100ML; 152MG/100ML
- Trophamine INJ 0.54GM/100ML; 1.2GM/100ML; 0.32GM/100ML; 0; 0; 0.5GM/100ML; 0.36GM/100ML; 0.48GM/100ML; 0.82GM/100ML; 1.4GM/100ML; 1.2GM/100ML; 0.34GM/100ML; 0.48GM/100ML; 0.68GM/100ML; 0.38GM/100ML; 5MEQ/L; 0.025GM/100ML; 0.42GM/100ML; 0.2GM/100ML; 0.24GM/100ML; 0.78GM/100ML

Details

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

INDEX

A

Abelcet	346
Abiraterone.....	1
Abiraterone Acetate.....	1
Accutane.....	151
Acetylcysteine	346
Acitretin.....	3
Actimmune	4
Acyclovir Sodium	346
Adalimumab-aacf (2 Pen)	138
Adempas.....	5
Aimovig.....	6
Akeega.....	7
Albuterol Sulfate	346
Alecensa	8
Alosetron	9
Alosetron Hydrochloride.....	9
Alpha1-proteinase Inhibitor	10
Alunbrig	11
Alyq.....	270
Ambrisentan	12
Amitriptyline Hcl	127
Amitriptyline Hydrochloride.....	127
Amnesteem.....	151
Amphotericin B.....	346
Amphotericin B Liposome	346
Aprepitant.....	346
Arcalyst	13
Armodafinil	14
Asparlas.....	15
Astagraf XL.....	346
Atovaquone	184
Aubagio	16
Augtyro	17
Austedo	18
Austedo Xr	18
Austedo Xr Patient Titration Kit.....	18

Auvelity.....	19
Avonex	20
Avonex Aet.....	20
Avonex Pen.....	20
Ayvakit.....	21
Azathioprine.....	346

B

Balversa.....	22
Banzel	23
Benlysta.....	24
Benztropine Mesylate	119
Berinert	25
Besremi	26
Betaseron.....	27
Bexarotene	28, 277
Bosentan.....	29
Bosulif.....	30
Braftovi	31
Briviact.....	32, 33
Briviact Inj	33
Bronchitol	34
Bronchitol Tolerance Test.....	34
Brukinsa	35
Budesonide.....	346
Buprenorphine.....	36, 37
Buprenorphine Hcl.....	36
Buprenorphine Patch.....	37
Bydureon.....	38
Bydureon Bcise.....	38
Byetta	39

C

Cabometyx	40
Calcipotriene.....	41
Calcitrene	41
Calcitriol	41
Calquence.....	42

Caprelsa.....	43
Carac.....	44
Carbaglu	45
Carbinoxamine Maleate	128
Carglumic Acid	45
Cayston.....	46
Cerdelga	47
Chantix Aet	48
Chlordiazepoxide - 65	49
Chlordiazepoxide Hcl.....	49
Chlordiazepoxide Hydrochloride	49
Chlorzoxazone.....	131
Claravis.....	151
Clemastine Fumarate.....	129
Clinimix 4.25%/dextrose 10%	346
Clinimix 4.25%/dextrose 5%	346
Clinimix 5%/dextrose 15%	346
Clinimix 5%/dextrose 20%	346
Clinimix 6/5	346
Clinimix 8/10	346
Clinimix 8/14	346
Clinisol Sf 15%	346
Clinolipid.....	346
Clobazam.....	50
Clomipramine.....	51
Clomipramine Hydrochloride	51
Clorazepate.....	52
Clorazepate Dipotassium.....	52
Clozapine Odt.....	53
Colistimethate Sodium	54
Coly-mycin.....	54
Cometriq.....	55
Copaxone.....	108
Copiktra.....	56
Cotellic	57
Cromolyn Sodium	346
Cyclobenzaprine Hydrochloride	131
Cyclophosphamide.....	58
Cyclophosphamide Oral.....	58
Cyclosporine	346
Cyclosporine Modified.....	346
Cyproheptadine Hcl.....	120
Cyproheptadine Hydrochloride.....	120
Cystagon.....	59

Cystaran	60
----------------	----

D

Dalfampridine	61
Dalfampridine Er	61
Daurismo.....	62
Deferasirox.....	63
Demser	64
Desipramine Hydrochloride.....	127
Desvenlafaxine.....	65
Desvenlafaxine Er.....	65
Dextrose 50%.....	346
Dextrose 70%.....	346
Dhe Nasal.....	66
Diacomit.....	67
Diazepam	68
Diazepam Aet.....	68
Diazepam Intensol	68
Dicyclomine Hcl.....	115
Dicyclomine Hydrochloride.....	115
Dihydroergotamine Mesylate.....	66, 115
Dipyridamole	115
Disopyramide Phosphate	115
Doptelet.....	69
Doxepin Hcl	121
Doxepin Hydrochloride	121, 291
Doxycycline	210
Drizalma.....	70
Drizalma Sprinkle	70
Dronabinol	71
Droxidopa	196
Dupixent.....	72

E

Eligard.....	74
Emend	346
Emsam.....	75
Enbrel.....	76
Enbrel Mini	76
Enbrel Sureclick.....	76
Endari.....	78
Engerix-b.....	346
Enstilar	41
Epclusa.....	79
Epidiolex.....	80

Epo	81
Epoprostenol Sodium	346
Eprontia	82
Ergotamine	83
Ergotamine Tartrate/caffeine.....	83
Erivedge	84
Erleada.....	85
Erlotinib.....	86
Erlotinib Hydrochloride	86
Esbriet.....	87
Everolimus	88, 346
Exkivity	90

F

Fanapt.....	91
Fanapt Titration Pack	91
Fasenra	92
Fasenra Pen	92
Fentanyl.....	93
Fentanyl Citrate Oral Transmucosal	211
Fentanyl Patch.....	93
Fetzima.....	94
Fetzima Titration Pack	94
Fiasp Pumpcart.....	346
Fingolimod	106
Fintepla.....	95
Firmagon	96
Flucytosine	97
Fluorouracil.....	44, 98
Forteo	99
Fotivda.....	101
Fruzaqla.....	102
Fycompa.....	103

G

Gamastan.....	346
Gammaked	154
Gamunex-c	154
Ganciclovir.....	346
Gattex	104
Gavreto.....	105
Gefitinib	150
Gengraf.....	346
Genotropin.....	109
Genotropin Miniquick.....	109

Gilenya.....	106
Gilotrif.....	107
Glatiramer	108
Granisetron Hydrochloride	346
Growth Hormone	109
Guanfacine Er	115
Guanfacine Hydrochloride.....	115

H

Haegarda	111
Harvoni	112
Hepatamine	346
Heplisav-b.....	346
Hetlioz.....	113
Hetlioz Lq	114
High Risk Medication - 65.....	115
Hrm - Anticonvulsants - 65.....	116
Hrm - Hypnotics - 65	117
Hrm - Scopolamine - 65.....	118
Hrm-antiparkinson - 65.....	119
Hrm-cyproheptadine - 65.....	120
Hrm-doxepin - 65.....	121
Hrm-hydroxyzine - 65.....	122
Hrm-hydroxyzine Inj - 65	124
Hrm-promethazine - 65.....	126
Hrms - Antidepressants.....	127
Hrms - Carbinoxamine.....	128
Hrms - Clemastine- 65	129
Hrms - Teas.....	130
Hrm-skeletal Muscle Relaxants - 65.....	131
Humira	132
Humira Pediatric Crohns Disease Starter Pack..	132
Humira Pen	132
Humira Pen-cd/uc/hs Starter	132
Humira Pen-pediatric Uc Starter Pack.....	132
Humira Pen-ps/uv Starter.....	132
Humulin R U-500 (concentrated)	346
Hydrocodone Bitartrate Er.....	149
Hydromorphone Hcl	346
Hydromorphone Hydrochloride.....	346
Hydroxyzine Hcl.....	122, 124
Hydroxyzine Hydrochloride	122, 124
Hydroxyzine Pamoate.....	122
Hypnotic-benzodiazepines - 65.....	134
Hysingla Er	149

I	
Ibrance.....	135
Icatibant.....	136
Icatibant Acetate.....	136
Iclusig.....	137
Idacio.....	138
Idacio (2 Pen).....	138
Idacio (2 Syringe).....	138
Idacio Starter Package For Crohns Disease.....	138
Idacio Starter Package For Plaque Psoriasis.....	138
Idhifa.....	140
Imatinib.....	141
Imatinib Mesylate.....	141
Imbruvica.....	142
Imipramine Hcl.....	127
Imipramine Hydrochloride.....	127
Imovax Rabies (h.d.c.v.).....	346
Inbrija.....	144
Increlex.....	145
Inlyta.....	146
Inqovi.....	147
Inrebic.....	148
Ipratropium Bromide.....	346
Ipratropium Bromide/albuterol Sulfate.....	346
Ir Before Er.....	149
Iressa.....	150
Isolyte-s.....	346
Isolyte-s Ph 7.4.....	346
Isotretinoin.....	151
Itraconazole.....	152
Ivermectin.....	153
Ivermectin Tab.....	153
Ivig.....	154
Iwilfin.....	155
J	
Jakafi.....	156
Javygtor.....	251
Jaypirca.....	157
Jynneos.....	346
K	
Kalydeco.....	158
Kesimpta.....	159
Ketoconazole.....	160
Ketorolac Tromethamine.....	115
Kevzara.....	161
Kisqali.....	162
Kisqali Femara 200 Dose.....	162
Kisqali Femara 400 Dose.....	162
Kisqali Femara 600 Dose.....	162
Korlym.....	163
Koselugo.....	164
Krazati.....	165
Kristalose.....	166
L	
Lapatinib.....	167
Lapatinib Ditosylate.....	167
Lenalidomide.....	240
Lenvima.....	168
Lenvima 10 Mg Daily Dose.....	168
Lenvima 12mg Daily Dose.....	168
Lenvima 14 Mg Daily Dose.....	168
Lenvima 18 Mg Daily Dose.....	168
Lenvima 20 Mg Daily Dose.....	168
Lenvima 24 Mg Daily Dose.....	168
Lenvima 4 Mg Daily Dose.....	168
Lenvima 8 Mg Daily Dose.....	168
Leuprolide.....	169
Leuprolide Acetate.....	169
Levalbuterol.....	346
Levalbuterol Hcl.....	346
Levalbuterol Hydrochloride.....	346
Lidocaine.....	170, 292
Lidocaine Patches.....	170
Lidocan III.....	170
Linezolid.....	171
Lonsurf.....	172
Lorbrena.....	173
Lumakras.....	174
Lupron Depot (1-month).....	176
Lupron Depot (3-month).....	176
Lupron Depot-ped (1-month).....	175
Lupron Depot-ped (3-month).....	175
Lupron Depot-ped (6-month).....	175
Lupron Ped.....	175
Lupron-endometriosis.....	176
Lynparza.....	177
Lyrica Cr.....	178
	352

Lytgobi 179

M

Mavyret 180
Mekinist..... 181
Mektovi 182
Memantine..... 183
Memantine Hcl Titration Pak..... 183
Memantine Hydrochloride 183
Memantine Hydrochloride Er 183
Mepron Aet 184
Metformin Er..... 185
Metformin Hydrochloride Er 185
Methadone Hcl 149
Methadone Hydrochloride 149
Methscopolamine Bromide 115
Methylprednisolone..... 346
Methylprednisolone Acetate 346
Methylprednisolone Sodium Succinate..... 346
Methylprednisolone Sodiumsuccinate 346
Methyltestosterone 186
Metyrosine..... 64
Mifepristone 163
Modafinil..... 187
Morphine Sulfate..... 346
Morphine Sulfate Er 149
Morphine Sulfate/sodium Chloride..... 347
Mounjaro 188
Mycophenolate Mofetil..... 347
Mycophenolic Acid Dr..... 347

N

Natpara 189
Nayzilam 190
Nerlynx..... 191
Nexavar 192
Ninlaro..... 194
Nitisinone 195
Northera..... 196
Nubeqa 197
Nuedexta..... 198
Nulojix..... 347
Nuplazid 199
Nurtec 200
Nutrilipid..... 347

O

Octagam 154
Octreotide..... 201
Octreotide Acetate 201
Odomzo..... 202
Ofev..... 203
Ogsiveo 204
Ojjaara..... 205
Omega-3..... 206
Omega-3-acid Ethyl Esters 206
Oncaspar 207
Ondansetron Hcl 347
Ondansetron Hydrochloride..... 347
Ondansetron Odt 347
Onureg..... 208
Opsumit..... 209
Oracea 210
Oracea Aet 210
Oral-intranasal Fentanyl..... 211
Orgovyx 212
Orkambi 213
Orserdu..... 214
Otezla 215
Oxandrolone..... 216
Oxazepam 217
Oxazepam - 65 217
Ozempic 218

P

Panretin 219
Part B Versus Part D 346
Pazopanib Hydrochloride..... 319
Pegasys..... 220
Pemazyre..... 221
Pentamidine Isethionate 347
Perphenazine/amitriptyline 115
Phenobarbital 116
Phenobarbital Sodium..... 116
Phenylbutyrate 222
Piqray 223
Piqray 200mg Daily Dose..... 223
Piqray 250mg Daily Dose..... 223
Piqray 300mg Daily Dose..... 223
Pirfenidone 87

Plenammine	347
Pomalyst	224
Posaconazole	225
Posaconazole Dr	225
Prednisolone	347
Prednisolone Sodium Phosphate	347
Prednisone	347
Prednisone Intensol	347
Pregabalin	226
Pregabalin Er	178
Prehevbrio	347
Premasol	347
Pretomanid	227
Pretomanid Aet	227
Prevymis	228
Privigen	154
Procrit	81
Prograf	347
Prolastin-c	10
Promacta	229
Promethazine Hcl	126
Promethazine Hydrochloride	126
Promethazine Hydrochloride Plain	126
Promethegan	126
Prosol	347
Protriptyline Hcl	130
Pulmozyme	231

Q

Qinlock	232
Quetiapine Fumarate Er	233
Quetiapine Xr	233
Quinine Sulfate	235
Qulipta	236

R

Rabavert	347
Recombivax Hb	347
Regranex	237
Repatha	238
Repatha Pushttronex System	238
Repatha Sureclick	238
Retevmo	239
Revlimid	240
Rezlidhia	241

Rezurock	242
Rinvoq	243
Rozlytrek	245
Rubraca	246
Rufinamide	23
Rybelsus	247
Rydapt	248

S

Sajazir	136
Sandimmune	347
Sandostatin Lar	249
Sandostatin Lar Depot	249
Sapropterin	251
Sapropterin Dihydrochloride	251
Scemblix	252
Scopolamine	118
Signifor	253
Sildenafil	254, 255
Sildenafil Citrate	254
Sildenafil Inj	255
Sirolimus	347
Sirturo	256
Skyrizi	257
Skyrizi Pen	257
Sodium Oxybate	334
Sodium Phenylbutyrate	222
Somatuline Depot	258
Somavert	259
Sorafenib Tosylate	192
Spritam	260
Sprycel	261
Stelara	263
Stivarga	264
Sunitinib Malate	265
Sutent	265
Symlin	267
Symlinpen 120	267
Symlinpen 60	267
Sympazan	268

T

Tabrecta	269
Tacrolimus	347
Tadalafil	270

Tadalafil (pah).....	270	Trophamine.....	347
Tafinlar.....	271	Trulicity.....	297
Tagrisso.....	273	Truqap.....	298
Taltz.....	274	Truseltiq.....	299
Talzenna.....	276	Truxima.....	300
Targretin Topical.....	277	Tukysa.....	302
Tasigna.....	278	Turalio.....	303
Tasimelteon.....	113	<i>U</i>	
Tazarotene.....	280	Ubrelvy.....	304
Tazorac.....	280	<i>V</i>	
Tazverik.....	281	Valchlor.....	305
Tecfidera.....	282	Valtoco.....	306
Tecfidera Starter Pack.....	282	Valtoco 10 Mg Dose.....	306
Tecvayli.....	283	Valtoco 15 Mg Dose.....	306
Temazepam.....	134, 284	Valtoco 20 Mg Dose.....	306
Temazepam 30mg - 65.....	284	Valtoco 5 Mg Dose.....	306
Tepmetko.....	285	Vanflyta.....	307
Teriparatide.....	99	Varenicline Starting Month Box.....	48
Testosterone Enanthate.....	286	Varenicline Tartrate.....	48
Testosterone Enanthate Inj.....	286	Venclexta.....	308
Tetrabenazine.....	287	Venclexta Starting Pack.....	308
Thalomid.....	288	Ventavis.....	309
Thioridazine Hcl.....	115	Verquvo.....	310
Tibsovo.....	289	Versacloz.....	311
Tobramycin.....	290	Verzenio.....	312
Topical Doxepin.....	291	Vigabatrin.....	313
Topical Lidocaine.....	292	Vigadrone.....	313
Topical Tretinoin.....	293	Vigpoder.....	313
Toremifene.....	294	Vitrakvi.....	314
Toremifene Citrate.....	294	Vizimpro.....	315
Tpn Electrolytes.....	347	Vonjo.....	316
Tracleer.....	29	Voriconazole.....	317
Travasol.....	347	Vosevi.....	318
Tretinoin.....	293	Votrient.....	319
Tretinoin Microsphere.....	293	Vumerity.....	320
Tretinoin Microsphere Pump.....	293	<i>W</i>	
Triazolam.....	134	Welireg.....	321
Trientine.....	295	<i>X</i>	
Trientine Hydrochloride.....	295	Xalkori.....	322
Trihexyphenidyl Hcl.....	119	Xeljanz.....	323
Trihexyphenidyl Hydrochloride.....	119	Xeljanz Xr.....	323
Trikafta.....	296		
Trimethobenzamide Hydrochloride.....	115		
Trimipramine Maleate.....	130		

Xermelo.....	325	Zarxio.....	335
Xgeva	326	Zejula	336
Xhance.....	327	Zelboraf.....	337
Xhance Mapd	327	Zenatane.....	151
Xifaxan	328	Zirabev	338
Xolair.....	329	Zolinza	339
Xospata.....	331	Zolpidem Tartrate	117
Xpovio.....	332	Zonisade.....	340
Xpovio 60 Mg Twice Weekly.....	332	Ztalmy	341
Xpovio 80 Mg Twice Weekly.....	332	Zurzuvae	342
Xtandi.....	333	Zydelig.....	343
Xyrem.....	334	Zykadia	344
Z		Zyprexa Relprevv.....	345
Zaleplon.....	117	Zytiga	1