

Aetna Better Health® of Florida (MEDICAID)

Antipsychotic (6 to < 18 Years of Age)

Maximum Length of Approval = 180 Days

Note: Form must be completed in full.

											Date	e of	Birt	h (M	M/DE	/////	Y)											
Reci	pient	<u>'s ru</u>	II Nam																									
Pres	cribe	r's Fi	ull Nar	ne	1		<u> </u>			I	1	1					1	1	1	1	1	1	1	1	1	1	1	1
Pres	cribe	r's N	PI																									
Pres	cribe	r's P	hone I	Numb	er	1					1						Pre	scrik	er's	Fax	Num	ber		7				
			-			-														-				-				
PRO	VIDEF	R TYP	E OR S	PECI	ALTY:										CI	ILD I	JNDE	R ST.	ATE (CARE	cus	тор	Y:		Yes			No
ΡΑΤ	ENT:			Male			Fema	le							ME	DICAT	ION F	REQU	EST:			New			Con	tinua	tion	
HEIC	SHT:				🗌 in	/	🗌 cr	n V	VEIG	HT:					🗌 lb	s /	🗌 kę	js	BMI:				*	BMI %	6: _			
_															BN	II Cal	culato	or: * <u>h</u>	ttps://	<u>www</u>	.cdc.g	<u>ov/he</u>	althy	weigł	<u>nt/bmi</u>	<u>/calcı</u>	<u>ılator.</u>	<u>html</u>
1.			tion R sted /	-			(c)		Strength Directions									Quantity										
			sieu r	mup	sycn	Ulic	(3)		ou c	FIIGL																		
		9440		_	-					<u> </u>							Dire	cuo	113						Qu	anner	•9	
																	Dire	:010	113						qu		.y	
2		-															Dire								44		<u>.</u>	
2.	Dia	gno:						Disru				/ior	Disc	rder						ood	Dys	regu	latio	n Di			.y	
2.	Dia	gno:		ectru	m			Disru Schiz	ıptiv	e Be	ehav	vior	Disc	rder			Disi	ruptiv	ve M		•	regu			sord	er		
2.	Dia	gno: ADH Autis	ID						ıptiv zopł	re Be	ehav			rder			Disi	ruptiv	ve M		•	Ũ			sord	er		
2. 3.	Dia Dia Dia	gnos ADF Autis Bipo	ID sm Sp	sorde t oms	r : [A¢		Schiz Schiz	ıptiv zopł zoaf	e Be nreni fecti	ehav	Diso	rder	rder	Irrita	ability er: _	Disi Oth	ruptiv er:	ve M	njuri	ous	Beha	avio	r	sord	er		-
	Dia Dia D Tar (che	gnos ADF Autis Bipo rget s	ID sm Sp Iar Dis Symp t	sorde toms appl	r : [y)		ggres	Schiz Schiz	ıptiv zopł zoaf	e Be nreni fecti	ehav ia ve D	Diso Isivi	rder ty		Irrita	er: _	Disi Oth	ruptiv er:	ve M	njuri	ous	Beha	avio	r	sord	er		-
3.	Dia Dia Dia Tar (cho	gnos ADH Autis Bipo get \$ eck a	ID sm Sp lar Dis Symp t all that	sorde toms appl arget	r : [y) Sym	ptic	ggres	Schiz Schiz	ıptiv zopł zoaf	e Be nreni fecti	ehav ia ve D mpul	Diso Isivi' I	rder ty	 2 Mo	Irrita Oth	er: _ ate	Disi Oth /		ve M	njuri	ous	Beha	avio	r	sord	er		
3 . 4 .	Dia Dia Dia C Tar (che Sev Fur	gnos ADH Autis Bipo get \$ eck a verity nctio	ID sm Sp Jar Dis Symp t all that / of T a	sorde toms appl arget	r [y) Sym ment	nptic ::	ggres	Schiz Schiz	uptiv zoph zoaf ı [[e Be nreni fecti	èhav ia ve D mpul Mild Mild	Diso Isivi I	rder ty	 2 Mo	Irrita Oth	er: _ ate	Disi Oth /		ve M	njuri	ous	Beha	avio	r	sord	er	reme	
3. 4. 5.	Dia Dia Dia C Tar (che Sev Fur	gnos ADH Autis Bipo get \$ eck a verity nctio	ID sm Sp lar Dis Sympt all that y of Ta nal In	sorde toms appl arget pair ipsyd	r [y) Sym ment choti	nptic :: c Tr	ggres oms:	Schiz Schiz	uptiv zoph zoaf ı [[e Be nreni fecti] Ir] 1] 1] 2 Mo	èhav ia ve D mpul Mild Mild	Diso Isivi I I	rder ty	 2 Mo	Irrita Oth	er: _ ate ate	Disi Oth /	ruptiv er: 3 Ma 3 Ma	ve M Self I	njuri	ous	Beha	eve eve	r re re	sord	er 5 Ext	reme	
3. 4. 5.	Dia Dia Dia C Tar (che Sev Fur	gnos ADH Autis Bipo get \$ eck a verity nctio	ID sm Sp Jar Dis Sympt all that y of Ta nal In s Ant	sorde toms appl arget pair ipsyd	r [y) Sym ment choti	nptic :: c Tr	ggres oms:	Schiz Schiz	uptiv zoph zoaf ı [[e Be nreni fecti] Ir] 1] 1] 2 Mo	ehav ia ve D mpul Mild Mild	Diso Isivi I I	rder ty	 2 Mo	Irrita Oth	er: _ ate ate	Disi Oth /	ruptiv er: 3 Ma 3 Ma	ve M Self I	njuri	ous	Beha] 4 S] 4 S	eve eve	r re re	sord	er 5 Ext	reme	
3. 4. 5.	Dia Dia Dia C Tar (che Sev Fur	gnos ADH Autis Bipo get \$ eck a verity nctio	ID sm Sp Jar Dis Sympt all that y of Ta nal In s Ant	sorde toms appl arget pair ipsyd	r [y) Sym ment choti	nptic :: c Tr	ggres oms:	Schiz Schiz	uptiv zoph zoaf ı [[e Be nreni fecti] Ir] 1] 1] 2 Mo	ehav ia ve D mpul Mild Mild	Diso Isivi I I	rder ty	 2 Mo	Irrita Oth	er: _ ate ate	Disi Oth /	ruptiv er: 3 Ma 3 Ma	ve M Self I	njuri	ous	Beha] 4 S] 4 S	eve eve	r re re	sord	er 5 Ext	reme	



CoverMyMeds® or SureScripts

Aetna Better Health® of Florida (MEDICAID)

Antipsychotic (6 to < 18 Years of Age)

Maximum Length of Approval = 180 Days

Note: Form must be completed in full.

	List all other psychotherapeutic r	nedications th	nationt is t	aking conc		with th	o an	tinev	choti			
	antidepressants, mood stabilizer				unentry			upsy	chou	C (1.C	•,	
	Psychotherapeutic Medication	Dose/day	<u>י</u> ף:	sychothera	peutic M	edicati	on	Dose/day				
	Rationale for prescribing antipsy	chotic above r	naximum rec	ommende	d dose?	(if appl	icabl	e)				
	Is your intent to target lower dose ☐ Yes ☐ No	e antipsychoti	c treatment?									
		ve entinevele			liaahla).							
	Rationale for prescribing 2 or mo	re antipsycho	tics for >60 d	ays (if app	licable):							
	If your request is for two antipsyc	chotics:										
	Is the plan to cross taper, with an		onotherapy r	esumed wi	thin the	next 60) day	s?				
		YES, please pro					-					
	Have metabolic monitoring labs*	(fasting lipids	and glucose) been perl	ormed w	vithin tl	he las	st 6 m	nonth	is?		
	*Official lab results (most recent) mu	st be attached.	For continuati					st 6 m	nonth	IS?		
	*Official lab results (most recent) mu		For continuati					st 6 m	nonth	ıs?		
	*Official lab results (most recent) mu	st be attached.	For continuati	on of therap	y, labs ar	e requii		st 6 m	nonth	IS?		
	*Official lab results (most recent) mu	st be attached. yskinesia (TD) DISCUS:	For continuati	on of therap	y, labs ar months	e requii	red.	st 6 m	nonth	IS?		
	*Official lab results (most recent) mu □Yes No Date: _ Has an assessment for Tardive D AIMs: Yes No	st be attached. yskinesia (TD) DISCUS: must be attache	For continuati	n the last 6	y, labs ar months Da	e requii ? .te:	red.				 	
	*Official lab results (most recent) mu □Yes No Date: Has an assessment for Tardive D AIMs: □Yes □ No *Official Form or notation (most recent)	st be attached. yskinesia (TD) DISCUS: must be attache	For continuati	n the last 6	y, labs ar months Da hs TD	e requii ? ite:) Scree	red. n: q_		m	nonths		
	*Official lab results (most recent) mu	st be attached. yskinesia (TD) DISCUS: must be attache CMP	For continuati	n the last 6	y, labs ar months Da hs TD	e requii ? ite:) Scree	red. n: q_		m	nonths		
	*Official lab results (most recent) mu YesNoDate: Has an assessment for Tardive D AIMs:YesNo *Official Form or notation (most recent) Monitoring Plan: RTC: Labs:CBCProlactin Next Appointment Date:	st be attached. yskinesia (TD) DISCUS: must be attache 	For continuati	n the last 6	y, labs ar months Da hs TD , specify:	e requii ? tte: Scree	r ed. n: q_		m	nonths		
-	*Official lab results (most recent) mu \[Yes \[No Date: Has an assessment for Tardive D AIMs: \[Yes \[No *Official Form or notation (most recent) Monitoring Plan: RTC: Labs: \[CBC \[Prolactin Next Appointment Date: scriber's Signature:	st be attached. yskinesia (TD) DISCUS: must be attache CMP	For continuati	n the last 6	y, labs ar months Da hs TD	e requii ? ite:) Scree : Date	r ed. n: q		m	onths		
	*Official lab results (most recent) mu YesNoDate: Has an assessment for Tardive D AIMs:YesNo *Official Form or notation (most recent) Monitoring Plan: RTC: Labs:CBCProlactin Next Appointment Date:	st be attached. yskinesia (TD) DISCUS: must be attache CMP	For continuati	n the last 6	y, labs ar months Da hs TD , specify: s and rec	e requii ? ite:) Scree : Date	r ed. n: q		m	onths		
	*Official lab results (most recent) mu \[Yes \[No Date: Has an assessment for Tardive D AIMs: \[Yes \[No *Official Form or notation (most recent) Monitoring Plan: RTC: Labs: \[CBC \[Prolactin Next Appointment Date: Scriber's Signature: QUIRED FOR REVIEW: All copies of m ies of related labs. The provider must	st be attached. yskinesia (TD) DISCUS: must be attache CMP	For continuati	n the last 6	y, labs ar months Da hs TD , specify: s and rec years.	e requir ? te:) Scree : Date cent cha	n: q	tes), a	m	nonths e mos		

immediately and arrange for the return or destruction of these documents. Distribution, reproduction or any

other use of this transmission by any party other than the intended recipient is strictly prohibited



Aetna Better Health® of Florida (MEDICAID)

Antipsychotic (6 to < 18 Years of Age)

Maximum Length of Approval = 180 Days

Note: Form must be completed in full.

FDA-approved agents and doses are considered most appropriate.

	FDA Information for 6–17 Age Group
Age Range	Dosing Instructions
	Aripiprazole
Pediatric age	Initial dose: 2 mg/day
10–17	Recommended dose: 10 mg/day
	Maximum dose: 30 mg/day
Pediatric age	Initial dose: 2 mg/day
13–17	Recommended dose: 10 mg/day
	Maximum dose: 30 mg/day
Pediatric age	Initial dose: 2 mg/day
6–17	Recommended dose: 5–10 mg/day
	Maximum dose: 15 mg/day
	Lurasidone
Pediatric age	Initial dose: 20 mg/day
10-17	Recommended dose: 20-80 mg/day
	Maximum dose: 80 mg/day
Pediatric age	Initial dose: 40 mg/day
10-17	Recommended dose: 40-80 mg/day
	Maximum dose: 80 mg/day
	Olanzapine
Pediatric age	Oral Formulation
13–17	Initial dose: 2.5–5 mg/day
	Target dose: 10 mg/day
Pediatric age	Initial dose: 2.5–5 mg/day
13–17	Target dose: 10 mg/day
	Paliperidone
Pediatric age	Weight < 51kg: Initial Dose (3 mg/day)
12–17	Recommended Dose (3–6 mg/day)
	Maximum Dose (6 mg/day)
	Weight ≥ 51kg: Initial Dose (3 mg/day)
	Recommended Dose (3–12 mg/day)
	Maximum Dose (12 mg/day) Risperidone
Pediatric age	Initial dose: 0.5 mg/day
•	Titration: 0.5–1 mg/day
10 17	Recommended dose: 2.5 mg/day
	Recommended doge. 2.0 mg/day
	Pediatric age 10–17 Pediatric age 13–17 Pediatric age 6–17 Pediatric age 10-17 Pediatric age 10-17 Pediatric age 13–17 Pediatric age 13–17 Pediatric age

Confidentiality Notice: The documents accompanying this transmission contain confidential health information that is legally privileged. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender (via return fax) immediately and arrange for the return or destruction of these documents. Distribution, reproduction or any other use of this transmission by any party other than the intended recipient is strictly prohibited.



Aetna Better Health® of Florida (MEDICAID)

Antipsychotic (6 to < 18 Years of Age)

Maximum Length of Approval = 180 Days

Note: Form must be completed in full.

FDA Information for 6–17 Age Group								
Medication and Approved Use	Age Range	Dosing Instructions						
Irritability associated	Pediatric age	Initial dose: 0.25 mg/day (< 20 kg); 0.5 mg/day (≥ 20 kg)						
with Autism	5–16	Titration: 0.25–0.5 mg at > or = 2 weeks						
		Recommended dose: 0.5 mg/day (< 20 kg; 1 mg/day (≥ 20 kg)						
		Effective dose range: 0.5–3 mg/day						
Schizophrenia	Pediatric age	Initial dose: 0.5 mg/day						
	13–17	Titration: 0.5–1 mg/day						
		Target dose: 3 mg/day						
		Effective dose range: 1–6 mg/day						
		Quetiapine						
Bipolar I Disorder	Pediatric age	Information provided is for the immediate release table formulation						
(mania)	10–17	Day 1: 25 mg twice a day						
		Day 2: Twice daily dosing totaling 100 mg						
		Day 3: Twice daily dosing totaling 200 mg						
		Day 4: Twice daily dosing totaling 300 mg						
		Day 5: Twice daily dosing totaling 400 mg						
		Further adjustments should be in increments no greater than 100 mg/per day						
		within the recommended dose range of 400–600 mg/per day. Based on response and tolerability, may be administered three times daily.						
Schizophrenia	Pediatric age	Information provided is for the immediate release tablet formulation						
	12–17	Day 1: 25 mg twice daily						
		Day 2: Twice daily dosing totaling 100 mg						
		Day 3: Twice daily dosing totaling 200 mg						
		Day 4: Twice daily dosing totaling 300 mg						
		Day 5: Twice daily dosing totaling 400 mg						
		Recommend dose range: 400–800 mg/day						
		Further adjustments should be in increments no greater than 100 mg/per day within the recommended dose range of 400–800 mg/per day. Based on response and tolerability, may be administered three times daily.						

Helpful Links:

- Access the following information at http://floridabhcenter.org/index.html:
 - Antispychotic High Dosing Table for Children and Adolescents
 - AIMS/DISCUS forms
 - Florida Medicaid Best Practice Psychotherapeutic Medication Guidelines for Children and Adolescents
- The Centers for Disease Control and Prevention (CDC) BMI Calculator for Children and Teens: <u>https://www.cdc.gov/healthyweight/bmi/calculator.html</u>

Confidentiality Notice: The documents accompanying this transmission contain confidential health information that is legally privileged. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender (via return fax) immediately and arrange for the return or destruction of these documents. Distribution, reproduction or any other use of this transmission by any party other than the intended recipient is strictly prohibited.