AETNA BE	TTER HEALTH®		* ae	etna [®]
Coverage	Policy/Guideline			
Name:	Adbry (tralokinuma	ab-ldrm)	Page:	1 of 5
Effective D	Date: 4/1/2023		Last Review Date:	4/2023
A musline	□Illinois	□Florida	□Virginia	a
Applies to:	☐New Jersey	⊠Maryland	□Michig	an
	⊠Pennsylvania Kids	⊠Florida Kids	□Texas	

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Adbry under the patient's prescription drug benefit.

Description:

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indication

Indicated for the treatment of moderate-to-severe atopic dermatitis in adult patients whose disease is not adequately controlled with topical prescription therapies or when those therapies are not advisable. Adbry can be used with or without topical corticosteroids.

All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Adbry

Policy/Guideline:

Documentation for all indications:

The patient is unable to take Dupixent and Rinvoq, where indicated, for the given diagnosis due to a trial and inadequate treatment response or intolerance, or a contraindication. Documentation is required for approval.

Documentation:

Submission of the following information is necessary to initiate the prior authorization review:

- A. For initial requests:
 - 1. Member's chart notes or medical records showing affected area(s) and body surface area (where applicable).
 - 2. Member's chart notes, medical record documentation, or claims history of prerequisite therapies including response to therapy. If prerequisite therapies are not advisable, documentation of why therapy is not advisable for the member.
- B. For continuation requests: Documentation (e.g., chart notes) that the member has experienced a positive clinical response to therapy as evidenced by low disease activity or improvement in signs or symptoms of atopic dermatitis.

	TTER HEALTH® Policy/Guideline		* a	etna™
Name: Adbry (tralokinumab-ldrm)		Page:	2 of 5	
Effective Date: 4/1/2023			Last Review Da	te: 4/2023
Applies	□Illinois	□Florida	□Virg	inia
Applies to:	☐New Jersey	⊠Maryland	□Micl	higan
	⊠Pennsylvania Kids	⊠Florida Kids	□Texa	as

Prescriber Specialties:

This medication must be prescribed by or in consultation with a dermatologist or allergist/immunologist.

Criteria for Initial Approval:

Atopic dermatitis

Authorization of 4 months may be granted for treatment of moderate-to-severe atopic dermatitis in members 18 years of age or older when all of the following criteria are met:

- A. Affected body surface is greater than or equal to 10% body surface area OR crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
- B. Member meets one of the following:
 - 1. Member has had an inadequate treatment response with one of the following in the past year:
 - i. A medium potency to super-high potency topical corticosteroid (see Appendix)
 - ii. A topical calcineurin inhibitor
 - 2. The use of medium potency to super-high potency topical corticosteroid and topical calcineurin inhibitor are not advisable for the member (e.g., due to contraindications, prior intolerances).

Criteria for Continuation of Therapy:

Atopic dermatitis

Authorization of 12 months may be granted for members 18 years of age or older (including new members) who are using the requested medication for moderate-to-severe atopic dermatitis when the member has achieved or maintained a positive clinical response as evidenced by low disease activity (i.e., clear or almost clear skin), or improvement in signs and symptoms of atopic dermatitis (e.g., redness, itching, oozing/crusting).

Note: Member cannot use Adbry concomitantly with any other biologic drug or targeted synthetic drug.

Approval Duration and Quantity Restrictions:

Approval:

Initial: 4 monthsRenewal: 12 months

Quantity Level Limit:

			♥ae	etna™
	TTER HEALTH®			
Coverage	Policy/Guideline			
Name: Adbry (tralokinumab-ldrm)		ab-ldrm)	Page:	3 of 5
Effective Date: 4/1/2023			Last Review Date:	4/2023
Amaliaa	□Illinois	□Florida	□Virginia	
Applies to:	☐New Jersey	⊠Maryland	□Michiga	ın
	⊠Pennsylvania Kids	⊠Florida Kids	□Texas	

Loading Dose: Four 150 mg syringes for the first 14 days

Standard Limit: 4 syringes per 28 days

Adult Dosing: Initial dose of 600 mg (four 150 mg injections), followed by 300 mg (two 150 mg injections) administered every other week

12-17 Years Old Dosing: Initial dose of 300 mg (two 150 mg injections), followed by 150 (one 150 mg injection) every other week.

Appendix:

Table. Relative potency of select topical corticosteroid products

Potency	Drug	Dosage form	Strength
I. Super- high	Augmented betamethasone dipropionate	Ointment, Lotion, Gel	0.05%
potency (group 1)	Clobetasol propionate	Cream, Gel, Ointment, Solution, Cream (emollient), Lotion, Shampoo, Foam, Spray	0.05%
	Fluocinonide	Cream	0.1%
ı	Flurandrenolide	Tape	4 mcg/cm ²
	Halobetasol propionate	Cream, Lotion, Ointment, Foam	0.05%
II. High potency (group 2)	Amcinonide	Ointment	0.1%
	Augmented betamethasone dipropionate	Cream	0.05%
	Betamethasone dipropionate Ointment		0.05%
	Clobetasol propionate Cream		0.025%
	Desoximetasone	Cream, Ointment, Spray	0.25%
		Gel	0.05%
	Diflorasone diacetate Ointment, Cream (emollien		0.05%
	Fluocinonide	Cream, Ointment, Gel, Solution	0.05%
	Halcinonide	Cream, Ointment	0.1%
	Halobetasol propionate	Lotion	0.01%
Potency	Drug	Dosage form	Strength
	Amcinonide	Cream, Lotion	0.1%
	Betamethasone dipropionate	Cream, hydrophilic emollient	0.05%

	TTER HEALTH® Policy/Guideline		*ae	etna"
Name:	Adbry (tralokinuma	ıb-ldrm)	Page:	4 of 5
Effective Date: 4/1/2023			Last Review Date:	4/2023
Applies	□Illinois	□Florida	□Virginia	
Applies to:	☐New Jersey	⊠Maryland	□Michiga	ın
	⊠Pennsylvania Kids	⊠Florida Kids	□Texas	

Potency	Drug	Dosage form	Strength
	Betamethasone valerate	Ointment	0.1%
III. High		Foam	0.12%
potency	Desoximetasone	Cream, Ointment	0.05%
(group 3)	Diflorasone diacetate	Cream	0.05%
	Fluocinonide	Cream, aqueous emollient	0.05%
	Fluticasone propionate	Ointment	0.005%
	Mometasone furoate	Ointment	0.1%
	Triamcinolone acetonide	Cream, Ointment	0.5%
IV. Medium	Betamethasone dipropionate	Spray	0.05%
potency	Clocortolone pivalate	Cream	0.1%
(group 4)	Fluocinolone acetonide	Ointment	0.025%
	Flurandrenolide	Ointment	0.05%
	Hydrocortisone valerate	Ointment	0.2%
	Mometasone furoate	Cream, Lotion, Solution	0.1%
	Triamcinolone acetonide	Cream	0.1%
		Ointment	0.05% and 0.1%
		Aerosol Spray	0.2 mg per 2- second spray
V. Lower-	Betamethasone dipropionate	Lotion	0.05%
mid potency	Betamethasone valerate	Cream	0.1%
	Desonide	Ointment, Gel	0.05%
(group 5)	Fluocinolone acetonide	Cream	0.025%
	Flurandrenolide	Cream, Lotion	0.05%
	Fluticasone propionate	Cream, Lotion	
	Hydrocortisone butyrate	Cream, Lotion, Ointment, Solution	0.1%
	Hydrocortisone probutate	Cream	0.1%
	Hydrocortisone valerate	Cream	0.2%
	Prednicarbate	Cream (emollient), Ointment	0.1%
	Triamcinolone acetonide	Lotion	0.1%
		Ointment	0.025%
	Alclometasone dipropionate	Cream, Ointment	0.05%
	Betamethasone valerate	Lotion	0.1%

	TER HEALTH®		♥ae	etna [™]
Coverage F	Policy/Guideline			
Name: Adbry (tralokinumab-ldrm)		b-ldrm)	Page:	5 of 5
Effective Date: 4/1/2023			Last Review Date:	4/2023
Applies	□Illinois	□Florida	□Virginia	
Applies to:	□New Jersey	⊠Maryland	□Michiga	n
	⊠Pennsylvania Kids	⊠Florida Kids	□Texas	

Potency	Drug	Dosage form	Strength
VI. Low	Desonide	Cream, Lotion, Foam	0.05%
potency	Fluocinolone acetonide	Cream, Solution, Shampoo, Oil	0.01%
(group 6)	Triamcinolone acetonide	Cream, lotion	0.025%
	Hydrocortisone (base, greater than or	Cream, Ointment, Solution	2.5%
VII. Least potent (group 7)	equal to 2%)	Lotion	2%
	, ,	Cream, Ointment, Gel, Lotion, Spray, Solution	1%
		Cream, Ointment	0.5%
	Hydrocortisone acetate	Cream	2.5%
		Lotion	2%
		Cream	1%

References:

- 1. Adbry [package insert]. Madison, NJ: LEO Pharma Inc.; July 2022.
- 2. Eichenfield LF, Tom WL, Chamlin SL, et. al. Guidelines of care for the management of atopic dermatitis: Section 1. Diagnosis and Assessment of Atopic Dermatitis. *J Am Acad Dermatol*. 2014;70:338-351.
- 3. Topical Corticosteroids. *Drug Facts and Comparisons*. Facts & Comparisons [database online]. St. Louis, MO: Wolters Kluwer Health Inc; December 1, 2021. Accessed November 7, 2022.