



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name:	Eltrombopag olamine & Alvaiz (eltrombopag choline)	Page:	1 of 6
Effective Date:	7/3/2025	Last Review Date:	6/2025
Applies to:	<input type="checkbox"/> Illinois <input checked="" type="checkbox"/> Pennsylvania Kids <input checked="" type="checkbox"/> Virginia	<input checked="" type="checkbox"/> New Jersey <input type="checkbox"/> Michigan	<input checked="" type="checkbox"/> Florida Kids <input checked="" type="checkbox"/> Maryland

**Intent:**

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for eltrombopag olamine and Alvaiz under the patient's prescription drug benefit.

**Description:**

**A. FDA-Approved Indications**

- Treatment of thrombocytopenia in adult and pediatric patients 1 year and older with persistent or chronic immune thrombocytopenia (ITP) who have had an insufficient response to corticosteroids, immunoglobulins, or splenectomy.
- Treatment of thrombocytopenia in patients with chronic hepatitis C to allow the initiation and maintenance of interferon-based therapy.
- First-line treatment of severe aplastic anemia in adult and pediatric patients 2 years and older in combination with standard immunosuppressive therapy
- Treatment of patients with severe aplastic anemia who have had an insufficient response to immunosuppressive therapy.

**B. Alvaiz is indicated for:**

- Treatment of thrombocytopenia in adult and pediatric patients 6 years and older with persistent or chronic ITP who have had an insufficient response to corticosteroids, immunoglobulins, or splenectomy.
- Treatment of thrombocytopenia in adult patients with chronic hepatitis C to allow the initiation and maintenance of interferon-based therapy.
- Treatment of adult patients with severe aplastic anemia who have had an insufficient response to immunosuppressive therapy.

**C. Compendial Uses**

1. MYH9-related disease with thrombocytopenia (eltrombopag olamine only)
2. Myelodysplastic syndromes (MDS) (eltrombopag olamine only)
3. Thrombocytopenia post-hematopoietic cell transplant

D. All other indications are considered experimental/investigational and not medically necessary.

**Applicable Drug List:**

Eltrombopag olamine  
Alvaiz



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name:	Eltrombopag olamine & Alvaiz (eltrombopag choline)	Page:	2 of 6
Effective Date:	7/3/2025	Last Review Date:	6/2025
Applies to:	<input type="checkbox"/> Illinois <input checked="" type="checkbox"/> Pennsylvania Kids <input checked="" type="checkbox"/> Virginia	<input checked="" type="checkbox"/> New Jersey <input type="checkbox"/> Michigan	<input checked="" type="checkbox"/> Florida Kids <input checked="" type="checkbox"/> Maryland

**Policy/Guideline:**

The patient unable to take the preferred formulary alternative for the given diagnosis due to a trial and inadequate treatment response or intolerance, or a contraindication.  
Documentation is required for approval.

**Documentation**

Submission of the following information is necessary to initiate the prior authorization review:

- A. Persistent or chronic immune thrombocytopenia (ITP):
  - 1. For initial requests: pretreatment platelet count
  - 2. For continuation requests: current platelet count
- B. Aplastic anemia continuation of therapy: current platelet count

**Exclusions**

Coverage will not be provided when the requested drug will be used concomitantly with other thrombopoietin receptor agonists (e.g., Nplate, Doptelet, Mulpleta) or with spleen tyrosine kinase inhibitors (e.g., Tavalisse).

**Prescriber Specialties:**

This medication must be prescribed by or in consultation with EITHER of the following:

- Persistent or chronic immune thrombocytopenia (ITP), aplastic anemia, MYH9-related disease with thrombocytopenia, myelodysplastic syndromes, and thrombocytopenia post-hematopoietic cell transplant: hematologist or oncologist
- Thrombocytopenia with hepatitis C: hematologist or a prescriber specializing in infectious disease, gastroenterology, hepatology, or transplant

**Criteria for Initial Approval:**

**A. Persistent or chronic immune thrombocytopenia (ITP)**

Authorization of 6 months may be granted for treatment of persistent or chronic ITP when BOTH of the following criteria are met:

- 1. Member has had an inadequate response or intolerance to prior therapy with corticosteroids, immunoglobulins, or splenectomy.
- 2. Member has an untransfused platelet count at any point prior to the initiation of the requested medication of EITHER of the following:
  - a) Less than  $30 \times 10^9/L$
  - b)  $30 \times 10^9/L$  to  $50 \times 10^9/L$  with symptomatic bleeding (e.g., significant mucous membrane bleeding, gastrointestinal bleeding, or trauma) or risk factors for bleeding (see Appendix).



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name:	Eltrombopag olamine & Alvaiz (eltrombopag choline)	Page:	3 of 6
Effective Date:	7/3/2025	Last Review Date:	6/2025
Applies to:	<input type="checkbox"/> Illinois <input checked="" type="checkbox"/> Pennsylvania Kids <input checked="" type="checkbox"/> Virginia	<input checked="" type="checkbox"/> New Jersey <input type="checkbox"/> Michigan	<input checked="" type="checkbox"/> Florida Kids <input checked="" type="checkbox"/> Maryland

**B. Thrombocytopenia associated with chronic hepatitis C**

Authorization of 12 months may be granted to members who are prescribed the requested drug for the initiation and maintenance of interferon-based therapy for the treatment of thrombocytopenia associated with chronic hepatitis C.

**C. Severe Aplastic anemia**

1. Eltrombopag olamine

- i. Authorization of 6 months may be granted for first-line treatment of severe aplastic anemia when the requested drug will be used in combination with standard immunosuppressive therapy (e.g., horse antithymocyte globulin [h-ATG] and cyclosporine).
- ii. Authorization of 6 months may be granted for treatment of severe aplastic anemia in members who have had an insufficient response to immunosuppressive therapy.

2. Alvaiz

Authorization of 6 months may be granted for treatment of aplastic anemia in members who have had an insufficient response to immunosuppressive therapy.

**D. MYH9-related disease with thrombocytopenia (eltrombopag olamine only)**

Authorization of 12 months may be granted to members with thrombocytopenia associated with MYH9-related disease.

**E. Myelodysplastic syndromes (eltrombopag olamine only)**

Authorization of 12 months may be granted for treatment of myelodysplastic syndromes (MDS).

**F. Thrombocytopenia post-hematopoietic cell transplant**

Authorization of 6 months may be granted for treatment of prolonged thrombocytopenia in members who are post-allogeneic transplant and have poor graft function.

**Criteria for Continuation of Therapy**

**A. Persistent or chronic ITP**

1. Authorization of 3 months may be granted to members with current platelet count less than  $50 \times 10^9/L$  for whom the platelet count is not sufficient to prevent clinically important bleeding and who have not received a maximal dose of the requested drug for at least 4 weeks.
2. Authorization of 12 months may be granted to members with current platelet count less than  $50 \times 10^9/L$  for whom the current platelet count is sufficient to prevent clinically important bleeding.
3. Authorization of 12 months may be granted to members with current platelet count of  $50 \times 10^9/L$  to  $200 \times 10^9/L$ .
4. Authorization of 12 months may be granted to members with current platelet count greater than  $200 \times 10^9/L$  to less than or equal to  $400 \times 10^9/L$  for whom dosing for the



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name:	Eltrombopag olamine & Alvaiz (eltrombopag choline)	Page:	4 of 6
Effective Date:	7/3/2025	Last Review Date:	6/2025
Applies to:	<input type="checkbox"/> Illinois <input checked="" type="checkbox"/> Pennsylvania Kids <input checked="" type="checkbox"/> Virginia	<input checked="" type="checkbox"/> New Jersey <input type="checkbox"/> Michigan	<input checked="" type="checkbox"/> Florida Kids <input checked="" type="checkbox"/> Maryland

requested drug will be adjusted to achieve a platelet count sufficient to avoid clinically important bleeding.

**B. Thrombocytopenia associated with chronic hepatitis C**

Authorization of 6 months may be granted to members who are continuing to receive interferon-based therapy.

**C. Severe Aplastic anemia**

1. Authorization of up to 16 weeks total may be granted to members with current platelet count less than  $50 \times 10^9/L$  who have not received appropriately titrated therapy with the requested drug for at least 16 weeks.
2. Authorization of 12 months total may be granted to members with current platelet count less than  $50 \times 10^9/L$  who are transfusion independent.
3. Authorization of 12 months may be granted to members with current platelet count of  $50 \times 10^9/L$  to  $200 \times 10^9/L$ .
4. Authorization of 12 months may be granted to members with current platelet count greater than  $200 \times 10^9/L$  to less than or equal to  $400 \times 10^9/L$  for whom dosing for the requested drug will be adjusted to achieve and maintain an appropriate target platelet count.

**D. MYH9-related disease with thrombocytopenia (eltrombopag olamine only)**

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

**E. Myelodysplastic syndromes (eltrombopag olamine only) and chemotherapy-induced thrombocytopenia (CIT)**

Authorization of 12 months may be granted for continued treatment of myelodysplastic syndromes or thrombocytopenia post-hematopoietic cell transplant in members who experience benefit from therapy (e.g., increased platelet counts, decreased bleeding events, reduced need for platelet transfusions).

**Appendix**

**Examples of risk factors for bleeding (not all inclusive)**

- Undergoing a medical or dental procedure where blood loss is anticipated
- Comorbidity (e.g., peptic ulcer disease, hypertension)
- Mandated anticoagulation therapy
- Profession (e.g., construction worker) or lifestyle (e.g., plays contact sports) that predisposes member to trauma

**Quantity Restrictions**

**Quantity Level Limit:**



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Eltrombopag olamine & Alvaiz  
(eltrombopag choline)

Page: 5 of 6

Effective Date: 7/3/2025

Last Review Date: 6/2025

Applies to: ☐ Illinois ☒ New Jersey ☒ Florida Kids  
☒ Pennsylvania Kids ☐ Michigan ☒ Maryland  
☒ Virginia

Medication	Standard Limit	FDA-recommended dosing
Eltrombopag olamine 12.5 mg tablets	60 per 30 days	<p>Persistent or chronic immune thrombocytopenia (ITP): Initiate at 50 mg once daily for most adult and pediatric patients 6 years and older and at 25 mg once daily for pediatric patients aged 1 to 5 years. Dose reductions are needed for patients of East-/Southeast-Asian ancestry or those with hepatic impairment. Adjust to maintain platelet count greater than or equal to <math>50 \times 10^9/L</math>. Do not exceed 75 mg per day.</p> <p>Chronic Hep C-associated thrombocytopenia: Initiate at 25 mg once daily. Adjust to achieve target platelet count required to initiate antiviral therapy. Do not exceed a daily dose of 100 mg.</p> <p>First-line severe aplastic anemia: Initiate once daily at 2.5mg/kg (in pediatric patients aged 2 to 5 years old), 75 mg (pediatric patients aged 6 to 11 years old), or 150mg for patients aged 12 years and older concurrently with standard immunosuppressive therapy. Reduce initial dose in patients of East-/Southeast-Asian ancestry or those with hepatic impairment. Modify dosage for toxicity or elevated platelet counts.</p> <p>Refractory severe aplastic anemia: Initiate at 50mg once daily for most patients. Reduce initial dose in patients with hepatic impairment or patients of East-/Southeast-Asian ancestry. Adjust to maintain platelet count greater than or equal to <math>50 \times 10^9/L</math>. Do not exceed 150 mg per day.</p>
Eltrombopag olamine 25 mg tablets	90 per 30 days	
Eltrombopag olamine 12.5 mg oral susp pkts	120 packets per 30 days	
Eltrombopag olamine 25 mg oral susp pkts	180 packets per 30 days	
Eltrombopag olamine 50 mg tablets	90 per 30 days	
Eltrombopag olamine 75 mg tablets	60 per 30 days	
Alvaiz 9 mg tablets	60 per 30 days	<p>Persistent or chronic ITP: Initiate at 36mg once daily for most adult and pediatric patients 6 years and older. Dose reductions are needed for patients of East-/Southeast-Asian ancestry or those with hepatic impairment. Adjust to maintain platelet count greater than or equal to <math>50 \times 10^9/L</math>. Do not exceed 54mg per day.</p> <p>Chronic hepatitis C-associated thrombocytopenia: Initiate at 18mg once daily. Adjust to achieve target</p>
Alvaiz 18 mg tablets	90 per 30 days	
Alvaiz 36 mg tablets	90 per 30 days	



**AETNA BETTER HEALTH®**  
Coverage Policy/Guideline

Name:	Eltrombopag olamine & Alvaiz (eltrombopag choline)	Page:	6 of 6
Effective Date:	7/3/2025	Last Review Date:	6/2025
Applies to:	<input type="checkbox"/> Illinois <input checked="" type="checkbox"/> New Jersey <input checked="" type="checkbox"/> Florida Kids <input checked="" type="checkbox"/> Pennsylvania Kids <input type="checkbox"/> Michigan <input checked="" type="checkbox"/> Maryland <input checked="" type="checkbox"/> Virginia		

Medication	Standard Limit	FDA-recommended dosing
Alvaiz 54 mg tablets	60 per 30 days	platelet count required to initiate antiviral therapy. Do not exceed a daily dose of 72mg.  Refractory severe aplastic anemia: Initiate at 36mg once daily for most patients. Reduce initial dose in patients with hepatic impairment or patients of East-/Southeast-Asian ancestry. Adjust to maintain platelet count greater than or equal to $50 \times 10^9/L$ . Do not exceed 108 mg per day.

**References:**

1. Promacta [package insert]. East Hanover, NJ: Novartis Pharmaceuticals Corporation; March 2023.
2. Alvaiz [package insert]. Parsippany, NJ: Teva Pharmaceuticals; May 2024.
3. Pecci A, Gresele P, Klersy C, et al. Eltrombopag for the treatment of the inherited thrombocytopenia deriving from MYH9 mutations. *Blood*. 2010;116(26):5832-7.
4. The NCCN Drugs & Biologics Compendium® © 2024 National Comprehensive Cancer Network, Inc. <https://www.nccn.org>. Accessed June 10, 2024.
5. The NCCN Clinical Practice Guidelines in Oncology® Myelodysplastic Syndrome (Version 2.2024). © 2024 National Comprehensive Cancer Network, Inc. <https://www.nccn.org>. Accessed June 10, 2024.
6. Nuenert C, Terrel DR, Arnold DM, et al. American Society of Hematology 2019 guidelines for immune thrombocytopenia. *Blood Adv*. 2019;3(23):3829–3866.
7. Provan D, Arnold DM, Bussel JB, et al. Updated international consensus report on the investigation and management of primary immune thrombocytopenia. *Blood Adv*. 2019;3(22): 3780–3817.
8. Provan D, Stasi R, Newland AC, et al. International consensus report on the investigation and management of primary immune thrombocytopenia. *Blood*. 2010;115(2):168-186.
9. Rodeghiero F, Stasi R, Gernsheimer T, et al. Standardization of terminology, definitions and outcome criteria in immune thrombocytopenic purpura of adults and children: report from an international working group. *Blood*. 2009;113(11):2386-2393.
10. Olnes MJ, Scheinberg P, Calvo KR, et al. Eltrombopag and improved hematopoiesis in refractory aplastic anemia. *N Engl J Med*. 2012;367(1):11-19.
11. Townsley DM, Scheinberg P, Winkler T, et al. Eltrombopag added to standard immunosuppression for aplastic anemia. *N Engl J Med*. 2017;376:1540-1550.
12. Kulasekararaj A, Cavenagh J, Dokal I, et al. Guidelines for the diagnosis and management of adult aplastic anaemia: A British Society for Haematology guideline. *Br J Haematol*. 2024;204(3):784-804.