



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Filsuvez

Page: 1 of 2

Effective Date: 3/26/2024

Last Review Date: 02/08/2024

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> New Jersey
	<input checked="" type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Florida Kids	<input checked="" type="checkbox"/> Pennsylvania Kids
	<input type="checkbox"/> Michigan	<input checked="" type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

### Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Filsuvez under the patient's prescription drug benefit.

### Description:

#### FDA-Approved Indication

Indicated for the treatment of wounds associated with dystrophic and junctional epidermolysis bullosa (EB) in adults and pediatric patients 6 months of age and older.

All other indications are considered experimental/investigational and not medically necessary.

### Applicable Drug List:

Filsuvez

### Policy/Guideline:

Submission of the following information is necessary to initiate the prior authorization review:

- Medical records documenting clinical manifestations of disease.
- Laboratory test results supporting diagnosis.

### Prescriber Specialties

This medication must be prescribed by or in consultation with a dermatologist or wound care specialist.

### Criteria for Initial Approval

#### **Epidermolysis Bullosa (EB)**

Authorization may be granted for treatment of wounds in members with dystrophic epidermolysis bullosa (DEB) and junctional epidermolysis bullosa (JEB) when ALL the following criteria are met:

- Member is 6 months of age or older.
- Member has clinical manifestations of disease (e.g., extensive skin blistering, skin erosions, scarring).
- Member has laboratory test results confirming diagnosis (i.e., genetic testing, immunofluorescence mapping [IFM], or transmission electron microscopy [TEM]).
- Filsuvez will not be administered to wound(s) that are currently healed.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Filsuvez

Page: 2 of 2

Effective Date: 3/26/2024

Last Review Date: 02/08/2024

Applies to:	<input checked="" type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> New Jersey
	<input checked="" type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Florida Kids	<input checked="" type="checkbox"/> Pennsylvania Kids
	<input type="checkbox"/> Michigan	<input checked="" type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

### Continuation of Therapy

#### Epidermolysis Bullosa (EB)

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

#### Approval Duration and Quantity Restrictions:

**Approval:** 12 months

**Quantity Level Limit:** Reference Formulary for drug specific quantity level limits

#### References:

1. Filsuvez [package insert]. Wahlstedt, Germany: Lichtenheldt GmbH; December 2023.
2. Has C, Liu L, Bolling MC, et al. Clinical practice guidelines for laboratory diagnosis of epidermolysis bullosa. *Br J Dermatol.* 2020; 182: 574-592.