



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Kevzara

Page: 1 of 8

Effective Date: 7/31/2026

Last Review Date: 6/17/2026

Applies to: ☒ Illinois

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Kevzara under the patient's prescription drug benefit.

Description:

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-approved Indications

- Adult patients with moderately to severely active rheumatoid arthritis (RA) who have had an inadequate response or intolerance to one or more disease-modifying antirheumatic drugs (DMARDs).
- Adult patients with polymyalgia rheumatica (PMR) who have had an inadequate response to corticosteroids or who cannot tolerate corticosteroid taper.
- Patients with active polyarticular juvenile idiopathic arthritis (pJIA) who weigh 63 kg or greater.

Compendial Uses

- Immune checkpoint inhibitor-related toxicity - inflammatory arthritis
- Giant cell arteritis (GCA)

All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Non-Preferred Agent: Kevzara

Policy/Guideline:

Documentation for ALL indications:

The patient is unable to take THREE preferred products, where indicated, for the given diagnosis due to a trial and inadequate treatment response or intolerance, or a contraindication. Documentation is required for approval.

Documentation

Submission of the following information is necessary to initiate the prior authorization review:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Kevzara

Page: 2 of 8

Effective Date: 7/31/2026

Last Review Date: 6/17/2026

Applies to: ☒ Illinois

Rheumatoid arthritis

Initial requests

- Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
- Laboratory results, chart notes, or medical record documentation of biomarker testing (i.e., rheumatoid factor [RF], anti-cyclic citrullinated peptide [anti-CCP], and C-reactive protein [CRP] and/or erythrocyte sedimentation rate [ESR]) (if applicable).

Continuation requests

Chart notes or medical record documentation supporting positive clinical response.

Polymyalgia rheumatica and immune checkpoint inhibitor-related toxicity

Initial requests

Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

Continuation requests

Chart notes or medical record documentation supporting positive clinical response.

Polyarticular juvenile idiopathic arthritis

Initial requests

Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy.

Continuation requests

Chart notes or medical record documentation supporting positive clinical response.

Giant cell arteritis (GCA)

For continuation requests

Chart notes or medical record documentation supporting positive clinical response.

Prescriber Specialties

This medication must be prescribed by or in consultation with ONE of the following:

- Rheumatoid arthritis, polymyalgia rheumatica, polyarticular juvenile idiopathic arthritis, and giant cell arteritis: rheumatologist
- Immune checkpoint inhibitor-related toxicity: oncologist, hematologist, or rheumatologist

Coverage Criteria

Rheumatoid arthritis (RA)

Authorization of 12 months may be granted for adult members who have previously



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Kevzara

Page: 3 of 8

Effective Date: 7/31/2026

Last Review Date: 6/17/2026

Applies to: ☒ Illinois

received a biologic or targeted synthetic drug indicated for moderately to severely active rheumatoid arthritis (RA) within the past 120 days.

Authorization of 12 months may be granted for adult members for treatment of moderately to severely active RA when BOTH of the following criteria are met:

- Member meets EITHER of the following:
 - Member has been tested for either of the following biomarkers and the test was positive:
 - Rheumatoid factor (RF)
 - Anti-cyclic citrullinated peptide (anti-CCP)
 - Member has been tested for ALL of the following biomarkers:
 - RF
 - Anti-CCP
 - C-reactive protein (CRP) and/or erythrocyte sedimentation rate (ESR)
- Member meets ONE of the following:
 - Member has failed to achieve a low disease activity after a 3-month trial of methotrexate (MTX) monotherapy at a maximum titrated dose of at least 15 mg per week and meets ANY of the following conditions:
 - Member has had a documented inadequate response to MTX in combination with at least one other conventional synthetic drug (i.e., hydroxychloroquine and/or sulfasalazine) after a 3-month trial at a maximum tolerated dose(s).
 - Member has experienced a documented intolerable adverse event to hydroxychloroquine or sulfasalazine.
 - Member has a documented contraindication to hydroxychloroquine (see Appendix A) and sulfasalazine (e.g., porphyria, intestinal or urinary obstruction).
 - Member has moderate to high disease activity.
 - Member was unable to tolerate a 3-month trial of MTX monotherapy at a maximum titrated dose of at least 15 mg per week and meets ANY of the following conditions:
 - Member has had a documented inadequate response to MTX in combination with at least one other conventional synthetic drug (i.e., hydroxychloroquine and/or sulfasalazine) after a 3-month trial at a maximum tolerated dose(s).
 - Member has stopped taking MTX and has had a documented inadequate response to another conventional synthetic drug (i.e., leflunomide, hydroxychloroquine, and/or sulfasalazine) alone or in combination after a 3-month trial at a maximum tolerated dose(s).
 - Member has experienced a documented intolerable adverse event to leflunomide, hydroxychloroquine, or sulfasalazine.



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Kevzara

Page: 4 of 8

Effective Date: 7/31/2026

Last Review Date: 6/17/2026

Applies to: ☒ Illinois

- Member has a documented contraindication to leflunomide, hydroxychloroquine (see Appendix A), and sulfasalazine (e.g., porphyria, intestinal or urinary obstruction).
- Member has moderate to high disease activity.
- Member has experienced a documented intolerable adverse event or has a documented contraindication to MTX (see Appendix A), discontinues MTX, and meets ANY of the following conditions:
 - Member has had a documented inadequate response to another conventional synthetic drug (i.e., leflunomide, hydroxychloroquine, and/or sulfasalazine) alone or in combination after a 3-month trial at a maximum tolerated dose(s).
 - Member has experienced a documented intolerable adverse event to leflunomide, hydroxychloroquine, or sulfasalazine.
 - Member has a documented contraindication to leflunomide, hydroxychloroquine (see Appendix A), and sulfasalazine (e.g., porphyria, intestinal or urinary obstruction).
 - Member has moderate to high disease activity.

Polymyalgia rheumatica (PMR)

Authorization of 12 months may be granted for adult members for treatment of polymyalgia rheumatica when ANY of the following criteria is met:

- Member has experienced an inadequate response to systemic corticosteroids.
- Member has experienced a disease flare during a taper with systemic corticosteroids.
- Member has experienced an inadequate response to methotrexate.
- Member has experienced an intolerance or contraindication to both systemic corticosteroids and methotrexate (see Appendix A).

Polyarticular juvenile idiopathic arthritis (pJIA)

Authorization of 12 months may be granted for members weighing 63 kg and greater who have previously received a biologic or targeted synthetic drug indicated for active polyarticular juvenile idiopathic arthritis.

Authorization of 12 months may be granted for members weighing 63 kg and greater for treatment of active polyarticular juvenile idiopathic arthritis when ANY of the following criteria is met:

- Member has had an inadequate response to methotrexate or another conventional synthetic drug (e.g., leflunomide, sulfasalazine, hydroxychloroquine) administered at an adequate dose and duration.
- Member has had an inadequate response to a trial of scheduled non-steroidal anti-inflammatory drugs (NSAIDs) and/or intra-articular glucocorticoids (e.g., triamcinolone hexacetonide) and ONE of the following risk factors for poor outcome:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Kevzara

Page: 5 of 8

Effective Date: 7/31/2026

Last Review Date: 6/17/2026

Applies to: ☒ Illinois

- Involvement of ankle, wrist, hip, sacroiliac joint, and/or temporomandibular joint (TMJ)
 - Presence of erosive disease or enthesitis
 - Delay in diagnosis
 - Elevated levels of inflammation markers
 - Symmetric disease
2. Member has risk factors for disease severity and potentially a more refractory disease course (see Appendix B) and the member also meets ONE of the following:
- High-risk joints are involved (e.g., cervical spine, wrist, or hip)
 - High disease activity
 - Is judged to be at high risk for disabling joint disease

Immune checkpoint inhibitor-related toxicity

Authorization of 12 months may be granted for treatment of immune checkpoint inhibitor-related toxicity when then member has moderate or severe immunotherapy-related inflammatory arthritis and meets EITHER of the following:

- Member has had an inadequate response to corticosteroids or a conventional synthetic drug (e.g., methotrexate, sulfasalazine, leflunomide, hydroxychloroquine).
- Member has an intolerance or contraindication to corticosteroids and a conventional synthetic drug (e.g., methotrexate, sulfasalazine, leflunomide, hydroxychloroquine).

Giant cell arteritis (GCA)

Authorization of 12 months may be granted for treatment of giant cell arteritis when the member's diagnosis was confirmed by EITHER of the following:

- Temporal artery biopsy or cross-sectional imaging
- Acute-phase reactant elevation (i.e., high erythrocyte sedimentation rate [ESR] and/or high serum C-reactive protein [CRP])

Continuation of Therapy

Rheumatoid arthritis (RA)

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for moderately to severely active RA and who achieve or maintain a positive clinical response as evidenced by disease activity improvement of at least 20% from baseline in tender joint count, swollen joint count, pain, or disability.

Polymyalgia rheumatica (PMR)

Authorization of 12 months may be granted for all adult members (including new members) who are using the requested medication for PMR and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in ANY of the following from baseline:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Kevzara

Page: 6 of 8

Effective Date: 7/31/2026

Last Review Date: 6/17/2026

Applies to: ☒ Illinois

- Morning stiffness
- Hip or shoulder pain
- Hip or shoulder range of motion
- C-reactive protein (CRP) and/or erythrocyte sedimentation rate (ESR)

Polyarticular juvenile idiopathic arthritis

Authorization of 12 months may be granted for all members (including new members) weighing 63 kg and greater who are using the requested medication for active polyarticular juvenile idiopathic arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in ANY of the following from baseline:

- Number of joints with active arthritis (e.g., swelling, pain, limitation of motion)
- Number of joints with limitation of movement
- Functional ability

Immune checkpoint inhibitor-related toxicity

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for immunotherapy-related inflammatory arthritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition.

Giant cell arteritis (GCA)

Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for giant cell arteritis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in ANY of the following from baseline:

- Headaches
- Scalp tenderness
- Tenderness and/or thickening of superficial temporal arteries
- Constitutional symptoms (e.g., weight loss, fever, fatigue, night sweats)
- Jaw and/or tongue claudication
- Acute visual symptoms (e.g., amaurosis fugax, acute visual loss, diplopia)
- Symptoms of polymyalgia rheumatica (e.g., shoulder and/or hip girdle pain)
- Limb claudication
- Acute-phase reactants (i.e., erythrocyte sedimentation rate [ESR] or serum C-reactive protein [CRP])

Other

For all indications: Member has had a documented negative tuberculosis (TB) test (which can include a tuberculosis skin test [TST] or an interferon-release assay [IGRA]) within 12

	
AETNA BETTER HEALTH® Coverage Policy/Guideline	
Name: Kevzara	Page: 7 of 8
Effective Date: 7/31/2026	Last Review Date: 6/17/2026
Applies to: ☒ Illinois	

months of initiating therapy for persons who are naïve to biologic drugs or targeted synthetic drugs associated with an increased risk of TB.

If the screening testing for TB is positive, there must be further testing to confirm there is no active disease (e.g., chest x-ray). Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug.

Dosage and Administration

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

Appendix

Appendix A: Examples of Clinical Reasons to Avoid Methotrexate, Hydroxychloroquine, or Leflunomide

- Clinical diagnosis of alcohol use disorder, alcoholic liver disease, or other chronic liver disease
- Drug interaction
- Risk of treatment-related toxicity
- Pregnancy or currently planning pregnancy
- Breastfeeding
- Significant comorbidity prohibits use of systemic agents (e.g., liver or kidney disease, blood dyscrasias, uncontrolled hypertension)
- Hypersensitivity
- History of intolerance or adverse event

Appendix B: Risk Factors for Articular Juvenile Idiopathic Arthritis

- Positive rheumatoid factor
- Positive anti-cyclic citrullinated peptide antibodies
- Pre-existing joint damage

Approval Duration and Quantity Restrictions:

Initial Approval: 12 months

Renewal Approval: 12 months

Quantity Level Limit:

- 150 mg/1.14 mL single-dose prefilled syringe: 1 pack (2 x 150 mg syringe) per 4 weeks
- 150 mg/1.14 mL single-dose prefilled pen: 1 pack (2 x 150 mg pen) per 4 weeks
- 200 mg/1.14 mL single-dose prefilled syringe: 1 pack (2 x 200 mg syringe) per 4 weeks



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Kevzara

Page:

8 of 8

Effective Date: 7/31/2026

Last Review Date: 6/17/2026

Applies to: ☒ Illinois

- 200 mg/1.14 mL single-dose prefilled pen: 1 pack (2 x 200 mg pen) per 4 weeks

References:

1. Kevzara [package insert]. Morristown, NJ: Sanofi-aventis, U.S. LLC /Regeneron Pharmaceuticals, Inc.; May 2025.
2. Genovese MC, Fleischmann R, Kivitz AJ, et al. Sarilumab plus methotrexate in patients with active rheumatoid arthritis and inadequate response to methotrexate: results of a phase III study. *Arthritis Rheumatol.* June 2015;67(6):1424-37.
3. Strand V, Reaney M, Chen C, et al. Sarilumab improves patient-reported outcomes in rheumatoid arthritis patients with inadequate response/intolerance to tumour necrosis factor inhibitors. *RMD Open.* 2017; 3:e000416. doi: 10.1136/rmdopen-2016-000416.
4. Testing for TB Infection. Centers for Disease Control and Prevention. Retrieved on June 17, 2025 from: <https://www.cdc.gov/tb/testing/>.
5. Smolen JS, Landewé RBM, Bergstra SA, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2022 update. *Ann Rheum Dis.* 2023;82:3-18.
6. Aletaha D, Neogi T, Silman, et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum.* 2010;62(9):2569-81.
7. Smolen JS, Aletaha D. Assessment of rheumatoid arthritis activity in clinical trials and clinical practice. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. Available with subscription. URL: www.uptodate.com. Accessed March 19, 2021.
8. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Care Res.* 2021;0:1-16.
9. Dasgupta B, Cimmino MA, Kremers HM, et al. 2012 provisional classification criteria for polymyalgia rheumatica: a European League Against Rheumatism/American College of Rheumatology collaborative initiative. *Arthritis Rheum.* 2012 Apr;64(4):943-54.
10. Menter A, Gelfand JM, Connor C, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management of psoriasis with systemic nonbiologic therapies. *J Am Acad Dermatol.* 2020;82(6):1445-1486.
11. Ringold S, Angeles-Han S, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Treatment of Juvenile Idiopathic Arthritis: Therapeutic Approaches for Non-Systemic Polyarthritis, Sacroiliitis, and Enthesitis. American College of Rheumatology. 2019;1-18.
12. Onel KB, Horton DB, Lovell DJ, et al. 2021 American College of Rheumatology guideline for the treatment of juvenile idiopathic arthritis: therapeutic approaches for oligoarthritis, temporomandibular joint arthritis, and systemic juvenile idiopathic arthritis. *Arthritis Rheumatol.* 2022;74(4):553-569.
13. The NCCN Drugs & Biologics Compendium. © 2025 National Comprehensive Cancer Network, Inc. <https://www.nccn.org>. Accessed June 18, 2025.