



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Yesintek

Page: 1 of 9

Effective Date: 5/23/2025

Last Review Date: 4/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

**Intent:**

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Yesintek under the patient's prescription drug benefit.

**Description:**

**A. FDA-Approved Indications**

1. Moderate to severe plaque psoriasis (PsO) in patients 6 years or older who are candidates for phototherapy or systemic therapy
2. Active psoriatic arthritis (PsA) in patients 6 years or older
3. Moderately to severely active Crohn's disease (CD) in adults
4. Moderately to severely active ulcerative colitis (UC) in adults

**B. Compendial Uses**

Immune checkpoint inhibitor-related toxicity

All other indications are considered experimental/investigational and not medically necessary.

**Applicable Drug List:**

**Preferred:**

Yesintek (ustekinumab-kfce)

**Policy/Guideline:**

**Documentation:**

**A. **Plaque psoriasis****

1. Initial requests:
  - i. Chart notes or medical record documentation of affected area(s) and body surface area (BSA) affected.
  - ii. Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
2. Continuation requests: Chart notes or medical record documentation of decreased body surface area (BSA) affected and/or improvement in signs and symptoms.

**B. **Psoriatic arthritis****



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Yesintek

Page: 2 of 9

Effective Date: 5/23/2025

Last Review Date: 4/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

1. Initial requests: Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
2. Continuation requests: Chart notes or medical record documentation supporting positive clinical response.

**C. Crohn's disease and Ulcerative Colitis**

Continuation requests: Chart notes or medical record documentation supporting positive clinical response to therapy or remission.

**D. Immune checkpoint inhibitor-related toxicity**

Chart notes, medical record documentation, or claims history supporting previous medications tried, including response to therapy or intolerance to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

**Prescriber Specialty:**

The medication must be prescribed by or in consultation with one of the following:

- A. Plaque psoriasis: dermatologist
- B. Psoriatic arthritis: rheumatologist or dermatologist
- C. Crohn's disease and ulcerative colitis: gastroenterologist
- D. Immune checkpoint inhibitor-related toxicity: gastroenterologist, hematologist or oncologist

**Criteria for Initial Approval:**

**A. Plaque psoriasis (PsO)**

1. Authorization of 12 months may be granted for members 6 years of age or older who previously received a biologic or targeted synthetic drug indicated for the treatment of moderate to severe plaque psoriasis.
2. Authorization of 12 months may be granted for treatment of moderate to severe plaque psoriasis in members 6 years of age or older when any of the following criteria is met:
  - i. Crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
  - ii. At least 10% of the body surface area (BSA) is affected
  - iii. At least 3% of body surface area (BSA) is affected and the member meets any of the following criteria:



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Yesintek

Page: 3 of 9

Effective Date: 5/23/2025

Last Review Date: 4/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- a. Member has had an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin.
- b. Member has a clinical reason to avoid pharmacologic treatment with methotrexate, cyclosporine, and acitretin (see Appendix).

**B. Psoriatic arthritis (PsA)**

1. Authorization of 12 months may be granted for members 6 years of age or older who have previously received a biologic or targeted synthetic drug indicated for active psoriatic arthritis.
2. Authorization of 12 months may be granted for members 6 years of age or older for treatment of active psoriatic arthritis when either of the following criteria is met:
  - i. Member has mild to moderate disease and meets one of the following criteria:
    - a. Member has had an inadequate response to methotrexate, leflunomide, or another conventional synthetic drug (e.g., sulfasalazine) administered at an adequate dose and duration.
    - b. Member has an intolerance or contraindication to methotrexate or leflunomide (see Appendix), or another conventional synthetic drug (e.g., sulfasalazine).
    - c. Member has enthesitis
  - ii. Member has severe disease.

**C. Crohn's disease (CD)**

Authorization of 12 months may be granted for treatment of moderately to severely active Crohn's disease.

**D. Ulcerative colitis (UC)**

Authorization of 12 months may be granted for treatment of moderately to severely active ulcerative colitis.

**E. Immune checkpoint inhibitor-related toxicity**

Authorization of 6 months may be granted for the treatment of immune checkpoint inhibitor-related diarrhea or colitis when the member has experienced an inadequate response, intolerance, or contraindication to infliximab or vedolizumab.

**Continuation of Therapy:**

**A. Plaque psoriasis (PsO)**



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Yesintek

Page: 4 of 9

Effective Date: 5/23/2025

Last Review Date: 4/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

Authorization of 12 months may be granted for all members 6 years of age or older (including new members) who are using the requested medication for moderate to severe plaque psoriasis and who achieve or maintain positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when any of the following is met:

1. Reduction in body surface area (BSA) affected from baseline
2. Improvement in signs and symptoms from baseline (e.g., itching, redness, flaking, scaling, burning, cracking, pain)

**B. Psoriatic arthritis (PsA)**

Authorization of 12 months may be granted for all members 6 years of age or older (including new members) who are using the requested medication for psoriatic arthritis and who achieve or maintain positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:

1. Number of swollen joints
2. Number of tender joints
3. Dactylitis
4. Enthesitis
5. Skin and/or nail involvement
6. Functional status
7. C-reactive protein (CRP)

**C. Crohn's Disease (CD)**

1. Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active or fistulizing Crohn's disease and who achieve or maintain remission.
2. Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active or fistulizing Crohn's disease and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
  - i. Abdominal pain or tenderness
  - ii. Diarrhea
  - iii. Body weight
  - iv. Abdominal mass



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Yesintek

Page: 5 of 9

Effective Date: 5/23/2025

Last Review Date: 4/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- v. Hematocrit
- vi. Endoscopic appearance of the mucosa
- vii. Improvement on a disease activity scoring tool (e.g., Crohn's Disease Activity Index [CDAI] score)

#### D. Ulcerative colitis

1. Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active ulcerative colitis and who achieve or maintain remission.
2. Authorization of 12 months may be granted for all members (including new members) who are using the requested medication for moderately to severely active ulcerative colitis and who achieve or maintain a positive clinical response as evidenced by low disease activity or improvement in signs and symptoms of the condition when there is improvement in any of the following from baseline:
  - i. Stool frequency
  - ii. Rectal bleeding
  - iii. Urgency of defecation
  - iv. C-reactive protein (CRP)
  - v. Fecal calprotectin (FC)
  - vi. Endoscopic appearance of the mucosa
  - vii. Improvement on a disease activity scoring tool (e.g., Ulcerative Colitis Endoscopic Index of Severity [UCEIS], Mayo score)

#### E. Immune checkpoint inhibitor-related toxicity

All members (including new members) requesting authorization for continuation of therapy must meet all initial authorization criteria.

##### Other Criteria:

For all indications: Member has had a documented negative tuberculosis (TB) test (which can include a tuberculosis skin test [TST] or an interferon-release assay [IGRA]) within 12 months of initiating therapy for persons who are naïve to biologic drugs or targeted synthetic drugs associated with an increased risk of TB.

If the screening testing for TB is positive, there must be further testing to confirm there is no active disease (e.g., chest x-ray). Do not administer the requested medication to members with active TB infection. If there is latent disease, TB treatment must be started before initiation of the requested medication.



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Yesintek

Page: 6 of 9

Effective Date: 5/23/2025

Last Review Date: 4/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

For all indications: Member cannot use the requested medication concomitantly with any other biologic drug or targeted synthetic drug for the same indication.

**Dosage and Administration:**

Approvals may be subject to dosing limits in accordance with FDA-approved labeling, accepted compendia, and/or evidence-based practice guidelines.

**Appendix**

**Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Cyclosporine, Acitretin, or Leflunomide**

1. Clinical diagnosis of alcohol use disorder, alcoholic liver disease, or other chronic liver disease
2. Drug interaction
3. Risk of treatment-related toxicity
4. Pregnancy or currently planning pregnancy
5. Breastfeeding
6. Significant comorbidity prohibits use of systemic agents (e.g., liver or kidney disease, blood dyscrasias, uncontrolled hypertension)
7. Hypersensitivity
8. History of intolerance or adverse event

**Approval Duration and Quantity Restrictions:**

**Approval:**

Initial Approval:

- 6 months for Immune checkpoint inhibitor-related toxicity
- 12 months for all other indications

Renewal Approval:

- 6 months for Immune checkpoint inhibitor-related toxicity
- 12 months for all other indications

**Quantity Level Limit:**

- Yesintek (ustekinumab-kfce) 130 mg/26 mL single-dose vial
  - 4 vials (1 dose)
- Yesintek (ustekinumab-kfce) subcutaneous injection 45 mg/0.5 mL single-dose vial/prefilled syringe
  - 1 vial/syringe per 84 days
  - Exception limit: 2 vials/syringes per 28 days
- Yesintek (ustekinumab-kfce) subcutaneous injection 90 mg/mL prefilled syringe



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Yesintek

Page: 7 of 9

Effective Date: 5/23/2025

Last Review Date: 4/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- 1 syringe per 56 days
- Exception limit: 2 syringes per 28 days

\*Coverage up to the exception limits may be provided with prior authorization

**FDA-Recommended Dosing:**

130 mg/26 mL single-dose vial

**Crohn's Disease (CD), Ulcerative Colitis (UC) Intravenous Induction**

- ≤ 55 kg: 260 mg (2 vials)
- 55 kg to 85 kg: 390 mg (3 vials)
- > 85 kg: 520 mg (4 vials)

45 mg/0.5 mL single-dose vial/prefilled syringe, 90 mg/mL prefilled syringe

**Plaque Psoriasis (PsO)**

**Adult**

- ≤ 100 kg: 45 mg initially and 4 weeks later, followed by 45 mg every 12 weeks
- > 100 kg: 90 mg initially and 4 weeks later, followed by 90 mg every 12 weeks

**Pediatric (6-17 years old)**

Weight based dosing at the initial dose, 4 weeks later, and then every 12 weeks thereafter

- < 60 kg: 0.75 mg/kg
- 60 kg to 100 kg: 45 mg
- > 100 kg: 90 mg

**Psoriatic Arthritis (PsA)**

**Adult**

- 45 mg initially and 4 weeks later, followed by 45 mg every 12 weeks
- > 100 kg with co-existent PsO: 90 mg initially and 4 weeks later, followed by 90 mg every 12 weeks

**Pediatric (6-17 years old)**

Weight based dosing at the initial dose, 4 weeks later, and then every 12 weeks thereafter



AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Yesintek

Page: 8 of 9

Effective Date: 5/23/2025

Last Review Date: 4/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- < 60 kg: 0.75 mg/kg
- 60 kg or more: 45 mg
- > 100 kg with co-existent PsO: 90 mg

### **Crohn's Disease (CD), Ulcerative Colitis (UC)**

#### **Maintenance Dose**

- 90 mg 8 weeks after the initial intravenous dose, then every 8 weeks thereafter

#### **Inadequate Response After Induction (4 weeks after first maintenance dose)**

- 90 mg every 4 weeks

\*Coverage up to the exception limits may be provided with prior authorization

#### **References:**

1. Stelara [package insert]. Horsham, PA: Janssen Biotech, Inc.; November 2024.
2. Imuldosa [package insert]. Raleigh, NC: Accord BioPharma Inc.; October 2024.
3. Otulfi [package insert]. Lake Zurich, IL: Fresenius Kabi USA, LLC; September 2024.
4. Pyzchiva [package insert]. Princeton, NJ: Sandoz Inc.; December 2024.
5. Selarsdi [package insert]. Leesburg, VA: Alvotect USA Inc.; October 2024.
6. Steqeyma [package insert]. Jersey City, NJ: Celltrion USA, Inc.; December 2024.
7. ustekinumab-aekn [package insert]. Leesburg, VA: Alvotect USA Inc.; October 2024.
8. ustekinumab-ttwe [package insert]. Grand Cayman, Cayman Islands: Quallent Pharmaceuticals Health LLC; March 2025.
9. Wezlana [package insert]. Thousand Oaks, CA: Amgen Inc.; December 2024.
10. Yesintek [package insert]. Cambridge, MA: Biocon Biologics Inc.; November 2024.
11. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. J Am Acad Dermatol. 2011;65(1):137-174.
12. Gossec L, Baraliakos X, Kerschbaumer A, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies: 2019 update. Ann Rheum Dis. 2020;79(6):700-712.
13. Gladman DD, Antoni C, Mease P, et al. Psoriatic arthritis: epidemiology, clinical features, course, and outcome. Ann Rheum Dis 2005;64(Suppl II):ii14-ii17.
14. Talley NJ, Abreu MT, Achkar J, et al. An evidence-based systematic review on medical therapies for inflammatory bowel disease. Am J Gastroenterol. 2011;106(Suppl 1):S2-S25.
15. Lichtenstein GR, Loftus Jr EV, Isaacs KI, et al. ACG Clinical Guideline: Management of Crohn's Disease in Adults. Am J Gastroenterol. 2018;113:481-517.
16. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. J Am Acad Dermatol. 2019;80(4):1029-1072.





AETNA BETTER HEALTH®  
Coverage Policy/Guideline

Name: Yesintek

Page: 9 of 9

Effective Date: 5/23/2025

Last Review Date: 4/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input checked="" type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

17. Testing for TB Infection. Centers for Disease Control and Prevention. Retrieved on January 21, 2025 from: <https://www.cdc.gov/tb/testing/index.html>.
18. Rubin DT, Ananthakrishnan AN, et al. 2019 ACG Clinical Guideline: Ulcerative Colitis in Adults. Am J Gastroenterol. 2019;114:384-413.
19. Feuerstein JD, Isaacs KL, Schneider Y, et al. AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. Gastroenterology. 2020;158:1450.
20. Singh JA, Guyatt G, Ogdie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. Arthritis Rheumatol. 2019;71(1):5-32. doi:10.1002/art.40726.
21. Menter A, Cordero KM, Davis DM, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis in pediatric patients. J Am Acad Dermatol. 2020;82(1):161-201.
22. Menter A, Gelfand JM, Connor C, et al. Joint AAD-NPF guidelines of care for the management of psoriasis with systemic nonbiologic therapies. J Am Acad Dermatol. 2020;82(6): 1445-86.
23. Feuerstein JD, Ho EY, Shmidt E, et al. AGA Clinical Practice Guidelines on the Medical Management of Moderate to Severe Luminal and Perianal Fistulizing Crohn's Disease. Gastroenterology. 2021; 160: 2496- 2508.
24. The NCCN Drugs & Biologics Compendium® © 2025 National Comprehensive Cancer Network, Inc. Available at: <http://www.nccn.org>. Accessed January 14, 2025.
25. Coates LC, Soriano ER, Corp N, et al. Group for Research and Assessment of Psoriasis and Psoriatic Arthritis (GRAPPA): updated treatment recommendations for psoriatic arthritis 2021. Nat Rev Rheumatol. 2022;18(8):465-479.
26. NCCN Clinical Practice Guidelines in Oncology® (NCCN Guidelines®). Management of Immunotherapy-related Toxicities. Version 1.2025. Available at: [www.nccn.org](http://www.nccn.org). Accessed January 21, 2025.