



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Epoprostenol Page: 1 of 4

Effective Date: 2/28/2025 Last Review Date: 1/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input checked="" type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for epoprostenol under the patient’s prescription drug benefit.

Description:

Indications

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications¹⁻³

Flolan/Veletri/epoprostenol is indicated for the treatment of pulmonary arterial hypertension (PAH) (World Health Organization [WHO] Group I) to improve exercise capacity. Studies establishing effectiveness included predominantly patients with New York Heart Association (NYHA) Functional Class III-IV symptoms and etiologies of idiopathic or heritable PAH or PAH associated with connective tissue diseases. All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Epoprostenol
Flolan
Veletri

Policy/Guideline:

Prescriber Specialty:

This medication must be prescribed by or in consultation with a pulmonologist or cardiologist.

Coverage Criteria

Pulmonary Arterial Hypertension (PAH)¹⁻⁷

12-month authorization may be granted for treatment of PAH when ALL of the following criteria are met:

- Member has PAH defined as WHO Group 1 class of pulmonary hypertension (refer to Appendix).
- PAH was confirmed by either of the following criteria:



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Epoprostenol Page: 2 of 4

Effective Date: 2/28/2025 Last Review Date: 1/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input checked="" type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- Pretreatment right heart catheterization with all of the following results.
 - Mean pulmonary arterial pressure (mPAP) > 20 mmHg
 - Pulmonary capillary wedge pressure (PCWP) ≤ 15 mmHg
 - Pulmonary vascular resistance (PVR) > 2 Wood units. For pediatric members, pulmonary vascular resistance index (PVRI) > 3 Wood units x m² is also acceptable.
- For infants less than one year of age, PAH was confirmed by Doppler echocardiogram if right heart catheterization cannot be performed.

Continuation of Therapy

12-month authorization may be granted for members with an indication listed in the coverage criteria section who are currently receiving the requested medication through a paid pharmacy or medical benefit, and who are experiencing benefit from therapy as evidenced by disease stability or disease improvement.

Appendix

WHO Classification of Pulmonary Hypertension (PH)⁵

Note: Patients with heritable PAH or PAH associated with drugs and toxins might be long-term responders to calcium channel blockers.

Group 1: Pulmonary Arterial Hypertension (PAH)

- Idiopathic
 - Long-term responders to calcium channel blockers
- Heritable
- Associated with drugs and toxins
- Associated with:
 - Connective tissue disease
 - Human immunodeficiency virus (HIV) infection
 - Portal hypertension
 - Congenital heart disease
 - Schistosomiasis
- PAH with features of venous/capillary (pulmonary veno-occlusive disease [PVOD]/pulmonary capillary hemangiomatosis [PCH]) involvement
- Persistent PH of the newborn

Group 2: PH associated with Left Heart Disease

- Heart failure:
 - With preserved ejection fraction



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Epoprostenol Page: 3 of 4

Effective Date: 2/28/2025 Last Review Date: 1/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input checked="" type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- With reduced or mildly reduced ejection fraction
- Cardiomyopathies with specific etiologies (i.e., hypertrophic, amyloid, Fabry disease, and Chagas disease)
- Valvular heart disease:
 - Aortic valve disease
 - Mitral valve disease
 - Mixed valvular disease
- Congenital/acquired cardiovascular conditions leading to post-capillary PH

Group 3: PH associated with Lung Diseases and/or Hypoxia

- Chronic obstructive pulmonary disease (COPD) and/or emphysema
- Interstitial lung disease
- Combined pulmonary fibrosis and emphysema
- Other parenchymal lung diseases (i.e., parenchymal lung diseases not included in Group 5)
- Nonparenchymal restrictive diseases:
 - Hypoventilation syndromes
 - Pneumonectomy
- Hypoxia without lung disease (e.g., high altitude)
- Developmental lung diseases

Group 4: PH associated with Pulmonary Artery Obstructions

- Chronic thromboembolic PH
- Other pulmonary artery obstructions:
 - Sarcomas (high- or intermediate-grade or angiosarcoma)
 - Other malignant tumors (e.g., renal carcinoma, uterine carcinoma, germ-cell tumors of the testis)
 - Non-malignant tumors (e.g., uterine leiomyoma)
 - Arteritis without connective tissue disease
 - Congenital pulmonary artery stenoses
 - Hydatidosis

Group 5: PH with Unclear and/or Multifactorial Mechanisms

- Hematological disorders, including inherited and acquired chronic hemolytic anemia and chronic myeloproliferative disorders
- Systemic disorders: Sarcoidosis, pulmonary Langerhans cell histiocytosis, and neurofibromatosis type 1
- Metabolic disorders, including glycogen storage diseases and Gaucher disease



AETNA BETTER HEALTH®
Coverage Policy/Guideline

Name: Epoprostenol Page: 4 of 4

Effective Date: 2/28/2025 Last Review Date: 1/2025

Applies to:	<input type="checkbox"/> Illinois	<input type="checkbox"/> Florida	<input type="checkbox"/> Florida Kids
	<input type="checkbox"/> New Jersey	<input type="checkbox"/> Maryland	<input type="checkbox"/> Michigan
	<input type="checkbox"/> Pennsylvania Kids	<input checked="" type="checkbox"/> Virginia	<input type="checkbox"/> Kentucky PRMD

- Chronic renal failure with or without hemodialysis
- Pulmonary tumor thrombotic microangiopathy
- Fibrosing mediastinitis
- Complex congenital heart disease

Approval Duration and Quantity Restrictions:

Approval: 12 months

References:

1. Flolan [package insert]. Research Triangle Park, NC: GlaxoSmithKline; August 2021.
2. Veletri [package insert]. Titusville, NJ: Actelion Pharmaceuticals US, Inc.; July 2022.
3. Epoprostenol injection [package insert]. Cranbury, NJ: Sun Pharmaceutical Industries, Inc.; October 2024.
4. Simonneau G, Montani D, Celermajer DS, et al. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur Respir J.* 2019;53(1):1801913. doi:10.1183/13993003.01913-2018
5. Kovacs G, Bartolome S, Denton CP, et al. Definition, classification and diagnosis of pulmonary hypertension. *Eur Respir J.* 2024;64(4):2401324. doi: 10.1183/13993003.01324-2024
6. Chin KM, Gaine SP, Gerges C, et al. Treatment algorithm for pulmonary arterial hypertension. *Eur Respir J.* 2024;64(4):2401325. doi: 10.1183/13993003.01325-2024
7. Ivy D, Rosenzweig EB, Abman SH, et al. Embracing the challenges of neonatal and paediatric pulmonary hypertension. *Eur Respir J.* 2024;64(4):2401345. doi: 10.1183/13993003.01345-2024