Osteoporosis Management in Women Who Had a Fracture (OMW)

2025

Women ages 67–85 MEDICARE

Measure definition

Women who had a fracture **and** either a bone mineral density (BMD) test **or** received a prescription to treat osteoporosis within 6 months of fracture (excludes fractures to the phalanges (fingers/toes), face and skull)

Intake period: From July 1 of year prior to measurement year to June 30 of measurement year

The intake period is used to capture the first fracture

Remove episode dates where any of the following are met:

- Members who had a BMD test during 24 months prior to fracture
- Members who had a claim or encounter for osteoporosis therapy within 365 days prior to fracture
- Members who received a dispensed prescription or had an active prescription to treat osteoporosis within 365 days prior to episode date



Medical record requirements

- Member legal name and date of birth
- Provider/practice identifier
- Provider Business Group (PBG) name and number
- Date of service (DOS)
- Applicable lab/test results and date collected



Commonly used claim codes*

• BMD test: 77080, 77081, 77085

Osteoporosis medication therapy:

- Injection, denosumab, 1 mg: J0897
- Injection, ibandronate sodium, 1 mg: J1740
- Injection, teriparatide, 10 mcg: J3110
- Injection, romosozumab-aqqg, 1 mg: J3111
- Injection, zoledronic acid, 1 mg: J3489

Medical record submission methods may not be applicable to all plan types. For more details, you can reach out to your HEDIS plan representative.



Required exclusions (Other exclusions may also apply)

- Members in hospice or using hospice services during measurement year
- Members who died anytime during measurement year
- Members who had a palliative care encounter or received palliative care anytime during intake period through end of measurement year
- Members ages 67 and older as of December 31 of measurement year who meet either of the following:
 - Enrolled in an Institutional SNP (I-SNP) anytime during the intake period through the end of the measurement year
 - Living long-term in an institution anytime during the intake period through end of measurement vear
- Members 67–80 years of age and older by the end of the measurement year with both frailty and advanced illness. Members must meet both frailty and advanced illness criteria to be excluded:
 - For frailty, at least 2 indications of frailty with different dates of service during the intake period through the end of measurement year
 - For advanced illness, either of the following during the measurement year or year prior:
 - Advanced illness on at least 2 different dates of service
 - Dispensed dementia medication
- Members ages 81 and older as of December 31 of measurement year with at least 2 indications of frailty during the intake period through the end of measurement year

Insights and recommendations

- Order a BMD test on all women with a diagnosis of a fracture or prescribe medication to prevent osteoporosis within 6 months.
- Schedule a follow-up to review BMD results and/or medication adherence.
- Ensure accurate coding of fractures.
- Let members know pharmacists can be a resource for questions about medications.
- Keep in mind that BMD testing in an inpatient setting satisfies this measure.

*FOR COMMONLY USED CODES: Not a comprehensive list of codes.

For measures that require claims data only, we cannot accept supplemental data sources, such as data feeds and medical record collection methods.

CPT® is a registered trademark of the American Medical Association. 2024 All rights reserved.

Aetna is the brand name used for products and services provided by one or more of the Aetna group of companies, including Aetna Life Insurance Company and its affiliates (Aetna).

The above information is not a complete list of services for this measure. For a complete list, please refer to the NCQA website at **NCQA.org**. HEDIS 2025 Volume 2: Technical Specifications for Health Plans by the National Committee for Quality Assurance (NCQA). HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA).

This message is for informational purposes only, is not medical advice and is not intended to be a substitute for proper medical care provided by a physician. Information is believed to be accurate as of the production date; however, it is subject to change. Refer **to Aetna.com** for more information about Aetna® plans.

