			⇔ aetna™		
AETNA BE	ETTER HEALTH®				
Coverage Policy/Guideline					
Name:	Egrifta		Page:	1 of 2	
Effective Date: 4/7/2024			Last Review Date:	4/2024	
Applies to:	⊠Illinois	□Florida	⊠Florida Kids		
	⊠New Jersey	⊠Maryland	⊠Michigan		
	⊠Pennsylvania Kids	⊠Virginia	□Arizona		

Intent:

The intent of this policy/guideline is to provide information to the prescribing practitioner outlining the coverage criteria for Egrifta under the patient's prescription drug benefit.

Description:

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indication

Egrifta SV is indicated for the reduction of excess abdominal fat in human immunodeficiency virus (HIV)-infected adult patients with lipodystrophy.

Limitations of Use:

- A. Long-term cardiovascular safety of Egrifta SV has not been established.
- B. Egrifta SV is not indicated for weight loss management as it has a weight neutral effect.
- C. There are no data to support improved compliance with anti-retroviral therapies in HIV-positive patients taking Egrifta SV.

All other indications are considered experimental/investigational and not medically necessary.

Applicable Drug List:

Non-Preferred: Egrifta

Policy/Guideline:

Prescriber Specialty:

This medication must be prescribed by or in consultation with an infectious disease specialist.

Criteria for Initial Approval:

Authorization of 6 months may be granted for reduction of excess abdominal fat in HIV-infected patients with lipodystrophy when the patient is currently receiving anti-retroviral therapy.

Continuation of Therapy:

			♦	etna [®]	
AETNA BE	TTER HEALTH®				
Coverage Policy/Guideline					
Name:	Egrifta		Page:	2 of 2	
Effective Date: 4/7/2024			Last Review Date:	4/2024	
Applies to:	⊠Illinois	□Florida	⊠Florida Kids		
	⊠New Jersey	⊠Maryland	⊠Michigan		
	⊠Pennsylvania Kids	⊠Virginia	□Arizona		

Authorization of 6 months may be granted for continued treatment in members requesting reauthorization for reduction of excess abdominal fat when all of the following criteria are met:

- A. The member has HIV infection and lipodystrophy
- B. The member is currently receiving anti-retroviral therapy
- C. The member has demonstrated a clear clinical improvement from baseline that is supported by waist circumference measurement or computed tomography (CT) scan

Note: Coverage will not be provided for weight loss.

Approval Duration and Quantity Restrictions:

Approval:

Initial Approval: 6 months Renewal Approval: 6 months

Quantity Level Limit: 30 vials per 30 days

Reference Formulary for drug specific quanitity level limits

References:

- 1. Egrifta SV [package insert]. Montreal, Québec: Theratechnologies, Inc.; October 2019.
- 2. Brown TT. Approach to the human immunodeficiency virus-infected patient with lipodystrophy. *J Clin Endocrinol Metab.* 2008;93(8):2937-2945.